



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 01:00 PM UTC

PDB ID : 2VQZ / pdb_00002vqz
Title : Structure of the cap-binding domain of influenza virus polymerase subunit PB2 with bound m7GTP
Authors : Guilligay, D.; Tarendeau, F.; Resa-Infante, P.; Coloma, R.; Crepin, T.; Sehr, P.; Lewis, J.; Ruigrok, R.W.H.; Ortin, J.; Hart, D.J.; Cusack, S.
Deposited on : 2008-03-21
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	166	 87% 10% ..
1	B	166	 87% 10% ..
1	D	166	 89% 10% ..
1	E	166	 84% 13% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	166	6% 78% 17% .. .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6920 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called POLYMERASE BASIC PROTEIN 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	164	Total	C	N	O	S	Se	0	2	0
			1303	815	238	238	2	10			
1	B	162	Total	C	N	O	S	Se	0	2	0
			1294	811	235	236	2	10			
1	D	165	Total	C	N	O	S	Se	0	2	0
			1321	827	242	240	2	10			
1	E	160	Total	C	N	O	S	Se	0	1	0
			1270	797	229	232	2	10			
1	F	160	Total	C	N	O	S	Se	0	5	0
			1295	813	235	234	1	12			

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	389	LYS	ARG	engineered mutation	UNP P31345
B	389	LYS	ARG	engineered mutation	UNP P31345
D	389	LYS	ARG	engineered mutation	UNP P31345
E	389	LYS	ARG	engineered mutation	UNP P31345
F	389	LYS	ARG	engineered mutation	UNP P31345

- Molecule 2 is 7N-METHYL-8-HYDROGUANOSINE-5'-TRIPHOSPHATE (CCD ID: MGT) (formula: C₁₁H₂₀N₅O₁₄P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total 33	C 11	N 5	O 14	P 3	0	0
2	B	1	Total 33	C 11	N 5	O 14	P 3	0	0
2	D	1	Total 33	C 11	N 5	O 14	P 3	0	0
2	E	1	Total 33	C 11	N 5	O 14	P 3	0	0
2	F	1	Total 33	C 11	N 5	O 14	P 3	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	65	Total O 65 65	0	0
3	B	66	Total O 66 66	0	0
3	D	58	Total O 58 58	0	0
3	E	51	Total O 51 51	0	0
3	F	32	Total O 32 32	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: POLYMERASE BASIC PROTEIN 2

Chain A: 



- Molecule 1: POLYMERASE BASIC PROTEIN 2

Chain B: 



- Molecule 1: POLYMERASE BASIC PROTEIN 2

Chain D: 




- Molecule 1: POLYMERASE BASIC PROTEIN 2

Chain E: 



- Molecule 1: POLYMERASE BASIC PROTEIN 2

Chain F: 



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	92.20Å 94.44Å 220.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.53 – 2.30 29.53 – 2.30	Depositor EDS
% Data completeness (in resolution range)	99.7 (29.53-2.30) 99.6 (29.53-2.30)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.39 (at 2.31Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.186 , 0.235 0.182 , 0.230	Depositor DCC
R_{free} test set	2165 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	32.1	Xtriage
Anisotropy	0.240	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 49.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.047 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6920	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.20% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MGT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.81	0/1310	0.91	0/1734
1	B	0.79	0/1301	0.90	0/1721
1	D	0.81	0/1329	0.92	0/1760
1	E	0.78	0/1276	0.90	0/1687
1	F	0.70	0/1303	0.91	3/1724 (0.2%)
All	All	0.78	0/6519	0.91	3/8626 (0.0%)

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	468	THR	CB-CA-C	6.00	117.34	108.87
1	F	429	ASN	CA-C-N	5.15	126.28	119.84
1	F	429	ASN	C-N-CA	5.15	126.28	119.84

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1303	0	1328	13	0
1	B	1294	0	1316	12	0
1	D	1321	0	1347	12	1

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	E	1270	0	1297	13	0
1	F	1295	0	1309	18	0
2	A	33	0	16	3	0
2	B	33	0	16	3	0
2	D	33	0	16	2	0
2	E	33	0	16	3	0
2	F	33	0	16	2	0
3	A	65	0	0	3	0
3	B	66	0	0	0	0
3	D	58	0	0	3	1
3	E	51	0	0	5	0
3	F	32	0	0	2	0
All	All	6920	0	6677	67	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

The worst 5 of 67 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:355:ARG:HD2	2:B:1483:MGT:O1G	1.66	0.95
1:D:355:ARG:HD2	2:D:1483:MGT:O1G	1.69	0.91
1:F:451:ILE:HG21	1:F:473[B]:MSE:HE3	1.58	0.83
1:B:421:VAL:HG13	1:B:425:ASN:HA	1.67	0.77
1:A:355:ARG:HD2	2:A:1483:MGT:O1G	1.87	0.75

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:458:MSE:SE	3:D:2027:HOH:O[4_555]	2.14	0.06

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	163/166 (98%)	159 (98%)	4 (2%)	0	100	100
1	B	160/166 (96%)	158 (99%)	2 (1%)	0	100	100
1	D	165/166 (99%)	161 (98%)	4 (2%)	0	100	100
1	E	157/166 (95%)	153 (98%)	4 (2%)	0	100	100
1	F	159/166 (96%)	156 (98%)	3 (2%)	0	100	100
All	All	804/830 (97%)	787 (98%)	17 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	144/135 (107%)	140 (97%)	4 (3%)	38	56
1	B	143/135 (106%)	138 (96%)	5 (4%)	32	48
1	D	146/135 (108%)	142 (97%)	4 (3%)	39	58
1	E	141/135 (104%)	137 (97%)	4 (3%)	38	56
1	F	143/135 (106%)	134 (94%)	9 (6%)	16	23
All	All	717/675 (106%)	691 (96%)	26 (4%)	32	47

5 of 26 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	391	GLU
1	F	341	GLU
1	F	457	VAL
1	E	422	ASN
1	F	342	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 8 such sidechains are listed below:

Mol	Chain	Res	Type
1	F	439	GLN
1	F	429	ASN
1	F	357	HIS
1	E	422	ASN
1	F	422	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MGT	A	1483	-	34,35,35	3.64	11 (32%)	45,56,56	2.15	12 (26%)
2	MGT	B	1483	-	34,35,35	3.47	11 (32%)	45,56,56	1.87	12 (26%)
2	MGT	E	1483	-	34,35,35	3.75	9 (26%)	45,56,56	1.82	10 (22%)
2	MGT	F	1483	-	34,35,35	3.64	10 (29%)	45,56,56	1.94	15 (33%)
2	MGT	D	1483	-	34,35,35	3.59	11 (32%)	45,56,56	2.26	15 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.
'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MGT	A	1483	-	-	1/22/50/50	0/3/3/3
2	MGT	B	1483	-	-	0/22/50/50	0/3/3/3
2	MGT	E	1483	-	-	2/22/50/50	0/3/3/3
2	MGT	F	1483	-	-	5/22/50/50	0/3/3/3
2	MGT	D	1483	-	-	7/22/50/50	0/3/3/3

The worst 5 of 52 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1483	MGT	C8-N9	-16.82	1.35	1.45
2	A	1483	MGT	C8-N9	-16.66	1.35	1.45
2	E	1483	MGT	C8-N9	-16.16	1.35	1.45
2	B	1483	MGT	C8-N9	-15.99	1.35	1.45
2	D	1483	MGT	C8-N9	-15.60	1.35	1.45

The worst 5 of 64 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	1483	MGT	C2-N3-C4	5.99	122.62	112.30
2	A	1483	MGT	C2-N3-C4	5.62	121.98	112.30
2	F	1483	MGT	C2-N3-C4	5.51	121.80	112.30
2	B	1483	MGT	C2-N3-C4	5.34	121.50	112.30
2	A	1483	MGT	C5-C6-N1	5.33	120.32	110.94

There are no chirality outliers.

5 of 15 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	D	1483	MGT	C5'-O5'-PA-O3A
2	D	1483	MGT	C5'-O5'-PA-O2A
2	F	1483	MGT	C5'-O5'-PA-O3A
2	D	1483	MGT	PB-O3A-PA-O1A
2	D	1483	MGT	PG-O3B-PB-O1B

There are no ring outliers.

5 monomers are involved in 13 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1483	MGT	3	0

Continued on next page...


Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1483	MGT	3	0
2	E	1483	MGT	3	0
2	F	1483	MGT	2	0
2	D	1483	MGT	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




Ligand MGT B 1483




Ligand MGT E 1483



Ligand MGT F 1483



Ligand MGT D 1483



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/166 (92%)	-0.35	0 100 100	12, 26, 43, 64	1 (0%)
1	B	152/166 (91%)	-0.20	2 (1%) 75 76	12, 29, 48, 83	2 (1%)
1	D	155/166 (93%)	-0.22	2 (1%) 75 76	14, 27, 59, 80	2 (1%)
1	E	150/166 (90%)	-0.18	2 (1%) 75 76	16, 31, 54, 70	1 (0%)
1	F	150/166 (90%)	0.24	10 (6%) 24 26	29, 47, 74, 100	1 (0%)
All	All	761/830 (91%)	-0.15	16 (2%) 63 65	12, 31, 65, 100	7 (0%)

The worst 5 of 16 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	453[A]	HIS	4.1
1	B	424	ALA	3.7
1	B	426	GLN	2.8
1	F	422	ASN	2.8
1	F	426	GLN	2.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,




median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MGT	F	1483	33/33	0.81	0.12	54,65,97,100	0
2	MGT	E	1483	33/33	0.90	0.10	30,43,68,70	0
2	MGT	D	1483	33/33	0.91	0.09	27,41,64,67	0
2	MGT	B	1483	33/33	0.94	0.07	23,38,52,53	0
2	MGT	A	1483	33/33	0.94	0.07	25,34,50,51	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around MGT F 1483:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGT E 1483:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGT D 1483:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around MGT B 1483:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MGT A 1483:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.