



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 18, 2026 – 01:11 PM UTC

PDB ID : 2VRJ / pdb_00002vrj
Title : Beta-glucosidase from *Thermotoga maritima* in complex with N-octyl-5- deoxy-6-oxa-N-(thio)carbamoylecystine
Authors : Aguilar, M.; Gloster, T.M.; Garcia-Moreno, M.I.; Ortiz Mellet, C.; Davies, G.J.; Llebaria, A.; Casas, J.; Egido-Gabas, M.; Garcia Fernandez, J.M.
Deposited on : 2008-04-09
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	468	
1	B	468	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	NCW	A	1446[A]	-	-	X	-
4	ACT	B	1448	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called BETA-GLUCOSIDASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	438	Total	C	N	O	S	0	22	0
			3720	2424	628	661	7			
1	B	443	Total	C	N	O	S	0	10	0
			3608	2342	595	664	7			

- Molecule 2 is (1S,2R,3S,4R,5R)-2,3,4-trihydroxy-N-octyl-6-oxa-8-azabicyclo[3.2.1]octane-8-carbothioamide (CCD ID: NCW) (formula: C₁₅H₂₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	1
			30	23	2	4	1		
2	B	1	Total	C	N	O	S	0	1
			25	18	2	4	1		

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0
3	B	1	Total Ca 1 1	0	0

- Molecule 4 is ACETATE ION (CCD ID: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 4 4	0	1
4	A	1	Total C O 4 2 2	0	0
4	B	1	Total C O 4 2 2	0	0

- Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	410	Total O 410 410	0	6
5	B	259	Total O 259 259	0	12

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: BETA-GLUCOSIDASE A

Chain A: 



• Molecule 1: BETA-GLUCOSIDASE A

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	93.70Å 94.41Å 113.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	72.55 – 1.90 72.46 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.7 (72.55-1.90) 99.7 (72.46-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.14 (at 1.90Å)	Xtriage
Refinement program	REFMAC 5.4.0065	Depositor
R, R_{free}	0.195 , 0.239 0.195 , 0.236	Depositor DCC
R_{free} test set	3981 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å ²)	32.5	Xtriage
Anisotropy	0.020	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 63.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.013 for k,h,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8070	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.50% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, CA, NCW

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.85	2/3891 (0.1%)	0.88	3/5277 (0.1%)
1	B	0.75	1/3751 (0.0%)	0.88	1/5108 (0.0%)
All	All	0.80	3/7642 (0.0%)	0.88	4/10385 (0.0%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	344	PRO	CA-C	7.44	1.55	1.51
1	A	403	ASN	C-O	-5.51	1.18	1.24
1	A	24	SER	C-O	5.31	1.26	1.23

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	220	VAL	N-CA-C	5.72	116.17	108.17
1	A	231	GLU	N-CA-C	-5.49	103.38	110.19
1	A	416	ILE	N-CA-C	-5.28	106.73	112.80
1	A	22	GLU	N-CA-C	5.07	116.61	111.14

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3720	0	3627	33	0
1	B	3608	0	3378	40	0
2	A	30	0	36	12	0
2	B	25	0	18	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	12	0	9	0	0
4	B	4	0	3	2	0
5	A	410	0	0	0	0
5	B	259	0	0	6	0
All	All	8070	0	7071	74	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (74) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:231:GLU:O	1:A:232:LYS:HB2	1.39	1.15
1:B:52:ASP:OD2	5:B:2045[A]:HOH:O	1.81	0.95
1:A:406:TRP:HB2	2:A:1446[A]:NCW:H122	1.52	0.91
1:A:405:GLU:OE2	2:A:1446[A]:NCW:H101	1.74	0.87
1:B:270[B]:GLU:OE2	5:B:2189[B]:HOH:O	1.92	0.86
1:B:299:LEU:HD23	1:B:313:VAL:HG22	1.60	0.83
1:A:406:TRP:HB2	2:A:1446[A]:NCW:C12	2.09	0.82
1:B:270[B]:GLU:CD	5:B:2189[B]:HOH:O	2.31	0.74
1:B:270[A]:GLU:HG2	5:B:2180:HOH:O	1.86	0.73
1:A:407:ALA:HB3	2:A:1446[A]:NCW:H132	1.72	0.72
1:A:231:GLU:O	1:A:232:LYS:CB	2.28	0.68
1:B:315:ARG:HB3	1:B:317:LEU:HD13	1.76	0.66
1:B:358:ASP:OD1	1:B:429:LYS:NZ	2.29	0.64
1:A:405:GLU:OE2	2:A:1446[A]:NCW:C10	2.45	0.64
1:B:333:TRP:CZ2	1:B:337:LYS:HD3	2.35	0.61
1:A:407:ALA:H	2:A:1446[A]:NCW:H122	1.67	0.60
1:A:237:ARG:NH2	1:A:266[A]:GLU:OE2	2.35	0.60
1:B:421:TYR:HB3	4:B:1448:ACT:H2	1.86	0.58
1:A:160:ASN:OD1	1:A:216:LYS:HE2	2.03	0.58
1:A:406:TRP:CB	2:A:1446[A]:NCW:H122	2.31	0.57
1:B:299:LEU:HD23	1:B:313:VAL:CG2	2.33	0.55
1:B:369:GLN:HA	1:B:372:ILE:HD12	1.89	0.54
1:B:299:LEU:HB2	1:B:315:ARG:HD3	1.90	0.54
1:B:40:HIS:NE2	5:B:2045[A]:HOH:O	2.08	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:233:GLU:O	1:B:237[A]:ARG:HG3	2.07	0.54
1:A:407:ALA:N	2:A:1446[A]:NCW:H122	2.24	0.52
1:A:14[B]:VAL:HG23	1:A:73:VAL:HG11	1.92	0.52
1:B:315:ARG:CB	1:B:317:LEU:HD13	2.39	0.52
1:A:300:VAL:HG13	1:A:310:VAL:HG13	1.92	0.51
1:B:402:ASP:OD2	1:B:418:TYR:HA	2.11	0.51
1:B:20:GLN:O	1:B:403:ASN:HB2	2.11	0.50
1:B:402:ASP:OD1	1:B:413:ARG:HB3	2.11	0.50
1:A:406:TRP:HB2	2:A:1446[A]:NCW:H121	1.92	0.50
1:A:14[B]:VAL:HG11	1:A:400:LEU:HB2	1.93	0.49
1:B:77:ARG:NH1	1:B:351:GLU:HG3	2.28	0.49
1:B:184[A]:MET:SD	1:B:189:VAL:HG11	2.52	0.49
1:B:63:GLU:O	1:B:67:ILE:HG23	2.13	0.48
2:A:1446[B]:NCW:H111	2:A:1446[B]:NCW:H8C1	1.65	0.48
1:B:421:TYR:CB	4:B:1448:ACT:H2	2.44	0.47
1:A:303:ASP:O	1:A:309:LYS:HA	2.13	0.47
1:B:243:HIS:O	1:B:247:ASN:HB2	2.14	0.47
1:B:315:ARG:HB3	1:B:317:LEU:CD1	2.42	0.47
1:B:418:TYR:HB2	1:B:429:LYS:HE2	1.97	0.46
1:A:20:GLN:O	1:A:403:ASN:HB2	2.16	0.46
1:B:398:TRP:HA	1:B:399:SER:HA	1.73	0.45
1:B:67:ILE:HD11	1:B:417:VAL:CG2	2.46	0.45
1:A:144:ALA:HB2	1:A:203[B]:ARG:HG2	1.98	0.45
1:B:110:GLU:C	1:B:112:GLY:H	2.24	0.44
1:A:109:LEU:HD21	1:A:157:ARG:HB3	1.98	0.44
1:B:216:LYS:NZ	1:B:346:GLU:OE1	2.30	0.44
1:A:14[B]:VAL:HG11	1:A:400:LEU:CA	2.48	0.44
1:A:144:ALA:HB2	1:A:203[B]:ARG:CG	2.48	0.44
1:A:361:SER:HB2	1:A:363:ASP:OD1	2.17	0.44
1:A:14[B]:VAL:CG1	1:A:400:LEU:HB2	2.48	0.43
1:A:185[B]:ARG:HE	1:A:185[B]:ARG:HB2	1.56	0.43
1:A:192[A]:ARG:NH1	1:A:271:TYR:O	2.53	0.42
1:B:267:PHE:CE1	1:B:308:ALA:HB2	2.55	0.42
1:B:248:TYR:CD1	1:B:249:PRO:HD3	2.55	0.42
1:B:71:LEU:O	1:B:436:SER:HB2	2.20	0.42
1:B:192:ARG:HG2	1:B:273:PRO:HG3	2.02	0.42
1:B:26:LEU:HD21	1:B:32:MET:HG2	2.00	0.42
1:B:65:ILE:HG21	1:B:107:THR:HG22	2.02	0.42
1:B:355:ALA:HA	1:B:413:ARG:O	2.19	0.41
1:A:398:TRP:HA	1:A:399:SER:HA	1.83	0.41
1:B:67:ILE:HD13	1:B:428:VAL:HG22	2.02	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:407:ALA:CB	2:A:1446[A]:NCW:H132	2.47	0.41
1:A:168:TRP:CZ3	2:A:1446[B]:NCW:H153	2.56	0.41
1:A:14[B]:VAL:HG11	1:A:400:LEU:HA	2.02	0.41
1:A:243:HIS:O	1:A:247:ASN:HB2	2.21	0.41
1:A:266[B]:GLU:HA	1:A:266[B]:GLU:OE2	2.21	0.41
1:A:280:MET:O	1:A:284:GLN:HG3	2.21	0.41
1:B:103:ARG:NH2	5:B:2085[A]:HOH:O	2.42	0.41
1:B:328:PRO:HA	1:B:378:HIS:NE2	2.36	0.41
1:B:139:ILE:HA	1:B:142:TRP:CE3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	455/468 (97%)	445 (98%)	9 (2%)	1 (0%)	43	36
1	B	451/468 (96%)	428 (95%)	22 (5%)	1 (0%)	43	36
All	All	906/936 (97%)	873 (96%)	31 (3%)	2 (0%)	43	36

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	213	LYS
1	B	7	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	390/399 (98%)	385 (99%)	5 (1%)	61	61
1	B	366/399 (92%)	355 (97%)	11 (3%)	36	30
All	All	756/798 (95%)	740 (98%)	16 (2%)	47	44

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	109	LEU
1	A	203[A]	ARG
1	A	203[B]	ARG
1	A	236	ILE
1	A	317	LEU
1	B	67	ILE
1	B	69	GLU
1	B	234	GLU
1	B	296	SER
1	B	313	VAL
1	B	336	LYS
1	B	350	THR
1	B	359	VAL
1	B	375	LEU
1	B	381	GLN
1	B	444	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	129	GLN
1	A	387	GLN
1	B	40	HIS
1	B	94	ASN
1	B	95	GLN
1	B	160	ASN
1	B	253	ASN
1	B	369	GLN
1	B	387	GLN
1	B	441	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 10 ligands modelled in this entry, 2 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	ACT	B	1448	-	3,3,3	0.84	0	3,3,3	0.42	0
4	ACT	A	1449	-	3,3,3	0.82	0	3,3,3	0.75	0
2	NCW	B	1446[B]	-	22,23,23	1.12	1 (4%)	26,31,31	1.22	2 (7%)
2	NCW	B	1446[A]	-	22,23,23	1.12	1 (4%)	26,31,31	1.21	2 (7%)
2	NCW	A	1446[B]	-	22,23,23	1.28	1 (4%)	26,31,31	1.43	5 (19%)
2	NCW	A	1446[A]	-	22,23,23	1.31	1 (4%)	26,31,31	1.45	5 (19%)
4	ACT	A	1448[B]	-	3,3,3	0.95	0	3,3,3	0.60	0
4	ACT	A	1448[A]	-	3,3,3	0.91	0	3,3,3	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NCW	B	1446[A]	-	-	6/13/42/42	0/3/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NCW	A	1446[B]	-	-	4/13/42/42	0/3/2/2
2	NCW	A	1446[A]	-	-	6/13/42/42	0/3/2/2
2	NCW	B	1446[B]	-	-	7/13/42/42	0/3/2/2

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	1446[A]	NCW	C7-S1	-5.03	1.59	1.67
2	A	1446[B]	NCW	C7-S1	-5.03	1.59	1.67
2	B	1446[A]	NCW	C7-S1	-4.60	1.60	1.67
2	B	1446[B]	NCW	C7-S1	-4.60	1.60	1.67

All (14) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1446[A]	NCW	C6-C5-C4	-4.17	108.70	113.30
2	B	1446[B]	NCW	C6-C5-C4	-4.17	108.70	113.30
2	A	1446[A]	NCW	C6-C5-C4	-3.64	109.28	113.30
2	A	1446[B]	NCW	C6-C5-C4	-3.64	109.28	113.30
2	A	1446[A]	NCW	S1-C7-N2	-2.42	117.94	122.47
2	A	1446[B]	NCW	S1-C7-N2	-2.42	117.94	122.47
2	A	1446[A]	NCW	O3-C3-C2	-2.40	104.72	110.38
2	A	1446[B]	NCW	O3-C3-C2	-2.40	104.72	110.38
2	B	1446[A]	NCW	C6-C5-N1	-2.24	98.24	99.99
2	B	1446[B]	NCW	C6-C5-N1	-2.24	98.24	99.99
2	A	1446[A]	NCW	N2-C7-N1	2.10	119.79	116.15
2	A	1446[B]	NCW	N2-C7-N1	2.10	119.79	116.15
2	A	1446[A]	NCW	O4-C4-C5	-2.10	106.03	109.87
2	A	1446[B]	NCW	O4-C4-C5	-2.10	106.03	109.87

There are no chirality outliers.

All (23) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1446[A]	NCW	N1-C7-N2-C8
2	B	1446[A]	NCW	S1-C7-N2-C8
2	B	1446[A]	NCW	N1-C7-N2-C8
2	B	1446[B]	NCW	S1-C7-N2-C8
2	B	1446[B]	NCW	N1-C7-N2-C8
2	A	1446[A]	NCW	C11-C12-C13-C14

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1446[A]	NCW	C10-C11-C12-C13
2	A	1446[B]	NCW	C9-C10-C11-C12
2	B	1446[A]	NCW	C9-C10-C11-C12
2	B	1446[B]	NCW	C9-C10-C11-C12
2	B	1446[A]	NCW	C11-C12-C13-C14
2	A	1446[A]	NCW	C11-C10-C9-C8
2	B	1446[A]	NCW	C11-C10-C9-C8
2	B	1446[B]	NCW	C11-C10-C9-C8
2	B	1446[B]	NCW	C10-C11-C12-C13
2	B	1446[B]	NCW	C11-C12-C13-C14
2	B	1446[A]	NCW	C10-C11-C12-C13
2	B	1446[B]	NCW	C12-C13-C14-C15
2	A	1446[B]	NCW	C10-C11-C12-C13
2	A	1446[B]	NCW	C9-C8-N2-C7
2	A	1446[A]	NCW	S1-C7-N2-C8
2	A	1446[B]	NCW	C11-C10-C9-C8
2	A	1446[A]	NCW	N2-C8-C9-C10


There are no ring outliers.

3 monomers are involved in 14 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	1448	ACT	2	0
2	A	1446[B]	NCW	2	0
2	A	1446[A]	NCW	10	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.

Ligand NCW B 1446 (B)



Ligand NCW B 1446 (A)





5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	438/468 (93%)	-0.16	5 (1%) 78 80	10, 29, 44, 58	22 (5%)
1	B	443/468 (94%)	0.72	51 (11%) 9 10	17, 42, 66, 78	10 (2%)
All	All	881/936 (94%)	0.28	56 (6%) 25 27	10, 33, 62, 78	32 (3%)

All (56) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	439	VAL	5.2
1	B	383	TRP	3.8
1	B	438	VAL	3.8
1	B	440	LYS	3.7
1	B	444	LEU	3.6
1	B	382	ALA	3.6
1	B	229	ALA	3.5
1	B	112	GLY	3.5
1	B	317	LEU	3.4
1	B	359	VAL	3.4
1	B	434	TRP	3.4
1	B	333	TRP	3.3
1	A	333	TRP	3.2
1	B	433	TYR	3.1
1	B	327	VAL	3.0
1	B	416	ILE	2.9
1	B	12	TRP	2.9
1	B	385	ALA	2.8
1	B	375	LEU	2.7
1	B	73	VAL	2.7
1	A	236	ILE	2.6
1	B	374	TYR	2.6
1	B	443	GLY	2.6
1	B	417	VAL	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	229	ALA	2.5
1	B	3	VAL	2.5
1	B	300	VAL	2.5
1	B	306	ALA	2.5
1	B	299	LEU	2.5
1	A	362	GLU	2.4
1	B	113	ILE	2.4
1	B	68	ILE	2.4
1	B	372	ILE	2.4
1	B	356	PHE	2.4
1	B	4	LYS	2.3
1	B	111	LYS	2.3
1	B	212	VAL	2.3
1	B	316	ASP	2.3
1	B	360	VAL	2.3
1	B	397	VAL	2.3
1	B	386	ILE	2.2
1	B	367	HIS	2.2
1	A	300	VAL	2.2
1	B	389	GLY	2.2
1	B	381	GLN	2.2
1	B	418	TYR	2.1
1	B	366	VAL	2.1
1	B	428	VAL	2.1
1	B	7	PRO	2.1
1	B	430	ASP	2.1
1	B	361[A]	SER	2.0
1	B	441	ASN	2.0
1	B	392	LEU	2.0
1	B	311	SER	2.0
1	B	370	ASN	2.0
1	B	435	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	ACT	A	1449	4/4	0.58	0.23	48,49,49,49	0
4	ACT	B	1448	4/4	0.83	0.15	51,52,52,53	0
2	NCW	B	1446[A]	22/22	0.86	0.14	39,47,49,50	3
2	NCW	B	1446[B]	22/22	0.86	0.14	39,48,51,52	3
4	ACT	A	1448[A]	4/4	0.88	0.12	29,29,29,30	4
4	ACT	A	1448[B]	4/4	0.88	0.12	30,31,32,33	4
2	NCW	A	1446[A]	22/22	0.88	0.13	12,33,38,39	8
2	NCW	A	1446[B]	22/22	0.88	0.13	27,34,38,39	8
3	CA	A	1447	1/1	0.94	0.14	48,48,48,48	0
3	CA	B	1447	1/1	0.98	0.09	38,38,38,38	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NCW B 1446 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NCW B 1446 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NCW A 1446 (A):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NCW A 1446 (B):**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.