



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 09:12 PM UTC

PDB ID : 5VRB / pdb_00005vrb
Title : Crystal structure of a transketolase from *Neisseria gonorrhoeae*
Authors : Seattle Structural Genomics Center for Infectious Disease (SSGCID)
Deposited on : 2017-05-10
Resolution : 1.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	667	
1	B	667	
1	C	667	
1	D	667	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 21287 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transketolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	627	Total	C	N	O	S	0	14	0
			4786	3040	816	906	24			
1	B	646	Total	C	N	O	S	0	10	0
			4915	3118	840	933	24			
1	C	615	Total	C	N	O	S	0	6	0
			4651	2956	798	875	22			
1	D	640	Total	C	N	O	S	0	5	0
			4839	3075	832	909	23			

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP B4RKU9
A	-6	ALA	-	expression tag	UNP B4RKU9
A	-5	HIS	-	expression tag	UNP B4RKU9
A	-4	HIS	-	expression tag	UNP B4RKU9
A	-3	HIS	-	expression tag	UNP B4RKU9
A	-2	HIS	-	expression tag	UNP B4RKU9
A	-1	HIS	-	expression tag	UNP B4RKU9
A	0	HIS	-	expression tag	UNP B4RKU9
B	-7	MET	-	initiating methionine	UNP B4RKU9
B	-6	ALA	-	expression tag	UNP B4RKU9
B	-5	HIS	-	expression tag	UNP B4RKU9
B	-4	HIS	-	expression tag	UNP B4RKU9
B	-3	HIS	-	expression tag	UNP B4RKU9
B	-2	HIS	-	expression tag	UNP B4RKU9
B	-1	HIS	-	expression tag	UNP B4RKU9
B	0	HIS	-	expression tag	UNP B4RKU9
C	-7	MET	-	initiating methionine	UNP B4RKU9
C	-6	ALA	-	expression tag	UNP B4RKU9
C	-5	HIS	-	expression tag	UNP B4RKU9
C	-4	HIS	-	expression tag	UNP B4RKU9
C	-3	HIS	-	expression tag	UNP B4RKU9

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-2	HIS	-	expression tag	UNP B4RKU9
C	-1	HIS	-	expression tag	UNP B4RKU9
C	0	HIS	-	expression tag	UNP B4RKU9
D	-7	MET	-	initiating methionine	UNP B4RKU9
D	-6	ALA	-	expression tag	UNP B4RKU9
D	-5	HIS	-	expression tag	UNP B4RKU9
D	-4	HIS	-	expression tag	UNP B4RKU9
D	-3	HIS	-	expression tag	UNP B4RKU9
D	-2	HIS	-	expression tag	UNP B4RKU9
D	-1	HIS	-	expression tag	UNP B4RKU9
D	0	HIS	-	expression tag	UNP B4RKU9

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	C	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0
2	D	1	Total C O 4 2 2	0	0

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	3	Total 3	Ca 3	0	0
3	B	3	Total 3	Ca 3	0	0
3	C	3	Total 3	Ca 3	0	0
3	D	3	Total 3	Ca 3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	492	Total 498	O 498	0	6
4	B	583	Total 592	O 592	0	9
4	C	509	Total 513	O 513	0	4
4	D	455	Total 457	O 457	0	2

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Transketolase

Chain A: 




• Molecule 1: Transketolase

Chain B: 



• Molecule 1: Transketolase

Chain C: 



• Molecule 1: Transketolase

Chain D: 

MET	ALA	HIS	HIS	HIS	HIS	HIS	M1	N61	I81	L90	H97	T110	D143	C154	S165	T169	K174	N182	ASN	ILE	SER	ILE	ASP	GLY	LYS	VAL	D191	K241	S254	H255	K256	Q312	L330	K348	ALA	GLU	THR	VAL	ALA	THR	ARG	LYS
ALA	SER	GLN	N360	L365	V373	D396	P426	A429	V455	T469	K505	L520	A537	Y538	V539	S556	V567	R576	M580	S584	V585	F586	D587	I605	A609	K617	F646	T647	V648	V655	I659											

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	82.45Å 114.21Å 125.69Å 90.00° 90.11° 90.00°	Depositor
Resolution (Å)	37.59 – 1.85 37.59 – 1.85	Depositor EDS
% Data completeness (in resolution range)	98.7 (37.59-1.85) 98.7 (37.59-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.67 (at 1.86Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.150 , 0.191 0.152 , 0.192	Depositor DCC
R_{free} test set	1970 reflections (0.99%)	wwPDB-VP
Wilson B-factor (Å ²)	18.4	Xtriage
Anisotropy	0.310	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 36.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.45$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.087 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	21287	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.62% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: OCS, EDO, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.27	0/4921	0.47	0/6696
1	B	0.27	0/5045	0.47	0/6855
1	C	0.26	0/4762	0.46	0/6476
1	D	0.25	0/4954	0.44	0/6731
All	All	0.26	0/19682	0.46	0/26758

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4786	0	4623	23	0
1	B	4915	0	4762	26	0
1	C	4651	0	4482	24	0
1	D	4839	0	4693	21	0
2	A	4	0	6	0	0
2	C	8	0	10	2	0
2	D	12	0	16	0	0
3	A	3	0	0	0	0
3	B	3	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	C	3	0	0	0	0
3	D	3	0	0	0	0
4	A	498	0	0	3	0
4	B	592	0	0	4	0
4	C	513	0	0	6	0
4	D	457	0	0	2	0
All	All	21287	0	18592	86	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 86 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:639[B]:LEU:HD21	1:B:90:LEU:HD22	1.56	0.87
1:B:258:HIS:ND1	4:B:801:HOH:O	2.19	0.76
1:C:639:LEU:HD21	1:D:90:LEU:HD22	1.74	0.69
1:C:548:GLN:NE2	4:C:803:HOH:O	2.20	0.66
1:A:137:ASP:OD1	1:A:315:TYR:OH	2.14	0.65

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	634/667 (95%)	621 (98%)	12 (2%)	1 (0%)	43	31
1	B	649/667 (97%)	634 (98%)	14 (2%)	1 (0%)	43	31
1	C	608/667 (91%)	595 (98%)	12 (2%)	1 (0%)	43	31
1	D	638/667 (96%)	624 (98%)	14 (2%)	0	100	100
All	All	2529/2668 (95%)	2474 (98%)	52 (2%)	3 (0%)	48	35

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	64	ALA
1	C	111	GLY
1	A	111	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	486/528 (92%)	484 (100%)	2 (0%)	84	82
1	B	501/528 (95%)	496 (99%)	5 (1%)	68	60
1	C	468/528 (89%)	465 (99%)	3 (1%)	78	73
1	D	488/528 (92%)	484 (99%)	4 (1%)	73	67
All	All	1943/2112 (92%)	1929 (99%)	14 (1%)	76	70

5 of 14 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	61	ASN
1	C	365	LEU
1	D	584	SER
1	D	312	GLN
1	D	365	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 12 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	548	GLN
1	C	17	GLN
1	D	310	GLN
1	C	63	HIS
1	B	115	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCS	D	154	1	6,8,9	1.61	1 (16%)	7,11,13	1.40	1 (14%)
1	OCS	C	154	1	6,8,9	1.12	0	7,11,13	1.21	1 (14%)
1	OCS	A	154	1,3	6,8,9	1.33	1 (16%)	7,11,13	1.09	1 (14%)
1	OCS	B	154	1	6,8,9	1.34	1 (16%)	7,11,13	1.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCS	D	154	1	-	0/4/7/9	-
1	OCS	C	154	1	-	0/4/7/9	-
1	OCS	A	154	1,3	-	0/4/7/9	-
1	OCS	B	154	1	-	3/4/7/9	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	154	OCS	CB-CA	-3.27	1.51	1.53
1	A	154	OCS	CB-CA	-2.16	1.52	1.53
1	B	154	OCS	CB-CA	-2.02	1.52	1.53

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	154	OCS	OD3-SG-CB	3.27	111.63	106.76
1	C	154	OCS	OD3-SG-CB	2.41	110.36	106.76
1	A	154	OCS	OD3-SG-CB	2.37	110.30	106.76

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	B	154	OCS	CA-CB-SG-OD2
1	B	154	OCS	CA-CB-SG-OD1
1	B	154	OCS	CA-CB-SG-OD3

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	154	OCS	1	0
1	B	154	OCS	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 12 are monoatomic - leaving 6 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	D	702	-	3,3,3	0.41	0	2,2,2	0.45	0
2	EDO	D	703	-	3,3,3	0.45	0	2,2,2	0.46	0
2	EDO	C	701	3	3,3,3	0.43	0	2,2,2	0.47	0
2	EDO	D	701	3	3,3,3	0.41	0	2,2,2	0.35	0
2	EDO	A	701	-	3,3,3	0.39	0	2,2,2	0.66	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	C	702	-	3,3,3	0.29	0	2,2,2	0.23	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	D	702	-	-	0/1/1/1	-
2	EDO	D	703	-	-	0/1/1/1	-
2	EDO	C	701	3	-	0/1/1/1	-
2	EDO	D	701	3	-	0/1/1/1	-
2	EDO	A	701	-	-	0/1/1/1	-
2	EDO	C	702	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	C	702	EDO	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	626/667 (93%)	-1.59	0 100 100	7, 19, 37, 70	14 (2%)
1	B	645/667 (96%)	-1.61	0 100 100	7, 18, 35, 68	10 (1%)
1	C	614/667 (92%)	-1.59	0 100 100	8, 19, 40, 71	6 (0%)
1	D	639/667 (95%)	-1.56	0 100 100	11, 21, 39, 74	5 (0%)
All	All	2524/2668 (94%)	-1.59	0 100 100	7, 19, 38, 74	35 (1%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OCS	A	154	9/10	1.00	0.03	10,12,17,37	0
1	OCS	B	154	9/10	1.00	0.03	10,13,24,39	0
1	OCS	C	154	9/10	1.00	0.02	12,15,23,38	0
1	OCS	D	154	9/10	1.00	0.03	10,12,20,42	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	EDO	D	702	4/4	0.99	0.04	30,30,35,44	0
2	EDO	C	701	4/4	1.00	0.02	21,23,24,26	0
2	EDO	C	702	4/4	1.00	0.03	20,23,29,31	0
2	EDO	D	701	4/4	1.00	0.02	27,27,28,33	0
2	EDO	A	701	4/4	1.00	0.01	19,22,22,23	0
2	EDO	D	703	4/4	1.00	0.03	20,22,25,28	0
3	CA	A	702	1/1	1.00	0.02	22,22,22,22	0
3	CA	A	703	1/1	1.00	0.07	50,50,50,50	0
3	CA	A	704	1/1	1.00	0.02	31,31,31,31	0
3	CA	B	701	1/1	1.00	0.02	25,25,25,25	0
3	CA	B	702	1/1	1.00	0.04	22,22,22,22	0
3	CA	B	703	1/1	1.00	0.03	30,30,30,30	0
3	CA	C	703	1/1	1.00	0.01	19,19,19,19	0
3	CA	C	704	1/1	1.00	0.02	32,32,32,32	0
3	CA	C	705	1/1	1.00	0.03	47,47,47,47	0
3	CA	D	704	1/1	1.00	0.05	39,39,39,39	0
3	CA	D	705	1/1	1.00	0.04	34,34,34,34	0
3	CA	D	706	1/1	1.00	0.02	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.