



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 03:05 PM UTC

PDB ID : 5VRD / pdb_00005vrd
Title : Crystal structure for Methylobacterium extorquens PqqCD (natural fusion)
Authors : Evans III, R.L.; Wilmot, C.M.; Esler, M.A.
Deposited on : 2017-05-10
Resolution : 2.85 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1407 (2.88-2.84)
Clashscore	190562	1446 (2.88-2.84)
Ramachandran outliers	187476	1406 (2.88-2.84)
Sidechain outliers	187428	1407 (2.88-2.84)
RSRZ outliers	180081	1408 (2.88-2.84)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	392	 3% 30% 19% 2% 48%
1	B	392	 2% 35% 10% 2% 50%
1	C	392	 3% 40% 9% 2% 48%
1	D	392	 4% 34% 15% 2% 47%

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 6570 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Bifunctional coenzyme PQQ synthesis protein C/D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	D	209	Total	C	N	O	S	0	0	0
			1671	1075	292	299	5			
1	A	208	Total	C	N	O	S	0	0	0
			1678	1077	298	297	6			
1	B	195	Total	C	N	O	S	0	0	0
			1579	1015	281	278	5			
1	C	203	Total	C	N	O	S	0	0	0
			1642	1055	288	294	5			

There are 80 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	-19	MET	-	initiating methionine	UNP Q49150
D	-18	GLY	-	expression tag	UNP Q49150
D	-17	SER	-	expression tag	UNP Q49150
D	-16	SER	-	expression tag	UNP Q49150
D	-15	HIS	-	expression tag	UNP Q49150
D	-14	HIS	-	expression tag	UNP Q49150
D	-13	HIS	-	expression tag	UNP Q49150
D	-12	HIS	-	expression tag	UNP Q49150
D	-11	HIS	-	expression tag	UNP Q49150
D	-10	HIS	-	expression tag	UNP Q49150
D	-9	SER	-	expression tag	UNP Q49150
D	-8	SER	-	expression tag	UNP Q49150
D	-7	GLY	-	expression tag	UNP Q49150
D	-6	LEU	-	expression tag	UNP Q49150
D	-5	VAL	-	expression tag	UNP Q49150
D	-4	PRO	-	expression tag	UNP Q49150
D	-3	ARG	-	expression tag	UNP Q49150
D	-2	GLY	-	expression tag	UNP Q49150
D	-1	SER	-	expression tag	UNP Q49150
D	0	HIS	-	expression tag	UNP Q49150
A	-19	MET	-	initiating methionine	UNP Q49150

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	GLY	-	expression tag	UNP Q49150
A	-17	SER	-	expression tag	UNP Q49150
A	-16	SER	-	expression tag	UNP Q49150
A	-15	HIS	-	expression tag	UNP Q49150
A	-14	HIS	-	expression tag	UNP Q49150
A	-13	HIS	-	expression tag	UNP Q49150
A	-12	HIS	-	expression tag	UNP Q49150
A	-11	HIS	-	expression tag	UNP Q49150
A	-10	HIS	-	expression tag	UNP Q49150
A	-9	SER	-	expression tag	UNP Q49150
A	-8	SER	-	expression tag	UNP Q49150
A	-7	GLY	-	expression tag	UNP Q49150
A	-6	LEU	-	expression tag	UNP Q49150
A	-5	VAL	-	expression tag	UNP Q49150
A	-4	PRO	-	expression tag	UNP Q49150
A	-3	ARG	-	expression tag	UNP Q49150
A	-2	GLY	-	expression tag	UNP Q49150
A	-1	SER	-	expression tag	UNP Q49150
A	0	HIS	-	expression tag	UNP Q49150
B	-19	MET	-	initiating methionine	UNP Q49150
B	-18	GLY	-	expression tag	UNP Q49150
B	-17	SER	-	expression tag	UNP Q49150
B	-16	SER	-	expression tag	UNP Q49150
B	-15	HIS	-	expression tag	UNP Q49150
B	-14	HIS	-	expression tag	UNP Q49150
B	-13	HIS	-	expression tag	UNP Q49150
B	-12	HIS	-	expression tag	UNP Q49150
B	-11	HIS	-	expression tag	UNP Q49150
B	-10	HIS	-	expression tag	UNP Q49150
B	-9	SER	-	expression tag	UNP Q49150
B	-8	SER	-	expression tag	UNP Q49150
B	-7	GLY	-	expression tag	UNP Q49150
B	-6	LEU	-	expression tag	UNP Q49150
B	-5	VAL	-	expression tag	UNP Q49150
B	-4	PRO	-	expression tag	UNP Q49150
B	-3	ARG	-	expression tag	UNP Q49150
B	-2	GLY	-	expression tag	UNP Q49150
B	-1	SER	-	expression tag	UNP Q49150
B	0	HIS	-	expression tag	UNP Q49150
C	-19	MET	-	initiating methionine	UNP Q49150
C	-18	GLY	-	expression tag	UNP Q49150
C	-17	SER	-	expression tag	UNP Q49150

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	-16	SER	-	expression tag	UNP Q49150
C	-15	HIS	-	expression tag	UNP Q49150
C	-14	HIS	-	expression tag	UNP Q49150
C	-13	HIS	-	expression tag	UNP Q49150
C	-12	HIS	-	expression tag	UNP Q49150
C	-11	HIS	-	expression tag	UNP Q49150
C	-10	HIS	-	expression tag	UNP Q49150
C	-9	SER	-	expression tag	UNP Q49150
C	-8	SER	-	expression tag	UNP Q49150
C	-7	GLY	-	expression tag	UNP Q49150
C	-6	LEU	-	expression tag	UNP Q49150
C	-5	VAL	-	expression tag	UNP Q49150
C	-4	PRO	-	expression tag	UNP Q49150
C	-3	ARG	-	expression tag	UNP Q49150
C	-2	GLY	-	expression tag	UNP Q49150
C	-1	SER	-	expression tag	UNP Q49150
C	0	HIS	-	expression tag	UNP Q49150


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Bifunctional coenzyme PQQ synthesis protein C/D



- Molecule 1: Bifunctional coenzyme PQQ synthesis protein C/D



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 21 2	Depositor
Cell constants a, b, c, α , β , γ	103.94Å 103.94Å 243.49Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	243.49 – 2.85 95.59 – 2.85	Depositor EDS
% Data completeness (in resolution range)	98.4 (243.49-2.85) 98.4 (95.59-2.85)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.31 (at 2.85Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.253 , 0.327 0.255 , 0.328	Depositor DCC
R_{free} test set	1597 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	77.6	Xtriage
Anisotropy	0.384	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 49.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	6570	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.66% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.32	11/1718 (0.6%)	1.42	18/2325 (0.8%)
1	B	1.15	4/1616 (0.2%)	1.31	5/2185 (0.2%)
1	C	1.10	3/1679 (0.2%)	1.28	15/2271 (0.7%)
1	D	1.33	10/1710 (0.6%)	1.37	14/2317 (0.6%)
All	All	1.23	28/6723 (0.4%)	1.35	52/9098 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2
1	B	0	1
All	All	0	3

The worst 5 of 28 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	18	SER	CA-C	10.65	1.67	1.52
1	A	102	ASP	CA-C	9.11	1.63	1.52
1	D	102	ASP	N-CA	8.36	1.56	1.46
1	A	247	PRO	CA-C	8.26	1.59	1.52
1	D	19	HIS	N-CA	8.25	1.56	1.46

The worst 5 of 52 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	104	GLY	N-CA-C	9.60	123.64	112.50
1	D	246	VAL	CA-C-N	8.84	129.49	120.38
1	D	246	VAL	C-N-CA	8.84	129.49	120.38
1	D	247	PRO	N-CA-C	7.80	120.22	110.70
1	A	105	ILE	CB-CA-C	-7.47	101.91	112.22

There are no chirality outliers.

All (3) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	GLY	Peptide
1	A	250	ALA	Peptide
1	B	245	MET	Peptide

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1678	0	1637	31	0
1	B	1579	0	1555	38	0
1	C	1642	0	1595	12	0
1	D	1671	0	1616	32	0
All	All	6570	0	6403	110	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 110 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:159:MET:HE3	1:B:202:LEU:HD22	1.58	0.84
1:B:227:ASN:O	1:B:231:THR:HG23	1.90	0.72
1:B:109:LEU:HD21	1:B:123:LEU:HD13	1.72	0.70
1:A:71:LYS:HE2	1:A:139:TYR:CZ	2.26	0.70
1:D:65:GLN:HA	1:D:68:ILE:HD12	1.73	0.68

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	200/392 (51%)	181 (90%)	14 (7%)	5 (2%)	4	10
1	B	185/392 (47%)	169 (91%)	14 (8%)	2 (1%)	11	24
1	C	193/392 (49%)	177 (92%)	13 (7%)	3 (2%)	7	17
1	D	199/392 (51%)	175 (88%)	16 (8%)	8 (4%)	2	4
All	All	777/1568 (50%)	702 (90%)	57 (7%)	18 (2%)	5	11

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	102	ASP
1	D	243	PRO
1	D	247	PRO
1	A	17	LEU
1	B	246	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	168/323 (52%)	133 (79%)	35 (21%)	1	1
1	B	161/323 (50%)	140 (87%)	21 (13%)	4	7
1	C	165/323 (51%)	140 (85%)	25 (15%)	3	5
1	D	166/323 (51%)	143 (86%)	23 (14%)	3	6
All	All	660/1292 (51%)	556 (84%)	104 (16%)	2	4

5 of 104 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	241	VAL
1	B	215	GLN
1	C	199	ASP
1	B	37	LEU
1	B	128	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	A	232	GLN
1	B	41	HIS
1	C	41	HIS
1	B	215	GLN
1	A	208	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	208/392 (53%)	0.23	13 (6%) 26 19	48, 72, 102, 133	0
1	B	195/392 (49%)	0.16	6 (3%) 51 43	46, 75, 98, 115	0
1	C	203/392 (51%)	0.36	11 (5%) 31 23	55, 85, 130, 155	0
1	D	209/392 (53%)	0.38	17 (8%) 18 13	52, 74, 113, 124	0
All	All	815/1568 (51%)	0.28	47 (5%) 29 21	46, 76, 113, 155	0

The worst 5 of 47 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	101	GLY	5.5
1	D	198	ALA	5.0
1	C	198	ALA	4.8
1	A	172	MET	4.8
1	C	172	MET	4.5

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.