



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:08 AM UTC

PDB ID : 7VR9 / pdb_00007vr9
Title : Crystal structure of human serum albumin complex with aripiprazole and myristic acid
Authors : Kawai, A.; Yamasaki, K.; Otagiri, M.
Deposited on : 2021-10-22
Resolution : 2.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6319 (2.30-2.30)
Clashscore	190562	6919 (2.30-2.30)
Ramachandran outliers	187476	6854 (2.30-2.30)
Sidechain outliers	187428	6854 (2.30-2.30)
RSRZ outliers	180081	6325 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	 4% 91% 8% .
1	B	585	 4% 89% 9% .

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 9314 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serum albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	578	Total	C	N	O	S	0	1	0
			4432	2802	750	839	41			
1	B	578	Total	C	N	O	S	0	0	0
			4421	2798	747	835	41			

- Molecule 2 is MYRISTIC ACID (CCD ID: MYR) (formula: C₁₄H₂₈O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	14	2		
2	A	1	Total	C	O	0	0
			13	11	2		
2	A	1	Total	C	O	0	0
			13	11	2		
2	A	1	Total	C	O	0	0
			13	11	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			16	14	2		
2	A	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			13	11	2		
2	B	1	Total	C	O	0	0
			13	11	2		
2	B	1	Total	C	O	0	0
			13	11	2		
2	B	1	Total	C	O	0	0
			16	14	2		
2	B	1	Total	C	O	0	0
			16	14	2		

- Molecule 3 is 7-[4-[4-[2,3-bis(chloranyl)phenyl]piperazin-1-yl]butoxy]-3,4-dihydro-1H-quinolin-2-one (CCD ID: 9SC) (formula: $C_{23}H_{27}Cl_2N_3O_2$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	0	0
			30	23	2	3	2		
3	B	1	Total	C	Cl	N	O	0	0
			30	23	2	3	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	106	Total 106	O 106	0	0
4	B	121	Total 121	O 121	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Serum albumin



• Molecule 1: Serum albumin



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	38.83Å 94.04Å 94.87Å 105.29° 89.84° 100.20°	Depositor
Resolution (Å)	35.69 – 2.30 35.69 – 2.30	Depositor EDS
% Data completeness (in resolution range)	97.3 (35.69-2.30) 97.5 (35.69-2.30)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.35 (at 2.29Å)	Xtriage
Refinement program	PHENIX 1.19.2_4158	Depositor
R, R_{free}	0.219 , 0.261 0.218 , 0.260	Depositor DCC
R_{free} test set	2758 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	53.1	Xtriage
Anisotropy	0.329	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 40.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.35$	Xtriage
Estimated twinning fraction	0.130 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	9314	wwPDB-VP
Average B, all atoms (Å ²)	67.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9SC, MYR

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.09	0/4521	0.24	0/6130
1	B	0.09	0/4509	0.23	0/6115
All	All	0.09	0/9030	0.24	0/12245

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4432	0	4169	28	0
1	B	4421	0	4175	30	0
2	A	87	0	135	6	0
2	B	87	0	135	5	0
3	A	30	0	0	2	0
3	B	30	0	0	2	0
4	A	106	0	0	3	0
4	B	121	0	0	0	0
All	All	9314	0	8614	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:B:607:9SC:C7	3:B:607:9SC:CL2	2.43	1.03
3:A:607:9SC:CL2	3:A:607:9SC:C7	2.46	1.00
3:B:607:9SC:CL2	3:B:607:9SC:C8	2.73	0.73
1:A:390:GLN:NE2	4:A:702:HOH:O	2.28	0.65
1:A:425:GLU:OE2	1:A:459:GLN:NE2	2.30	0.65
1:A:241:VAL:HG22	1:A:256:ASP:HB3	1.78	0.64
1:A:488:PHE:HB3	2:A:604:MYR:H71	1.78	0.64
1:B:513:ILE:O	1:B:521:ARG:NH1	2.33	0.62
1:A:212:LYS:NZ	4:A:706:HOH:O	2.32	0.62
1:B:426:VAL:HG21	1:B:460:LEU:HD12	1.82	0.62
1:B:224:PRO:HD2	1:B:296:ASP:HB3	1.83	0.61
1:A:218:ARG:NH1	4:A:708:HOH:O	2.34	0.60
1:A:426:VAL:HG21	1:A:460:LEU:HD13	1.84	0.59
1:B:131:GLU:OE2	1:B:162:LYS:NZ	2.36	0.59
1:B:458:ASN:HB2	1:B:484:ARG:HH22	1.68	0.58
1:B:548:MET:HE3	2:B:605:MYR:H31	1.86	0.57
1:A:414:LYS:O	1:A:472:ARG:NH2	2.38	0.56
1:A:42:LEU:HD22	1:A:73:LYS:HE3	1.88	0.56
1:A:392:CYS:HA	1:A:395:PHE:HB3	1.87	0.55
1:B:121:ASP:OD1	1:B:121:ASP:N	2.38	0.54
1:A:148:TYR:O	1:A:196:GLN:NE2	2.41	0.52
1:B:449:ALA:O	1:B:453:LEU:HB2	2.10	0.52
1:A:238:LEU:HD11	2:A:606:MYR:H121	1.91	0.51
1:B:417:GLN:NE2	1:B:494:ASP:OD2	2.36	0.51
1:B:138:TYR:O	1:B:142:ILE:HG12	2.11	0.51
1:B:100:GLU:OE2	1:B:247:HIS:NE2	2.31	0.51
1:A:10:ARG:NH2	1:A:255:ASP:OD1	2.39	0.49
1:B:567:CYS:O	1:B:571:GLU:N	2.44	0.49
1:B:504:ALA:O	1:B:508:THR:OG1	2.27	0.48
1:B:133:THR:O	1:B:137:LYS:HB2	2.13	0.47
1:B:342:SER:HB3	1:B:446:MET:HG2	1.95	0.47
1:A:186:ARG:NH2	1:A:187:ASP:OD1	2.46	0.47
1:B:513:ILE:HA	1:B:516:LEU:HD12	1.97	0.47
1:B:155:LEU:HD23	1:B:284:LEU:HD21	1.97	0.47
1:B:241:VAL:HG22	1:B:256:ASP:HB3	1.96	0.47
1:B:419:SER:OG	1:B:421:PRO:HD2	2.16	0.46
1:A:433:VAL:HG22	1:A:452:TYR:HB3	1.98	0.46
1:B:542:GLU:OE1	1:B:542:GLU:N	2.49	0.46
1:A:29:GLN:HG2	1:A:147:PRO:HA	1.97	0.45
1:B:532:LEU:HD13	2:B:605:MYR:H131	1.98	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:120:VAL:HG13	1:B:178:LEU:HD23	1.99	0.45
1:B:516:LEU:O	1:B:521:ARG:NH2	2.51	0.44
1:A:224:PRO:HD2	1:A:296:ASP:HB3	2.00	0.44
1:A:502:PHE:HB2	1:A:535:HIS:CE1	2.52	0.43
1:B:14:LEU:HD13	1:B:22:LEU:HD12	2.01	0.42
1:B:383:GLU:HB3	1:B:384:PRO:HD3	2.00	0.42
1:B:441:PRO:O	1:B:445:ARG:HG3	2.20	0.42
1:B:502:PHE:HB2	1:B:535:HIS:CE1	2.54	0.42
2:B:606:MYR:H62	2:B:606:MYR:H32	1.92	0.42
1:A:291:ALA:HA	2:A:606:MYR:H21	2.01	0.42
1:A:216:VAL:HG22	1:A:235:VAL:HG21	2.01	0.42
1:A:138:TYR:O	1:A:142:ILE:HG12	2.20	0.41
1:B:222:ARG:HD2	1:B:293:VAL:O	2.20	0.41
2:A:601:MYR:H131	2:A:601:MYR:H101	1.89	0.41
1:B:290:ILE:HG22	2:B:606:MYR:H61	2.02	0.41
1:A:291:ALA:HB1	2:A:606:MYR:H142	2.02	0.41
1:A:157:PHE:CG	3:A:607:9SC:CL1	3.11	0.41
1:A:414:LYS:HA	1:A:493:VAL:HA	2.02	0.41
1:A:22:LEU:HD13	2:A:602:MYR:H72	2.01	0.41
2:B:602:MYR:H72	2:B:602:MYR:H42	1.85	0.41
1:A:50:ALA:O	1:A:54:VAL:HG23	2.20	0.41
1:A:424:VAL:O	1:A:428:ARG:HG3	2.21	0.40
1:B:206:PHE:CZ	1:B:480:SER:HA	2.56	0.40
1:A:441:PRO:O	1:A:445:ARG:HG3	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	575/585 (98%)	556 (97%)	18 (3%)	1 (0%)	43 55

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	B	574/585 (98%)	561 (98%)	13 (2%)	0	100	100
All	All	1149/1170 (98%)	1117 (97%)	31 (3%)	1 (0%)	48	60

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	85	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	457/511 (89%)	455 (100%)	2 (0%)	84	92
1	B	457/511 (89%)	454 (99%)	3 (1%)	76	87
All	All	914/1022 (89%)	909 (100%)	5 (0%)	81	90

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	122	VAL
1	A	137	LYS
1	B	95	GLU
1	B	430	LEU
1	B	559	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	109	ASN
1	A	146	HIS
1	A	367	HIS
1	A	535	HIS
1	B	267	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	391	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

14 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	MYR	A	602	-	12,12,15	0.68	0	12,12,15	1.20	1 (8%)
2	MYR	A	606	-	15,15,15	0.62	0	15,15,15	1.10	0
3	9SC	A	607	-	33,33,33	0.28	0	45,45,45	0.56	0
2	MYR	A	603	-	12,12,15	0.70	0	12,12,15	1.09	0
2	MYR	B	602	-	12,12,15	0.69	0	12,12,15	1.11	1 (8%)
2	MYR	A	604	-	12,12,15	0.69	0	12,12,15	1.13	0
2	MYR	B	605	-	15,15,15	0.63	0	15,15,15	1.09	0
2	MYR	B	603	-	12,12,15	0.73	0	12,12,15	1.11	0
2	MYR	B	604	-	12,12,15	0.68	0	12,12,15	1.16	0
2	MYR	B	606	-	15,15,15	0.62	0	15,15,15	1.12	2 (13%)
3	9SC	B	607	-	33,33,33	0.31	0	45,45,45	0.56	0
2	MYR	A	601	-	15,15,15	0.62	0	15,15,15	1.16	2 (13%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	MYR	A	605	-	15,15,15	0.63	0	15,15,15	1.09	0
2	MYR	B	601	-	15,15,15	0.61	0	15,15,15	1.18	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	MYR	A	602	-	-	2/10/10/13	-
2	MYR	A	606	-	-	7/13/13/13	-
3	9SC	A	607	-	-	9/12/31/31	0/4/4/4
2	MYR	A	603	-	-	5/10/10/13	-
2	MYR	B	602	-	-	7/10/10/13	-
2	MYR	A	604	-	-	5/10/10/13	-
2	MYR	B	605	-	-	5/13/13/13	-
2	MYR	B	603	-	-	5/10/10/13	-
2	MYR	B	604	-	-	5/10/10/13	-
2	MYR	B	606	-	-	5/13/13/13	-
3	9SC	B	607	-	-	7/12/31/31	0/4/4/4
2	MYR	A	601	-	-	6/13/13/13	-
2	MYR	A	605	-	-	3/13/13/13	-
2	MYR	B	601	-	-	6/13/13/13	-

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	602	MYR	O2-C1-C2	2.21	121.00	114.00
2	B	601	MYR	C3-C2-C1	-2.13	108.95	114.51
2	A	601	MYR	C3-C2-C1	-2.05	109.16	114.51
2	B	606	MYR	O2-C1-C2	2.03	120.42	114.00
2	B	602	MYR	C3-C2-C1	-2.02	109.24	114.51
2	B	606	MYR	C3-C2-C1	-2.01	109.25	114.51
2	A	601	MYR	O2-C1-C2	2.01	120.34	114.00

There are no chirality outliers.

All (77) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	607	9SC	C2-C3-N1-C7
3	B	607	9SC	C2-C3-N1-C7
3	B	607	9SC	C4-C3-N1-C7
3	A	607	9SC	C4-C3-N1-C7
2	B	601	MYR	C1-C2-C3-C4
3	A	607	9SC	C12-C11-N2-C9
3	A	607	9SC	C12-C11-N2-C8
2	A	606	MYR	C11-C10-C9-C8
2	A	603	MYR	C7-C8-C9-C10
2	B	602	MYR	C3-C4-C5-C6
3	A	607	9SC	C11-C12-C13-C14
2	B	602	MYR	C7-C8-C9-C10
2	B	605	MYR	C10-C11-C12-C13
2	B	601	MYR	C5-C6-C7-C8
2	B	604	MYR	C5-C6-C7-C8
2	B	606	MYR	C10-C11-C12-C13
3	A	607	9SC	C13-C14-O1-C15
2	B	603	MYR	C1-C2-C3-C4
2	B	603	MYR	C4-C5-C6-C7
3	B	607	9SC	C13-C14-O1-C15
2	A	604	MYR	C6-C7-C8-C9
2	A	605	MYR	C3-C4-C5-C6
2	A	604	MYR	C3-C4-C5-C6
2	B	606	MYR	C11-C10-C9-C8
2	A	601	MYR	C9-C10-C11-C12
2	B	602	MYR	C6-C7-C8-C9
2	B	605	MYR	C1-C2-C3-C4
2	B	606	MYR	C11-C12-C13-C14
2	B	601	MYR	C10-C11-C12-C13
2	A	601	MYR	C11-C10-C9-C8
2	A	606	MYR	C1-C2-C3-C4
2	A	601	MYR	C3-C4-C5-C6
2	A	604	MYR	C7-C8-C9-C10
3	B	607	9SC	C2-C3-N1-C10
2	B	606	MYR	C6-C7-C8-C9
2	A	602	MYR	C7-C8-C9-C10
2	B	605	MYR	C7-C8-C9-C10
2	B	601	MYR	C11-C10-C9-C8
2	B	602	MYR	C11-C10-C9-C8
2	B	604	MYR	C1-C2-C3-C4
2	A	604	MYR	C4-C5-C6-C7
2	A	606	MYR	C7-C8-C9-C10
2	B	605	MYR	C3-C4-C5-C6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	B	601	MYR	C9-C10-C11-C12
2	A	606	MYR	C9-C10-C11-C12
2	A	605	MYR	C9-C10-C11-C12
2	B	604	MYR	C4-C5-C6-C7
3	B	607	9SC	C4-C3-N1-C10
2	A	603	MYR	C1-C2-C3-C4
2	B	606	MYR	C2-C3-C4-C5
2	A	601	MYR	C1-C2-C3-C4
2	A	606	MYR	O1-C1-C2-C3
2	A	605	MYR	C11-C10-C9-C8
2	B	604	MYR	C6-C7-C8-C9
2	A	603	MYR	O1-C1-C2-C3
2	A	606	MYR	O2-C1-C2-C3
2	B	603	MYR	O2-C1-C2-C3
2	B	601	MYR	C2-C3-C4-C5
3	A	607	9SC	C12-C13-C14-O1
2	A	603	MYR	O2-C1-C2-C3
2	A	603	MYR	C4-C5-C6-C7
2	B	603	MYR	O1-C1-C2-C3
2	B	602	MYR	O1-C1-C2-C3
2	A	601	MYR	C5-C6-C7-C8
3	B	607	9SC	C20-C15-O1-C14
2	A	601	MYR	C4-C5-C6-C7
2	A	604	MYR	C11-C10-C9-C8
2	B	602	MYR	O2-C1-C2-C3
2	B	604	MYR	C7-C8-C9-C10
3	A	607	9SC	C20-C15-O1-C14
2	A	606	MYR	C4-C5-C6-C7
2	B	602	MYR	C4-C5-C6-C7
3	B	607	9SC	C16-C15-O1-C14
2	A	602	MYR	C3-C4-C5-C6
2	B	605	MYR	C6-C7-C8-C9
3	A	607	9SC	C16-C15-O1-C14
2	B	603	MYR	C11-C10-C9-C8

There are no ring outliers.

9 monomers are involved in 15 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	602	MYR	1	0
2	A	606	MYR	3	0
3	A	607	9SC	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	602	MYR	1	0
2	A	604	MYR	1	0
2	B	605	MYR	2	0
2	B	606	MYR	2	0
3	B	607	9SC	2	0
2	A	601	MYR	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	578/585 (98%)	0.47	23 (3%)	42 44	28, 65, 107, 158	1 (0%)
1	B	578/585 (98%)	0.44	21 (3%)	46 48	36, 63, 102, 154	0
All	All	1156/1170 (98%)	0.46	44 (3%)	44 46	28, 64, 105, 158	1 (0%)

All (44) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	437	CYS	4.5
1	B	566	THR	4.1
1	B	563	ASP	4.1
1	A	278	CYS	3.9
1	B	565	GLU	3.8
1	B	138	TYR	3.6
1	A	566	THR	3.3
1	B	95	GLU	3.3
1	B	96	PRO	3.3
1	A	583	LEU	3.1
1	A	92	ALA	2.9
1	A	96	PRO	2.8
1	A	313	LYS	2.7
1	B	172	ALA	2.6
1	B	77	VAL	2.6
1	B	3	HIS	2.6
1	B	204	GLN	2.5
1	B	477	CYS	2.4
1	B	515	THR	2.4
1	B	546	ALA	2.4
1	A	478	THR	2.3
1	B	80	LEU	2.3
1	B	564	LYS	2.3
1	B	175	ALA	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	511	ALA	2.3
1	A	513	ILE	2.3
1	A	514	CYS	2.2
1	A	242[A]	HIS	2.2
1	A	471	ASP	2.2
1	A	461	CYS	2.2
1	A	95	GLU	2.2
1	A	94	GLN	2.2
1	A	108	ASP	2.2
1	A	474	THR	2.2
1	A	115	LEU	2.1
1	B	462	VAL	2.1
1	B	469	VAL	2.1
1	A	540	THR	2.1
1	B	115	LEU	2.1
1	A	438	CYS	2.1
1	B	206	PHE	2.1
1	A	443	ALA	2.0
1	B	578	ALA	2.0
1	A	301	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	9SC	B	607	30/30	0.70	0.26	71,85,90,91	30
3	9SC	A	607	30/30	0.74	0.26	80,88,90,91	30
2	MYR	B	606	16/16	0.79	0.18	61,66,72,73	0

Continued on next page...




Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	MYR	B	602	13/16	0.84	0.21	67,69,80,80	0
2	MYR	A	606	16/16	0.84	0.18	71,75,80,81	0
2	MYR	A	604	13/16	0.85	0.17	59,61,70,71	0
2	MYR	B	603	13/16	0.86	0.16	49,51,59,59	0
2	MYR	A	605	16/16	0.89	0.14	56,73,85,86	0
2	MYR	A	602	13/16	0.89	0.16	65,68,73,73	0
2	MYR	B	601	16/16	0.89	0.14	51,56,73,73	0
2	MYR	A	601	16/16	0.89	0.16	56,62,70,71	0
2	MYR	B	605	16/16	0.90	0.16	74,83,93,93	0
2	MYR	B	604	13/16	0.90	0.17	62,68,84,85	0
2	MYR	A	603	13/16	0.92	0.14	48,58,59,59	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 9SC B 607:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9SC A 607:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.