



Full wwPDB NMR Structure Validation Report ⓘ

Mar 6, 2026 – 07:17 AM UTC

PDB ID : 7VRL / pdb_00007vrl
BMRB ID : 36452
Title : Solution structure of Rbfox RRM bound to a non-cognate RNA
Authors : Yang, F.; Varani, G.
Deposited on : 2021-10-23

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 77%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823
RNA backbone	8273	777

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	7	71% 29%
2	B	102	57% 8% 33%

2 Ensemble composition and analysis

This entry contains 20 models. Model 18 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	B:114-B:148, B:158-B:190 (68)	0.33	18

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	1, 2, 5, 7, 8, 12, 17, 18, 19, 20
2	4, 9, 10, 14, 16
3	3, 11, 13, 15
Single-model clusters	6

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1855 atoms, of which 889 are hydrogens and 0 are deuteriums.

- Molecule 1 is a RNA chain called RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3').

Mol	Chain	Residues	Atoms						Trace
1	A	7	Total	C	H	N	O	P	0
			220	66	76	24	48	6	

- Molecule 2 is a protein called RNA binding protein fox-1 homolog 1.

Mol	Chain	Residues	Atoms						Trace
2	B	102	Total	C	H	N	O	S	0
			1635	511	813	156	152	3	

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
B	107	HIS	-	expression tag	UNP Q9NWB1
B	108	MET	-	expression tag	UNP Q9NWB1

4 Residue-property plots


4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1



4.2 Scores per residue for each member of the ensemble


Colouring as in section 4.1 above.

4.2.1 Score per residue for model 1

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.2 Score per residue for model 2

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')




- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.3 Score per residue for model 3

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.4 Score per residue for model 4

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')




- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.5 Score per residue for model 5

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.6 Score per residue for model 6

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')




- Molecule 2: RNA binding protein fox-1 homolog 1



4.2.7 Score per residue for model 7

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1





4.2.8 Score per residue for model 8

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 57% 43%



- Molecule 2: RNA binding protein fox-1 homolog 1

Chain B: 60% 6% 33%



4.2.9 Score per residue for model 9

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 71% 29%



- Molecule 2: RNA binding protein fox-1 homolog 1

Chain B: 50% 13% 33%




4.2.10 Score per residue for model 10

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 57% 43%




- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.11 Score per residue for model 11

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.12 Score per residue for model 12

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1



4.2.13 Score per residue for model 13

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')





- Molecule 2: RNA binding protein fox-1 homolog 1

Chain B: 58% 7% 33%



4.2.14 Score per residue for model 14

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 57% 43%



- Molecule 2: RNA binding protein fox-1 homolog 1

Chain B: 56% 9% 33%



4.2.15 Score per residue for model 15

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 57% 43%



- Molecule 2: RNA binding protein fox-1 homolog 1

Chain B: 62% 33%




4.2.16 Score per residue for model 16

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')

Chain A: 71% 29%



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.17 Score per residue for model 17

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')




- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.18 Score per residue for model 18 (medoid)

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.19 Score per residue for model 19

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')





- Molecule 2: RNA binding protein fox-1 homolog 1




4.2.20 Score per residue for model 20

- Molecule 1: RNA (5'-R(*UP*GP*CP*AP*UP*AP*U)-3')



- Molecule 2: RNA binding protein fox-1 homolog 1



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure calculation	
Amber	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1183
Number of shifts mapped to atoms	1183
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	77%

6 Model quality i

6.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.92±0.02	0±0/160 (0.0± 0.0%)	1.14±0.04	0±0/247 (0.1± 0.1%)
2	B	0.84±0.01	0±0/568 (0.0± 0.0%)	1.48±0.02	3±1/763 (0.4± 0.2%)
All	All	0.86	0/14560 (0.0%)	1.41	67/20200 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
2	B	0.0±0.0	0.7±0.7
All	All	0	13

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	158	PHE	CA-CB-CG	9.55	123.35	113.80	13	16
2	B	145	ASP	CA-CB-CG	7.94	120.54	112.60	19	1
1	A	5	U	C2'-C3'-O3'	6.71	119.57	109.50	1	2
2	B	130	ASP	CA-CB-CG	6.70	119.30	112.60	20	9
2	B	126	PHE	CA-CB-CG	6.30	120.10	113.80	4	3
2	B	160	PHE	CA-CB-CG	6.23	120.03	113.80	7	3
2	B	184	ARG	N-CA-C	6.10	120.34	113.02	6	1
2	B	180	VAL	CB-CA-C	6.05	121.22	111.29	7	4
2	B	171	ARG	NE-CZ-NH2	6.01	124.61	119.20	10	3
2	B	165	ASN	CA-CB-CG	5.89	118.49	112.60	8	19
2	B	185	LYS	CA-C-N	5.53	128.87	120.13	6	1
2	B	185	LYS	C-N-CA	5.53	128.87	120.13	6	1
2	B	186	ILE	N-CA-C	5.40	115.93	110.05	9	1
2	B	123	ASN	CA-CB-CG	5.31	117.91	112.60	6	1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	B	120	HIS	CA-CB-CG	5.24	119.04	113.80	19	1
1	A	4	A	C2'-C3'-O3'	5.19	117.29	109.50	10	1

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
2	B	158	PHE	Sidechain	8
2	B	160	PHE	Sidechain	2
2	B	118	ARG	Sidechain	2
2	B	173	ARG	Sidechain	1

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	144	76	76	1±1
2	B	556	549	549	2±2
All	All	14000	12500	12500	36

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:6:A:C2	2:B:160:PHE:CZ	0.56	2.94	4	1
2:B:144:LEU:H	2:B:144:LEU:HD22	0.54	1.62	17	5
1:A:6:A:C2	2:B:160:PHE:CE2	0.51	2.99	4	1
1:A:4:A:H1'	2:B:158:PHE:CE1	0.49	2.42	13	2
1:A:6:A:C2	2:B:160:PHE:CE1	0.48	3.01	14	2
2:B:144:LEU:HD23	2:B:162:THR:HG22	0.46	1.88	17	3
2:B:144:LEU:HD21	2:B:164:GLU:HA	0.45	1.87	12	5
2:B:144:LEU:HD22	2:B:144:LEU:N	0.43	2.28	11	4
2:B:134:ARG:HG2	2:B:146:VAL:HG21	0.42	1.91	11	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
2:B:134:ARG:CG	2:B:146:VAL:HG21	0.42	2.43	10	2
1:A:6:A:N3	2:B:160:PHE:CE1	0.42	2.87	13	1
2:B:144:LEU:HD21	2:B:164:GLU:N	0.42	2.30	3	2
1:A:6:A:N3	2:B:160:PHE:CE2	0.41	2.87	20	2
1:A:4:A:H1'	2:B:158:PHE:CE2	0.41	2.51	1	1
1:A:5:U:O3'	1:A:6:A:H4'	0.41	2.16	1	2
2:B:144:LEU:HD21	2:B:164:GLU:CA	0.41	2.46	12	1
2:B:123:ASN:HD22	2:B:186:ILE:HA	0.40	1.77	6	1

6.3 Torsion angles

6.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
2	B	68/102 (67%)	62±1 (91±2%)	6±1 (8±2%)	0±1 (1±1%)	26	74
All	All	1360/2040 (67%)	1242 (91%)	111 (8%)	7 (1%)	26	74

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	115	GLN	3
2	B	185	LYS	2
2	B	182	GLU	1
2	B	190	ASN	1

6.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	60/89 (67%)	51±2 (85±3%)	9±2 (15±3%)	5	43
All	All	1200/1780 (67%)	1023 (85%)	177 (15%)	5	43

All 28 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
2	B	165	ASN	20
2	B	171	ARG	18
2	B	180	VAL	18
2	B	163	PHE	15
2	B	161	VAL	12
2	B	120	HIS	10
2	B	121	VAL	10
2	B	181	VAL	10
2	B	127	ARG	8
2	B	182	GLU	8
2	B	158	PHE	6
2	B	144	LEU	6
2	B	173	ARG	5
2	B	122	SER	4
2	B	130	ASP	3
2	B	128	PHE	3
2	B	118	ARG	3
2	B	126	PHE	3
2	B	190	ASN	3
2	B	115	GLN	2
2	B	148	ILE	2
2	B	136	MET	2
2	B	123	ASN	1
2	B	129	ARG	1
2	B	188	VAL	1
2	B	146	VAL	1
2	B	134	ARG	1
2	B	145	ASP	1

6.3.3 RNA ⓘ

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers	Suiteness
1	A	6/7 (86%)	2±0 (33±0%)	0±1 (8±12%)	0.20±0.00
All	All	124/140 (89%)	40 (32%)	10 (8%)	0.20

The overall RNA backbone suiteness is 0.11.

All unique RNA backbone outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	3	C	20
1	A	7	U	20

All unique RNA pucker outliers are listed below:

Mol	Chain	Res	Type	Models (Total)
1	A	3	C	5
1	A	1	U	4
1	A	5	U	1

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

The completeness of assignment taking into account all chemical shift lists is 77% for the well-defined parts and 76% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: */Users/fan/PROJECTS/NMRdata/HIT600/Struct_calc/FoxRNA1/CS.str*

7.1.1 Bookkeeping

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1183
Number of shifts mapped to atoms	1183
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	3

7.1.2 Chemical shift referencing

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	102	-0.44 ± 0.17	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	94	0.14 ± 0.12	None needed (< 0.5 ppm)
$^{13}\text{C}'$	95	-0.17 ± 0.32	None needed (< 0.5 ppm)
^{15}N	97	0.05 ± 0.39	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 77%, i.e. 845 atoms were assigned a chemical shift out of a possible 1103. 0 out of 10 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	335/339 (99%)	137/138 (99%)	133/136 (98%)	65/65 (100%)
Sidechain	433/552 (78%)	284/353 (80%)	139/167 (83%)	10/32 (31%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	38/84 (45%)	19/43 (44%)	19/39 (49%)	0/2 (0%)
Sugar	29/77 (38%)	29/42 (69%)	0/35 (0%)	0/0 (—%)
Base	10/51 (20%)	10/30 (33%)	0/13 (0%)	0/8 (0%)
Overall	845/1103 (77%)	479/606 (79%)	291/390 (75%)	75/107 (70%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 76%, i.e. 1180 atoms were assigned a chemical shift out of a possible 1556. 0 out of 12 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	497/510 (97%)	203/208 (98%)	197/204 (97%)	97/98 (99%)
Sidechain	594/808 (74%)	387/516 (75%)	193/244 (79%)	14/48 (29%)
Aromatic	50/110 (45%)	25/56 (45%)	25/51 (49%)	0/3 (0%)
Sugar	29/77 (38%)	29/42 (69%)	0/35 (0%)	0/0 (—%)
Base	10/51 (20%)	10/30 (33%)	0/13 (0%)	0/8 (0%)
Overall	1180/1556 (76%)	654/852 (77%)	415/547 (76%)	111/157 (71%)

7.1.4 Statistically unusual chemical shifts ⓘ

The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	B	132	ASP	HB2	0.93	1.41 – 4.01	-6.8
1	A	3	C	H3'	3.31	3.45 – 5.36	-5.7
1	B	125	PRO	HD2	1.72	1.93 – 5.38	-5.6

7.1.5 Random Coil Index (RCI) plots ⓘ

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain B:

