



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 09:30 AM UTC

PDB ID : 1VSC / pdb_00001vsc
Title : VCAM-1
Authors : Wang, J.; Stehle, T.; Osborn, L.
Deposited on : 1995-04-27
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	196	9% 56% 36% 7% .
1	B	196	4% 60% 35% . .

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3359 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called VASCULAR CELL ADHESION MOLECULE-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	196	Total	C	N	O	S	0	0	0
			1528	961	250	308	9			
1	B	196	Total	C	N	O	S	0	0	0
			1528	961	250	308	9			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	142	Total	O	0	0
			142	142		
2	B	161	Total	O	0	0
			161	161		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: VASCULAR CELL ADHESION MOLECULE-1



• Molecule 1: VASCULAR CELL ADHESION MOLECULE-1



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	122.09Å 48.86Å 73.44Å 90.00° 117.42° 90.00°	Depositor
Resolution (Å)	15.00 – 1.90 15.00 – 1.90	Depositor EDS
% Data completeness (in resolution range)	82.9 (15.00-1.90) 78.6 (15.00-1.90)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.30 (at 1.88Å)	Xtriage
Refinement program	X-PLOR 3.1	Depositor
R, R_{free}	0.204 , 0.289 0.218 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	20.3	Xtriage
Anisotropy	0.703	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 62.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3359	wwPDB-VP
Average B, all atoms (Å ²)	31.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 37.01 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 4.5782e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.88	0/1556	1.30	20/2109 (0.9%)
1	B	0.95	0/1556	1.26	12/2109 (0.6%)
All	All	0.91	0/3112	1.28	32/4218 (0.8%)

There are no bond length outliers.

All (32) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	148	SER	N-CA-C	8.82	120.59	110.97
1	B	144	ALA	N-CA-C	8.18	128.22	110.80
1	B	68	SER	N-CA-C	-7.52	96.47	108.73
1	B	76	GLU	CA-C-N	-6.66	112.22	122.60
1	B	76	GLU	C-N-CA	-6.66	112.22	122.60
1	A	186	VAL	N-CA-C	6.64	118.36	108.46
1	A	177	ILE	CB-CA-C	-6.59	103.97	111.55
1	A	132	ASP	N-CA-C	-6.47	105.35	112.72
1	B	37	THR	N-CA-C	-6.32	100.63	110.42
1	B	153	SER	N-CA-C	6.13	118.73	109.23
1	B	143	ASP	N-CA-C	-6.09	97.83	110.80
1	A	169	LEU	N-CA-C	-6.08	99.75	109.96
1	A	133	HIS	N-CA-C	6.07	118.73	107.99
1	A	124	LEU	N-CA-C	6.05	118.82	109.07
1	A	76	GLU	CA-C-N	-5.78	113.40	122.65
1	A	76	GLU	C-N-CA	-5.78	113.40	122.65
1	A	162	ILE	N-CA-C	5.76	119.11	111.17
1	A	61	VAL	N-CA-C	-5.74	101.41	109.21
1	B	8	GLU	N-CA-C	5.66	117.90	111.11
1	B	195	VAL	N-CA-C	5.64	116.88	109.21
1	A	26	THR	N-CA-C	-5.60	100.65	109.50
1	B	187	ARG	N-CA-C	-5.59	99.90	109.24
1	A	135	MET	N-CA-C	-5.59	106.13	113.12
1	B	137	SER	N-CA-C	-5.45	100.90	109.50
1	A	68	SER	N-CA-C	-5.44	99.86	108.73

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	67	HIS	CB-CA-C	-5.39	100.60	111.17
1	B	130	LYS	N-CA-C	-5.31	98.72	108.02
1	A	78	ARG	N-CA-C	5.30	118.46	109.76
1	A	126	ILE	CB-CA-C	-5.26	102.99	110.83
1	A	93	LYS	N-CA-C	5.11	117.16	109.23
1	A	111	VAL	CB-CA-C	-5.11	103.98	110.98
1	A	114	SER	N-CA-C	5.08	117.85	109.72

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1528	0	1518	52	0
1	B	1528	0	1516	56	0
2	A	142	0	0	9	0
2	B	161	0	0	17	0
All	All	3359	0	3034	108	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (108) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:GLN:HB2	1:A:154:LEU:HD13	1.50	0.93
1:B:143:ASP:HB2	1:B:146:ARG:HD2	1.49	0.93
1:A:170:VAL:HG22	1:A:190:VAL:HG12	1.59	0.84
1:A:100:SER:OG	1:A:110:THR:HB	1.80	0.81
1:B:171:CYS:SG	2:B:673:HOH:O	2.37	0.81
1:B:113:CYS:SG	2:B:673:HOH:O	2.41	0.78
1:B:147:LYS:HB2	2:B:686:HOH:O	1.84	0.78
1:A:99:LEU:HG	1:A:191:LYS:HE3	1.71	0.72
1:A:145:ASP:HB3	1:A:148:SER:HB2	1.72	0.72

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:123:ARG:HD2	1:A:176:HIS:O	1.90	0.71
1:A:17:ASP:O	1:A:61:VAL:HG12	1.89	0.71
1:B:180:MET:HE3	1:B:180:MET:HA	1.74	0.69
1:B:168:VAL:HB	2:B:694:HOH:O	1.93	0.67
1:A:118:VAL:HG11	1:A:124:LEU:HD22	1.78	0.66
1:B:102:PRO:HB2	1:B:196:ASP:HB3	1.76	0.66
1:A:11:TYR:CE2	1:A:19:VAL:HG13	2.31	0.66
1:A:36:ARG:HG2	2:A:494:HOH:O	1.96	0.64
1:A:38:GLN:HG2	1:A:39:ILE:HG23	1.80	0.63
1:B:13:ALA:O	1:B:88:ILE:HA	1.98	0.63
1:A:136:LYS:HG2	1:A:154:LEU:HD11	1.81	0.62
1:B:163:GLU:O	1:B:163:GLU:HG2	2.01	0.61
1:B:128:LEU:HD23	1:B:156:VAL:HG11	1.82	0.61
1:B:14:GLN:NE2	2:B:445:HOH:O	2.34	0.60
1:A:103:LEU:HD23	1:A:194:GLN:O	2.02	0.58
1:A:148:SER:O	1:A:149:LEU:HB2	2.03	0.57
1:A:165:ILE:HG21	1:A:195:VAL:HB	1.85	0.57
1:A:136:LYS:HE3	2:A:625:HOH:O	2.04	0.56
1:B:76:GLU:HB3	2:B:690:HOH:O	2.04	0.56
1:B:176:HIS:HD2	2:B:606:HOH:O	1.88	0.56
1:B:46:LYS:O	1:B:46:LYS:HG3	2.05	0.55
1:B:6:THR:HG1	1:B:22:THR:HG1	1.55	0.55
1:B:36:ARG:NH1	2:B:506:HOH:O	2.39	0.54
1:B:46:LYS:HE2	2:B:470:HOH:O	2.08	0.54
1:B:90:SER:O	1:B:118:VAL:HA	2.08	0.53
1:B:142:GLU:O	1:B:144:ALA:N	2.42	0.53
1:B:63:PHE:CE1	1:B:147:LYS:O	2.61	0.53
1:B:118:VAL:O	1:B:121:PHE:CE1	2.61	0.53
1:B:138:GLN:HB2	1:B:154:LEU:HD13	1.91	0.52
1:B:136:LYS:HD3	1:B:154:LEU:HD11	1.91	0.52
1:A:167:LYS:O	1:A:193:LEU:HD12	2.09	0.52
1:A:70:LEU:HD11	2:A:447:HOH:O	2.10	0.52
1:A:179:GLU:HG2	2:A:531:HOH:O	2.10	0.52
1:A:195:VAL:CG1	1:A:196:ASP:N	2.73	0.52
1:B:28:CYS:SG	1:B:75:CYS:CB	2.99	0.51
1:A:172:ARG:NE	1:A:188:GLN:OE1	2.43	0.51
1:A:195:VAL:HA	2:A:671:HOH:O	2.11	0.50
1:B:12:LEU:HD13	1:B:87:GLU:HB3	1.93	0.50
1:B:5:THR:HB	1:B:84:ILE:HD11	1.92	0.50
1:A:99:LEU:CG	1:A:191:LYS:HE3	2.41	0.50
1:B:191:LYS:HE2	2:B:620:HOH:O	2.11	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:PRO:N	1:A:159:THR:HG22	2.27	0.50
1:B:147:LYS:NZ	2:B:580:HOH:O	2.45	0.50
1:B:165:ILE:HG23	1:B:195:VAL:HG23	1.94	0.50
1:B:95:PRO:HB2	1:B:171:CYS:SG	2.51	0.50
1:B:168:VAL:HG13	1:B:190:VAL:HG13	1.93	0.50
1:B:123:ARG:HD3	2:B:696:HOH:O	2.11	0.49
1:A:14:GLN:NE2	2:A:427:HOH:O	2.45	0.49
1:B:103:LEU:HD12	1:B:103:LEU:N	2.27	0.49
1:B:154:LEU:HD12	1:B:155:GLU:N	2.28	0.49
1:A:132:ASP:OD1	1:A:132:ASP:N	2.45	0.49
1:B:154:LEU:HD12	1:B:155:GLU:H	1.78	0.49
1:B:16:GLY:HA2	1:B:60:PRO:HB2	1.94	0.49
1:B:118:VAL:HG21	1:B:124:LEU:HD13	1.94	0.48
1:B:102:PRO:HB2	1:B:196:ASP:CB	2.41	0.48
1:A:36:ARG:HD2	1:A:40:ASP:OD1	2.14	0.48
1:B:170:VAL:HG22	1:B:190:VAL:HG22	1.96	0.48
1:A:82:LYS:NZ	2:A:518:HOH:O	2.46	0.47
1:A:135:MET:HE2	1:A:156:VAL:HG11	1.96	0.47
1:B:146:ARG:HB2	1:B:146:ARG:CZ	2.45	0.47
1:A:95:PRO:HB2	1:A:171:CYS:SG	2.54	0.47
1:A:101:GLY:O	1:A:109:ILE:HD11	2.14	0.47
1:B:176:HIS:CD2	2:B:606:HOH:O	2.64	0.47
1:A:152:LYS:HD3	1:A:152:LYS:HA	1.72	0.46
1:A:169:LEU:HA	1:A:169:LEU:HD23	1.64	0.46
1:A:117:ASP:HA	1:A:149:LEU:HD22	1.98	0.46
1:B:136:LYS:CD	1:B:154:LEU:HD11	2.47	0.45
1:B:109:ILE:HG23	2:B:493:HOH:O	2.16	0.45
1:A:103:LEU:HD21	1:A:195:VAL:HG23	1.99	0.45
1:A:8:GLU:OE1	1:A:82:LYS:NZ	2.50	0.44
1:B:105:ALA:HA	2:B:598:HOH:O	2.17	0.44
1:A:58:MET:HE3	1:A:58:MET:HB2	1.91	0.44
1:B:70:LEU:HD23	1:B:83:GLY:CA	2.47	0.44
1:B:121:PHE:HZ	1:B:148:SER:O	2.01	0.44
1:A:101:GLY:HA3	1:A:102:PRO:HD2	1.75	0.43
1:B:152:LYS:HD3	1:B:152:LYS:HA	1.73	0.43
1:A:172:ARG:HG3	1:A:188:GLN:OE1	2.17	0.43
1:A:183:VAL:HG21	2:A:555:HOH:O	2.18	0.43
1:B:38:GLN:OE1	1:B:67:HIS:HB3	2.17	0.43
1:B:70:LEU:HD23	1:B:83:GLY:HA3	2.00	0.43
1:B:100:SER:HB2	1:B:110:THR:H	1.83	0.43
1:A:35:TRP:HB2	1:A:43:LEU:HD11	2.00	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:146:ARG:HA	1:A:146:ARG:HD2	1.90	0.42
1:A:165:ILE:CG2	1:A:195:VAL:HB	2.48	0.42
1:B:165:ILE:HG12	1:B:195:VAL:CG2	2.50	0.42
1:B:78:ARG:HH11	1:B:78:ARG:HG3	1.85	0.42
1:A:185:THR:HA	2:A:618:HOH:O	2.20	0.41
1:B:123:ARG:NH1	2:B:652:HOH:O	2.52	0.41
1:A:18:SER:HB2	1:A:59:ASN:OD1	2.20	0.41
1:B:168:VAL:CG1	1:B:190:VAL:HG13	2.50	0.41
1:A:59:ASN:HA	1:A:60:PRO:HA	1.85	0.41
1:A:177:ILE:H	1:A:177:ILE:HG13	1.60	0.41
1:A:89:TYR:HB3	1:A:177:ILE:HG21	2.02	0.41
1:A:168:VAL:CG1	1:A:190:VAL:HB	2.51	0.41
1:B:20:SER:HB3	2:B:677:HOH:O	2.20	0.41
1:A:43:LEU:HD13	1:A:56:LEU:HD13	2.03	0.41
1:A:80:LEU:HD23	1:A:80:LEU:HA	1.92	0.40
1:B:146:ARG:NH2	1:B:150:GLU:HG3	2.36	0.40
1:A:123:ARG:HD3	1:A:123:ARG:HA	1.76	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	194/196 (99%)	184 (95%)	10 (5%)	0	100	100
1	B	194/196 (99%)	185 (95%)	8 (4%)	1 (0%)	24	16
All	All	388/392 (99%)	369 (95%)	18 (5%)	1 (0%)	36	29

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	144	ALA

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	179/179 (100%)	158 (88%)	21 (12%)	5	2
1	B	179/179 (100%)	163 (91%)	16 (9%)	9	4
All	All	358/358 (100%)	321 (90%)	37 (10%)	7	2

All (37) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	8	GLU
1	A	61	VAL
1	A	85	GLN
1	A	96	GLU
1	A	103	LEU
1	A	107	LYS
1	A	115	VAL
1	A	122	ASP
1	A	132	ASP
1	A	141	LEU
1	A	143	ASP
1	A	145	ASP
1	A	149	LEU
1	A	156	VAL
1	A	158	PHE
1	A	171	CYS
1	A	176	HIS
1	A	177	ILE
1	A	180	MET
1	A	193	LEU
1	A	195	VAL
1	B	15	ILE
1	B	52	THR
1	B	85	GLN
1	B	92	PRO
1	B	100	SER
1	B	115	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	129	LEU
1	B	139	GLU
1	B	141	LEU
1	B	143	ASP
1	B	162	ILE
1	B	163	GLU
1	B	171	CYS
1	B	174	LYS
1	B	190	VAL
1	B	196	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	14	GLN
1	B	176	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	196/196 (100%)	0.45	18 (9%) 14 15	11, 27, 72, 79	0
1	B	196/196 (100%)	0.17	7 (3%) 46 49	8, 24, 55, 77	0
All	All	392/392 (100%)	0.31	25 (6%) 25 27	8, 26, 68, 79	0

All (25) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	195	VAL	6.5
1	A	103	LEU	3.6
1	B	144	ALA	3.5
1	A	144	ALA	3.2
1	A	106	GLY	3.1
1	A	162	ILE	3.0
1	A	109	ILE	2.9
1	A	196	ASP	2.8
1	B	143	ASP	2.8
1	B	195	VAL	2.8
1	B	145	ASP	2.6
1	A	141	LEU	2.5
1	B	146	ARG	2.5
1	A	158	PHE	2.5
1	A	131	GLY	2.5
1	A	166	GLY	2.5
1	A	165	ILE	2.5
1	A	161	VAL	2.5
1	B	142	GLU	2.4
1	B	131	GLY	2.3
1	A	32	PHE	2.3
1	A	77	SER	2.3
1	A	194	GLN	2.2
1	A	78	ARG	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	177	ILE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.