



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 09:00 PM UTC

PDB ID : 1VSE / pdb_00001vse
Title : ASV INTEGRASE CORE DOMAIN WITH MG(II) COFACTOR AND
HEPES LIGAND, LOW MG CONCENTRATION FORM
Authors : Bujacz, G.; Jaskolski, M.; Alexandratos, J.; Wlodawer, A.
Deposited on : 1995-11-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	152	 59% 30% 6% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1263 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called INTEGRASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1124	709	210	200	5			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	ALA	VAL	conflict	UNP P03354
A	166	LYS	ARG	conflict	UNP P03354

- Molecule 2 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: $C_8H_{18}N_2O_4S$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	124	Total 124	O 124	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

• Molecule 1: INTEGRASE

Chain A: 



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	66.25Å 66.25Å 81.58Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	8.00 – 2.20	Depositor
% Data completeness (in resolution range)	86.1 (8.00-2.20)	Depositor
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	PROLSQ	Depositor
R, R_{free}	0.138 , 0.201	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	1263	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.92	0/1149	2.21	54/1558 (3.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (54) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	158	ARG	CD-NE-CZ	12.30	141.62	124.40
1	A	70	ARG	NE-CZ-NH2	-8.63	111.43	119.20
1	A	161	ARG	CB-CG-CD	8.56	130.98	111.30
1	A	95	ARG	NE-CZ-NH1	-8.53	112.97	121.50
1	A	148	GLY	CA-C-O	8.16	126.44	118.77
1	A	88	ILE	CB-CA-C	-7.72	99.07	110.81
1	A	131	THR	CA-C-O	-7.67	112.29	120.42
1	A	149	ASN	CA-CB-CG	7.49	120.08	112.60
1	A	67	LEU	CA-C-N	7.45	131.58	122.42
1	A	67	LEU	C-N-CA	7.45	131.58	122.42
1	A	160	ASN	OD1-CG-ND2	-7.43	115.17	122.60
1	A	113	GLY	CA-C-O	-7.40	114.94	122.56
1	A	160	ASN	CA-CB-CG	7.24	119.83	112.60
1	A	92	GLN	N-CA-CB	-7.18	98.92	110.63
1	A	179	ARG	NE-CZ-NH2	-7.16	112.76	119.20
1	A	144	THR	N-CA-C	-6.89	104.19	112.59

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	114	ARG	NE-CZ-NH1	6.87	128.37	121.50
1	A	148	GLY	N-CA-C	-6.71	105.00	116.01
1	A	178	LYS	CB-CA-C	-6.65	106.83	116.54
1	A	100	ALA	CA-C-N	6.38	129.35	120.29
1	A	100	ALA	C-N-CA	6.38	129.35	120.29
1	A	90	VAL	O-C-N	-6.35	116.31	123.10
1	A	83	THR	CA-C-O	-6.33	111.83	119.49
1	A	122	ASN	CA-CB-CG	-6.16	106.44	112.60
1	A	179	ARG	NE-CZ-NH1	6.15	127.65	121.50
1	A	100	ALA	O-C-N	-6.11	115.07	122.22
1	A	82	ASP	O-C-N	-6.09	115.62	122.93
1	A	193	MET	CA-C-N	6.00	128.24	120.44
1	A	193	MET	C-N-CA	6.00	128.24	120.44
1	A	134	TRP	CA-C-N	5.96	128.75	120.29
1	A	134	TRP	C-N-CA	5.96	128.75	120.29
1	A	70	ARG	NH1-CZ-NH2	5.81	126.85	119.30
1	A	128	SER	CA-C-N	5.77	127.94	120.44
1	A	128	SER	C-N-CA	5.77	127.94	120.44
1	A	95	ARG	NH1-CZ-NH2	5.71	126.73	119.30
1	A	185	GLN	N-CA-C	5.70	117.17	111.07
1	A	114	ARG	NE-CZ-NH2	-5.61	114.15	119.20
1	A	55	LEU	CA-C-O	5.57	125.95	119.32
1	A	96	VAL	CB-CA-C	5.57	119.44	111.81
1	A	103	HIS	N-CA-CB	5.48	118.27	110.16
1	A	162	LEU	CA-C-O	5.46	126.33	120.55
1	A	198	HIS	O-C-N	5.42	130.27	123.23
1	A	114	ARG	CD-NE-CZ	-5.41	116.83	124.40
1	A	161	ARG	CG-CD-NE	5.40	123.87	112.00
1	A	143	THR	O-C-N	-5.30	117.12	123.27
1	A	191	LYS	CA-C-O	-5.29	114.81	120.42
1	A	185	GLN	OE1-CD-NE2	5.25	127.85	122.60
1	A	113	GLY	N-CA-C	-5.21	104.87	112.17
1	A	135	LEU	CA-C-O	-5.18	114.93	120.42
1	A	148	GLY	CA-C-N	5.12	128.88	120.63
1	A	148	GLY	C-N-CA	5.12	128.88	120.63
1	A	89	VAL	CA-C-O	-5.10	114.95	120.36
1	A	160	ASN	CB-CG-ND2	5.09	124.04	116.40
1	A	178	LYS	N-CA-CB	-5.02	106.92	111.59

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	114	ARG	Sidechain
1	A	70	ARG	Sidechain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1124	0	1139	35	0
2	A	15	0	18	0	0
3	A	124	0	0	0	0
All	All	1263	0	1157	35	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 15.

All (35) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:ARG:HH12	1:A:165:ASP:HB2	1.01	1.10
1:A:161:ARG:NH1	1:A:165:ASP:HB2	1.70	1.06
1:A:161:ARG:HH12	1:A:165:ASP:CB	1.81	0.93
1:A:146:ILE:O	1:A:149:ASN:HB2	1.74	0.88
1:A:146:ILE:CG2	1:A:147:PRO:HD2	2.24	0.68
1:A:146:ILE:HD13	1:A:149:ASN:ND2	2.12	0.65
1:A:146:ILE:HG23	1:A:147:PRO:HD2	1.82	0.61
1:A:119:LYS:HD2	1:A:153:GLN:HB3	1.83	0.60
1:A:132:ARG:HG3	1:A:132:ARG:NH1	2.17	0.60
1:A:146:ILE:HD13	1:A:149:ASN:HD22	1.69	0.57
1:A:57:PRO:HB2	1:A:84:ALA:HB1	1.87	0.55
1:A:161:ARG:HH22	1:A:165:ASP:N	2.05	0.55
1:A:74:ARG:O	1:A:92:GLN:HG2	2.07	0.54
1:A:151:GLN:HG2	1:A:157:GLU:OE1	2.08	0.54
1:A:90:VAL:HG21	1:A:163:LEU:HD21	1.90	0.52
1:A:85:SER:O	1:A:86:SER:CB	2.58	0.51
1:A:57:PRO:HB2	1:A:84:ALA:CB	2.43	0.49
1:A:120:THR:O	1:A:144:THR:HA	2.13	0.49
1:A:146:ILE:HD12	1:A:146:ILE:N	2.29	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:62:GLN:NE2	1:A:156:VAL:H	2.11	0.47
1:A:161:ARG:NH2	1:A:165:ASP:N	2.62	0.47
1:A:161:ARG:HH22	1:A:165:ASP:CA	2.28	0.47
1:A:146:ILE:HD12	1:A:146:ILE:H	1.80	0.46
1:A:146:ILE:HG22	1:A:147:PRO:HD2	1.96	0.46
1:A:184:LYS:HA	1:A:184:LYS:HD3	1.66	0.44
1:A:114:ARG:NH1	1:A:138:TRP:O	2.51	0.44
1:A:151:GLN:C	1:A:153:GLN:H	2.26	0.43
1:A:151:GLN:CD	1:A:153:GLN:CG	2.92	0.43
1:A:132:ARG:NH1	1:A:132:ARG:CG	2.81	0.43
1:A:148:GLY:C	1:A:150:SER:H	2.27	0.42
1:A:146:ILE:CD1	1:A:149:ASN:ND2	2.83	0.42
1:A:168:ARG:HG2	1:A:168:ARG:HH11	1.86	0.41
1:A:68:GLU:OE1	1:A:164:LYS:NZ	2.50	0.41
1:A:148:GLY:C	1:A:150:SER:N	2.79	0.41
1:A:96:VAL:O	1:A:96:VAL:HG13	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/152 (95%)	137 (95%)	7 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	83/122 (68%)	80 (96%)	3 (4%)	31	42

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	86	SER
1	A	157	GLU
1	A	161	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	92	GLN
1	A	185	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EPE	A	261	-	15,15,15	1.67	2 (13%)	19,20,20	2.24	5 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EPE	A	261	-	-	4/9/19/19	0/1/1/1

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	261	EPE	O3S-S	4.96	1.65	1.47
2	A	261	EPE	C10-S	3.11	1.82	1.77

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	261	EPE	O2S-S-C10	6.86	117.09	106.73
2	A	261	EPE	O3S-S-C10	-4.20	97.78	106.00
2	A	261	EPE	C6-C5-N4	2.70	116.09	110.65
2	A	261	EPE	C7-N4-C3	2.53	117.99	111.24
2	A	261	EPE	O8-C8-C7	2.46	121.37	111.22

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	261	EPE	C10-C9-N1-C6
2	A	261	EPE	C9-C10-S-O2S
2	A	261	EPE	N4-C7-C8-O8
2	A	261	EPE	C10-C9-N1-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates ⓘ

EDS was not executed - this section is therefore empty.

6.4 Ligands ⓘ

EDS was not executed - this section is therefore empty.

6.5 Other polymers ⓘ

EDS was not executed - this section is therefore empty.