



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 03:49 AM UTC

PDB ID : 1VSI / pdb_00001vsi
Title : ASV INTEGRASE CORE DOMAIN WITH CA(II) COFACTOR
Authors : Bujacz, G.; Alexandratos, J.; Wlodawer, A.
Deposited on : 1997-03-04
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	152	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1242 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTEGRASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	146	Total	C	N	O	S	0	0	0
			1127	711	210	201	5			


There are 3 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	101	ALA	VAL	variant	UNP P03354
A	125	OCY	CYS	modified residue	UNP P03354
A	166	LYS	ARG	variant	UNP P03354

- Molecule 2 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Ca	0	0
			1	1		

- Molecule 3 is 4-(2-HYDROXYETHYL)-1-PIPERAZINE ETHANESULFONIC ACID (CCD ID: EPE) (formula: C₈H₁₈N₂O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	S	0	0
			15	8	2	4	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	99	Total	O	0	0
			99	99		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INTEGRASE



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	66.15Å 66.15Å 80.96Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	10.00 – 2.20 10.00 – 2.20	Depositor EDS
% Data completeness (in resolution range)	95.5 (10.00-2.20) 91.8 (10.00-2.20)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.11	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtriage
Refinement program	PROFFT	Depositor
R, R_{free}	0.171 , (Not available) 0.166 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.3	Xtriage
Anisotropy	0.018	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 81.5	EDS
L-test for twinning ¹	$\langle L \rangle = 0.37$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	1242	wwPDB-VP
Average B, all atoms (Å ²)	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.51% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: EPE, OCY, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.98	0/1142	2.43	69/1547 (4.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (69) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	149	ASN	CA-CB-CG	14.37	126.97	112.60
1	A	70	ARG	NE-CZ-NH2	-14.30	106.33	119.20
1	A	70	ARG	NH1-CZ-NH2	10.51	132.96	119.30
1	A	113	GLY	CA-C-O	-10.03	113.35	122.57
1	A	131	THR	CA-C-O	-9.63	110.34	120.55
1	A	89	VAL	CA-C-O	-9.35	110.45	120.36
1	A	199	PHE	CA-CB-CG	9.16	122.96	113.80
1	A	90	VAL	CB-CA-C	-9.15	96.41	110.50
1	A	62	GLN	CA-C-O	-8.70	110.66	120.32
1	A	72	ALA	CA-C-O	-8.45	111.49	120.70
1	A	127	THR	CA-CB-OG1	-8.11	97.44	109.60
1	A	161	ARG	CA-CB-CG	7.78	129.67	114.10
1	A	131	THR	O-C-N	7.58	130.16	122.12
1	A	168	ARG	CD-NE-CZ	7.57	135.00	124.40
1	A	165	ASP	CA-CB-CG	-7.46	105.14	112.60
1	A	93	HIS	O-C-N	-7.38	114.24	123.17

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	81	VAL	CA-C-O	7.37	128.29	120.48
1	A	81	VAL	O-C-N	-7.23	115.59	123.18
1	A	103	HIS	O-C-N	7.18	130.33	122.15
1	A	92	GLN	CB-CA-C	-7.00	96.57	109.37
1	A	96	VAL	CA-C-O	-6.88	112.30	120.76
1	A	102	GLN	OE1-CD-NE2	-6.75	115.85	122.60
1	A	116	LYS	N-CA-C	-6.58	104.27	112.90
1	A	126	PHE	CA-C-N	-6.45	112.54	122.60
1	A	126	PHE	C-N-CA	-6.45	112.54	122.60
1	A	114	ARG	CA-CB-CG	6.42	126.94	114.10
1	A	158	ARG	N-CA-C	-6.40	104.39	111.36
1	A	151	GLN	CB-CG-CD	6.32	123.34	112.60
1	A	100	ALA	CA-C-N	6.32	129.26	120.29
1	A	100	ALA	C-N-CA	6.32	129.26	120.29
1	A	169	VAL	CA-C-O	-6.29	114.18	120.85
1	A	103	HIS	CA-CB-CG	-6.29	107.51	113.80
1	A	65	PHE	CA-C-O	-6.28	114.06	121.16
1	A	144	THR	N-CA-C	-6.23	104.31	112.41
1	A	184	LYS	CA-C-O	-6.19	112.33	119.14
1	A	132	ARG	CA-CB-CG	-6.09	101.92	114.10
1	A	182	THR	CA-C-O	6.05	126.97	120.55
1	A	102	GLN	CA-C-O	6.01	126.79	120.42
1	A	154	ALA	CA-C-O	-5.96	113.37	120.10
1	A	129	LYS	CA-C-N	5.73	127.89	120.44
1	A	129	LYS	C-N-CA	5.73	127.89	120.44
1	A	100	ALA	O-C-N	-5.67	116.11	122.12
1	A	175	GLY	CA-C-N	-5.60	114.85	123.14
1	A	175	GLY	C-N-CA	-5.60	114.85	123.14
1	A	164	LYS	CA-CB-CG	-5.51	103.08	114.10
1	A	113	GLY	N-CA-C	-5.50	104.67	112.81
1	A	101	ALA	CA-C-O	-5.49	114.60	120.42
1	A	158	ARG	CB-CG-CD	5.48	123.90	111.30
1	A	72	ALA	O-C-N	5.47	127.55	121.53
1	A	70	ARG	CA-CB-CG	-5.47	103.16	114.10
1	A	62	GLN	N-CA-C	-5.46	99.52	108.76
1	A	86	SER	CA-C-O	-5.46	111.89	119.16
1	A	75	SER	N-CA-C	5.46	119.94	113.28
1	A	193	MET	CA-C-O	5.46	126.20	120.42
1	A	114	ARG	N-CA-CB	-5.40	101.21	110.01
1	A	148	GLY	CA-C-N	5.34	129.68	121.19
1	A	148	GLY	C-N-CA	5.34	129.68	121.19
1	A	172	GLU	CA-C-O	-5.34	113.83	119.97

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	SER	O-C-N	-5.34	116.51	122.75
1	A	192	ALA	CB-CA-C	-5.28	102.58	110.88
1	A	155	MET	CB-CA-C	-5.23	101.95	110.85
1	A	85	SER	CA-C-N	-5.19	115.89	125.02
1	A	85	SER	C-N-CA	-5.19	115.89	125.02
1	A	129	LYS	N-CA-C	5.15	117.56	111.33
1	A	137	ARG	CB-CA-C	-5.11	101.17	110.63
1	A	84	ALA	N-CA-CB	5.09	117.67	110.13
1	A	61	TRP	CA-CB-CG	5.01	123.12	113.60
1	A	184	LYS	CA-C-N	5.01	126.99	120.28
1	A	184	LYS	C-N-CA	5.01	126.99	120.28

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	70	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1127	0	1143	40	0
2	A	1	0	0	0	0
3	A	15	0	18	1	0
4	A	99	0	0	4	0
All	All	1242	0	1161	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ARG:HH11	1:A:96:VAL:HG23	1.09	1.15

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:95:ARG:NH1	1:A:96:VAL:HG23	1.78	0.98
1:A:129:LYS:HZ1	1:A:132:ARG:HH12	1.09	0.97
1:A:129:LYS:NZ	1:A:132:ARG:HH12	1.61	0.97
1:A:129:LYS:HE2	1:A:132:ARG:HH22	1.31	0.91
1:A:95:ARG:HD3	3:A:261:EPE:O3S	1.86	0.75
1:A:165:ASP:HB2	4:A:373:HOH:O	1.92	0.68
1:A:146:ILE:HD12	1:A:146:ILE:H	1.57	0.67
1:A:146:ILE:HG22	1:A:147:PRO:HD2	1.77	0.66
1:A:194:TYR:OH	1:A:199:PHE:HB2	1.96	0.65
1:A:129:LYS:HE2	1:A:132:ARG:NH2	2.11	0.64
1:A:122:ASN:O	1:A:123:GLY:C	2.42	0.62
1:A:165:ASP:O	1:A:169:VAL:HG13	2.01	0.60
1:A:129:LYS:HZ1	1:A:132:ARG:NH1	1.92	0.56
1:A:95:ARG:HD3	1:A:96:VAL:H	1.70	0.56
1:A:146:ILE:CG2	1:A:147:PRO:HD2	2.37	0.55
1:A:81:VAL:HG11	1:A:155:MET:HE2	1.89	0.55
1:A:120:THR:O	1:A:144:THR:HA	2.09	0.53
1:A:181:PRO:HB2	1:A:184:LYS:HG3	1.91	0.53
1:A:180:ILE:O	1:A:181:PRO:C	2.51	0.53
1:A:71:MET:HE2	1:A:180:ILE:HD13	1.92	0.50
1:A:129:LYS:NZ	1:A:132:ARG:NH1	2.44	0.50
1:A:194:TYR:CZ	1:A:199:PHE:HB2	2.47	0.49
1:A:125:OCY:H	1:A:125:OCY:HD3	1.76	0.49
1:A:59:GLN:HB3	4:A:304:HOH:O	2.14	0.47
1:A:95:ARG:HD3	1:A:96:VAL:N	2.29	0.47
1:A:85:SER:O	1:A:86:SER:HB2	2.13	0.47
1:A:146:ILE:HD12	1:A:146:ILE:N	2.28	0.47
1:A:99:VAL:HG23	4:A:328:HOH:O	2.16	0.46
1:A:146:ILE:HD13	1:A:149:ASN:HA	1.96	0.46
1:A:163:LEU:HA	1:A:196:LEU:HD11	1.99	0.45
1:A:129:LYS:HZ3	1:A:132:ARG:HH12	1.56	0.44
1:A:63:THR:HG22	1:A:64:ASP:N	2.34	0.43
1:A:135:LEU:HD13	1:A:142:HIS:HB2	1.99	0.43
1:A:125:OCY:HD3	1:A:125:OCY:N	2.35	0.42
1:A:194:TYR:CE1	1:A:199:PHE:CD1	3.08	0.42
1:A:165:ASP:CB	4:A:373:HOH:O	2.61	0.42
1:A:81:VAL:HG22	1:A:88:ILE:HG12	2.01	0.42
1:A:72:ALA:HB3	1:A:179:ARG:HD2	2.01	0.41
1:A:129:LYS:HA	1:A:129:LYS:HD3	1.77	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	143/152 (94%)	134 (94%)	8 (6%)	1 (1%)	18	19

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	123	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/121 (95%)	107 (93%)	8 (7%)	14	16

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	59	GLN
1	A	90	VAL
1	A	96	VAL
1	A	143	THR
1	A	146	ILE
1	A	158	ARG
1	A	169	VAL
1	A	199	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	153	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	OCY	A	125	1	7,8,9	0.48	0	4,8,10	1.91	1 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	OCY	A	125	1	-	1/5/7/9	-

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	125	OCY	CB-SG-CD	3.03	111.26	102.26

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	125	OCY	SG-CD-CE-OZ

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	125	OCY	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 2 ligands modelled in this entry, 1 is monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EPE	A	261	-	15,15,15	1.59	1 (6%)	19,20,20	1.95	6 (31%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EPE	A	261	-	-	2/9/19/19	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	261	EPE	O3S-S	5.46	1.67	1.47

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	261	EPE	O2S-S-C10	3.64	112.23	106.73
3	A	261	EPE	C3-C2-N1	3.63	117.98	110.65
3	A	261	EPE	O3S-S-O2S	-3.12	103.60	111.40
3	A	261	EPE	C6-N1-C2	2.53	114.28	108.84
3	A	261	EPE	O1S-S-C10	2.41	110.38	106.73
3	A	261	EPE	O3S-S-O1S	-2.02	106.35	111.40

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	261	EPE	N4-C7-C8-O8
3	A	261	EPE	C10-C9-N1-C6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	261	EPE	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	145/152 (95%)	-0.63	7 (4%) 35 32	3, 13, 59, 114	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	149	ASN	4.9
1	A	150	SER	4.0
1	A	199	PHE	3.3
1	A	148	GLY	2.6
1	A	145	GLY	2.6
1	A	146	ILE	2.5
1	A	147	PRO	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	OCY	A	125	9/10	0.95	0.06	15,19,37,40	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CA	A	280	1/1	0.86	0.08	41,41,41,41	0
3	EPE	A	261	15/15	0.98	0.04	14,19,22,23	0

6.5 Other polymers [i](#)

There are no such residues in this entry.