



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:57 PM UTC

PDB ID : 3VSC / pdb_00003vsc
Title : Crystal Structure of the K127A Mutant of O-Phosphoserine Sulfhydrylase Complexed with External Schiff Base of Pyridoxal 5'-Phosphate with O-Phospho-L-Serine
Authors : Nakamura, T.; Kawai, Y.; Kataoka, M.; Ishikawa, K.
Deposited on : 2012-04-24
Resolution : 2.07 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.07 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3774 (2.08-2.04)
Clashscore	190562	3883 (2.08-2.04)
Ramachandran outliers	187476	3860 (2.08-2.04)
Sidechain outliers	187428	3860 (2.08-2.04)
RSRZ outliers	180081	3775 (2.08-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	% 81% 14% ...
1	B	389	% 79% 16% ..

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 6159 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Protein CysO.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	382	Total	C	N	O	S	0	4	0
			2949	1869	518	553	9			
1	B	383	Total	C	N	O	S	0	5	0
			2961	1875	521	556	9			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	127	ALA	LYS	engineered mutation	UNP Q9YBL2
B	127	ALA	LYS	engineered mutation	UNP Q9YBL2

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (CCD ID: PLP) (formula: $C_8H_{10}NO_6P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	B	1	Total	C	N	O	P	0	0
			15	8	1	5	1		

- Molecule 3 is PHOSPHOSERINE (CCD ID: SEP) (formula: $C_3H_8NO_6P$).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			11	3	1	6	1		
3	B	1	Total	C	N	O	P	0	0
			11	3	1	6	1		

- Molecule 4 is (4S)-2-METHYL-2,4-PENTANEDIOL (CCD ID: MPD) (formula: $C_6H_{14}O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 8 6 2	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	102	Total O 102 102	0	0
5	B	87	Total O 87 87	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Protein CysO



• Molecule 1: Protein CysO



4 Data and refinement statistics

Property	Value	Source
Space group	P 43 21 2	Depositor
Cell constants a, b, c, α , β , γ	74.68Å 74.68Å 276.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.00 – 2.07 50.00 – 2.07	Depositor EDS
% Data completeness (in resolution range)	99.3 (50.00-2.07) 99.3 (50.00-2.07)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.22 (at 2.08Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.169 , 0.227 0.169 , 0.225	Depositor DCC
R_{free} test set	2439 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	32.3	Xtriage
Anisotropy	0.009	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 29.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6159	wwPDB-VP
Average B, all atoms (Å ²)	35.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.89% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PLP, MPD, SEP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.46	7/3009 (0.2%)	1.33	20/4085 (0.5%)
1	B	1.45	7/3021 (0.2%)	1.33	17/4101 (0.4%)
All	All	1.45	14/6030 (0.2%)	1.33	37/8186 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	342	GLY	N-CA	7.57	1.55	1.45
1	B	158	VAL	CA-CB	6.49	1.62	1.54
1	B	47	VAL	CA-C	6.00	1.60	1.52
1	A	97	THR	CA-C	5.93	1.60	1.53
1	A	244	VAL	CA-CB	5.79	1.61	1.54
1	B	252	ALA	CA-CB	5.77	1.60	1.52
1	B	366	VAL	CA-CB	5.69	1.60	1.53
1	B	344	ALA	CA-CB	-5.62	1.44	1.53
1	A	328	VAL	CA-CB	5.57	1.60	1.54
1	A	160	LEU	C-O	5.32	1.30	1.24
1	A	129	ARG	CA-C	5.22	1.58	1.52
1	B	132	VAL	C-O	-5.19	1.18	1.24
1	A	143	GLU	CA-C	5.13	1.59	1.52
1	A	338	ILE	N-CA	5.08	1.54	1.45

All (37) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	GLU	CA-C-N	-7.87	111.61	119.56
1	A	43	GLU	C-N-CA	-7.87	111.61	119.56
1	B	39	ASN	CA-C-N	7.68	127.57	119.05
1	B	39	ASN	C-N-CA	7.68	127.57	119.05
1	A	15	SER	N-CA-C	6.71	120.39	112.61
1	B	285	VAL	CB-CA-C	-6.67	102.64	110.91
1	A	299	VAL	N-CA-C	6.64	117.14	110.23
1	B	182	PHE	CA-C-N	6.60	127.31	119.98
1	B	182	PHE	C-N-CA	6.60	127.31	119.98
1	B	76	ILE	CB-CA-C	5.92	117.78	110.73
1	A	197	ASP	CA-C-N	-5.81	113.64	119.56
1	A	197	ASP	C-N-CA	-5.81	113.64	119.56
1	A	155	ASN	N-CA-C	5.70	117.17	111.07
1	B	138	LEU	N-CA-CB	-5.58	102.24	110.49
1	B	210	VAL	N-CA-C	-5.57	105.09	110.72
1	B	93	ARG	NE-CZ-NH2	5.50	124.15	119.20
1	B	339	GLY	CA-C-N	5.47	124.93	119.24
1	B	339	GLY	C-N-CA	5.47	124.93	119.24
1	A	45	GLU	CB-CA-C	-5.46	101.57	110.85
1	B	286	GLN	CB-CA-C	-5.45	104.12	110.17
1	A	359	GLU	CA-C-N	-5.38	114.41	119.85
1	A	359	GLU	C-N-CA	-5.38	114.41	119.85
1	B	82	MET	CB-CA-C	-5.36	104.68	111.86
1	A	212	LYS	CD-CE-NZ	5.35	129.03	111.90
1	B	300	GLU	N-CA-C	5.29	118.53	111.75
1	A	42	SER	CA-C-N	-5.28	116.53	123.34
1	A	42	SER	C-N-CA	-5.28	116.53	123.34
1	A	286	GLN	CA-C-N	-5.27	114.53	119.90
1	A	286	GLN	C-N-CA	-5.27	114.53	119.90
1	B	145	GLY	N-CA-C	-5.20	108.89	115.08
1	B	76	ILE	N-CA-C	-5.10	98.68	107.24
1	A	97	THR	CA-C-N	5.03	124.94	119.76
1	A	97	THR	C-N-CA	5.03	124.94	119.76
1	A	39	ASN	CA-C-N	5.02	124.62	119.05
1	A	39	ASN	C-N-CA	5.02	124.62	119.05
1	B	369	ASP	N-CA-C	5.02	115.16	108.38
1	A	138	LEU	N-CA-C	-5.01	107.01	113.23

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	382	LEU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2949	0	2948	34	0
1	B	2961	0	2956	32	0
2	A	15	0	7	0	0
2	B	15	0	7	0	0
3	A	11	0	3	0	0
3	B	11	0	3	2	0
4	A	8	0	14	1	0
5	A	102	0	0	2	0
5	B	87	0	0	3	0
All	All	6159	0	5938	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:212:LYS:HE3	1:A:216:ASN:ND2	1.76	1.01
1:A:254:ARG:HD2	5:A:572:HOH:O	1.69	0.92
1:A:138:LEU:C	1:A:138:LEU:HD23	2.07	0.79
1:B:286:GLN:HG2	1:B:315:LEU:HD22	1.68	0.75
1:B:35[B]:ARG:O	1:B:35[B]:ARG:HD3	1.87	0.74
1:A:212:LYS:HE3	1:A:216:ASN:HD22	1.50	0.74
1:A:139:SER:HA	1:A:169:TYR:OH	1.88	0.73
1:A:212:LYS:CE	1:A:216:ASN:HD22	2.02	0.72
1:A:212:LYS:CE	1:A:216:ASN:ND2	2.51	0.71
1:A:45:GLU:HG2	5:A:504:HOH:O	1.92	0.69
1:B:21:TYR:CB	1:B:45:GLU:HG3	2.29	0.63
1:B:21:TYR:HB3	1:B:45:GLU:HG3	1.83	0.61
1:B:261:GLY:CA	3:B:402:SEP:O3P	2.49	0.61
1:B:284:LEU:HD22	1:B:313:TYR:HB2	1.86	0.58
1:A:102:SER:OG	1:A:104:LEU:HB2	2.05	0.57
1:B:35[B]:ARG:HD2	1:B:46:TYR:CZ	2.40	0.57
1:A:212:LYS:NZ	1:A:216:ASN:HD22	2.03	0.56
1:A:27[A]:GLU:OE1	1:A:30:ARG:NH2	2.37	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:403:MPD:HM1	1:B:82:MET:HE3	1.89	0.55
1:B:29:LEU:HD13	1:B:53:ILE:HD12	1.90	0.54
1:A:305:TRP:HB3	1:A:309:LEU:HD22	1.91	0.53
1:A:329:GLU:O	1:A:333:SER:HB3	2.09	0.53
1:B:138:LEU:C	1:B:138:LEU:HD23	2.34	0.53
1:B:197:ASP:OD1	1:B:199:GLU:HB2	2.10	0.52
1:A:201:PRO:HD2	1:A:205[B]:HIS:CG	2.45	0.51
1:A:212:LYS:HE3	1:A:216:ASN:HD21	1.70	0.51
1:B:35[B]:ARG:HD3	1:B:35[B]:ARG:C	2.34	0.51
1:B:261:GLY:HA2	3:B:402:SEP:O3P	2.11	0.49
1:A:304:LEU:HD12	1:A:308:MET:HE3	1.94	0.49
1:B:258:GLY:HA3	1:B:266:MET:SD	2.52	0.49
1:B:201:PRO:HD2	1:B:205[B]:HIS:CG	2.47	0.48
1:B:78:VAL:CG1	1:B:82:MET:HE1	2.44	0.47
1:A:254:ARG:HA	1:A:254:ARG:NE	2.29	0.47
1:A:101:ARG:HG2	1:A:113:TRP:CE2	2.49	0.47
1:A:10:LEU:O	1:A:13:LEU:HB2	2.15	0.46
1:A:212:LYS:HE2	1:A:212:LYS:C	2.41	0.46
1:A:138:LEU:C	1:A:138:LEU:CD2	2.85	0.46
1:B:154:SER:OG	1:B:183:GLY:HA3	2.16	0.46
1:A:54:GLY:HA3	1:A:240:ARG:HD3	1.97	0.45
1:B:139:SER:HA	1:B:169:TYR:OH	2.15	0.45
1:A:104:LEU:HD22	1:A:104:LEU:HA	1.78	0.45
1:A:91:PHE:CD2	1:A:91:PHE:C	2.94	0.45
1:A:101:ARG:HG2	1:A:113:TRP:CD2	2.52	0.45
1:A:322:GLU:HB3	1:A:347:LYS:HG2	1.99	0.44
1:A:324:MET:HB3	1:A:381:ALA:HB2	1.99	0.44
1:B:143:GLU:HG2	5:B:520:HOH:O	2.17	0.44
1:A:81:GLU:O	1:B:245:GLN:HG2	2.17	0.44
1:B:95:LYS:HA	1:B:96:PRO:C	2.43	0.43
1:B:305:TRP:HB3	1:B:309:LEU:HD22	2.00	0.43
1:B:36:CYS:HA	1:B:59:PRO:HB2	2.01	0.43
1:B:69:GLU:HB2	5:B:509:HOH:O	2.19	0.42
1:A:158:VAL:HG22	1:A:187:PRO:HD3	2.02	0.42
1:A:96:PRO:O	1:B:82:MET:HG3	2.20	0.42
1:B:235:MET:HE3	1:B:235:MET:HB3	1.95	0.41
1:B:45:GLU:HB2	5:B:504:HOH:O	2.20	0.41
1:A:9:TYR:CD1	1:A:9:TYR:N	2.89	0.41
1:A:254:ARG:HD3	1:A:359:GLU:OE1	2.19	0.41
1:A:287:PRO:O	1:A:298:ARG:NH1	2.52	0.41
1:A:258:GLY:HA3	1:A:266:MET:SD	2.62	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:304:LEU:HD23	1:B:305:TRP:CE2	2.56	0.40
1:B:39:ASN:HA	1:B:40:PRO:HD3	1.55	0.40
1:B:90:PHE:CE2	1:B:121:PRO:HB3	2.56	0.40
1:B:133:GLU:HB2	1:B:233:ALA:HB2	2.03	0.40
1:B:259:SER:HB2	1:B:296:ILE:HG21	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	384/389 (99%)	369 (96%)	12 (3%)	3 (1%)	16	8
1	B	386/389 (99%)	376 (97%)	8 (2%)	2 (0%)	24	17
All	All	770/778 (99%)	745 (97%)	20 (3%)	5 (1%)	24	13

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	154	SER
1	B	154	SER
1	B	223	ASN
1	A	14[A]	ASP
1	A	14[B]	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	312/312 (100%)	297 (95%)	15 (5%)	23	16
1	B	313/312 (100%)	290 (93%)	23 (7%)	13	6
All	All	625/624 (100%)	587 (94%)	38 (6%)	17	9

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	13	LEU
1	A	20	SER
1	A	42	SER
1	A	62	ASP
1	A	88	LEU
1	A	101	ARG
1	A	104	LEU
1	A	138	LEU
1	A	154	SER
1	A	203	THR
1	A	209	ARG
1	A	212	LYS
1	A	284	LEU
1	A	309	LEU
1	A	310	ASP
1	B	29	LEU
1	B	43	GLU
1	B	45	GLU
1	B	69	GLU
1	B	71	LEU
1	B	75	ASP
1	B	78	VAL
1	B	88	LEU
1	B	138	LEU
1	B	140	ARG
1	B	141	ARG
1	B	143	GLU
1	B	209	ARG
1	B	212	LYS
1	B	236	ARG
1	B	284	LEU
1	B	289[A]	GLN
1	B	289[B]	GLN
1	B	297	ARG

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	309	LEU
1	B	312	SER
1	B	314	THR
1	B	370	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	39	ASN
1	A	216	ASN
1	B	39	ASN
1	B	286	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

5 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PLP	A	401	3	15,15,16	1.42	3 (20%)	21,22,23	3.21	9 (42%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	SEP	B	402	2	9,10,10	2.01	3 (33%)	9,14,14	2.55	4 (44%)
4	MPD	A	403	-	7,7,7	0.73	0	9,10,10	0.77	0
3	SEP	A	402	2	9,10,10	1.96	3 (33%)	9,14,14	3.91	4 (44%)
2	PLP	B	401	3	15,15,16	2.22	5 (33%)	21,22,23	3.09	11 (52%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PLP	A	401	3	-	1/6/6/8	0/1/1/1
3	SEP	B	402	2	-	2/10/10/10	-
4	MPD	A	403	-	-	1/5/5/5	-
3	SEP	A	402	2	-	5/10/10/10	-
2	PLP	B	401	3	-	1/6/6/8	0/1/1/1

All (14) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	401	PLP	P-O3P	-5.17	1.35	1.54
3	B	402	SEP	P-O1P	4.35	1.64	1.50
2	B	401	PLP	C2A-C2	4.08	1.56	1.50
2	B	401	PLP	O4P-C5A	3.25	1.56	1.44
3	A	402	SEP	P-O1P	3.12	1.60	1.50
3	B	402	SEP	P-OG	2.89	1.69	1.60
3	A	402	SEP	P-OG	2.87	1.69	1.60
2	A	401	PLP	O4P-C5A	2.72	1.54	1.44
3	A	402	SEP	OXT-C	-2.47	1.22	1.30
2	B	401	PLP	O3-C3	2.44	1.42	1.36
2	A	401	PLP	P-O2P	2.31	1.63	1.54
2	B	401	PLP	C3-C2	-2.24	1.38	1.41
3	B	402	SEP	OXT-C	-2.12	1.23	1.30
2	A	401	PLP	C5-C4	-2.04	1.38	1.40

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	402	SEP	OG-CB-CA	9.97	116.75	108.06
2	B	401	PLP	O3P-P-O4P	7.64	126.60	106.67
2	A	401	PLP	O4P-P-O1P	-6.32	89.35	106.44

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	401	PLP	O3P-P-O4P	5.99	122.28	106.67
3	B	402	SEP	OG-CB-CA	5.74	113.07	108.06
2	B	401	PLP	O2P-P-O1P	-5.66	88.79	110.83
2	A	401	PLP	O2P-P-O1P	-5.21	90.54	110.83
2	A	401	PLP	O3P-P-O2P	5.15	127.11	107.80
2	A	401	PLP	O3P-P-O1P	-5.04	91.22	110.83
2	B	401	PLP	O4P-P-O1P	-4.54	94.17	106.44
2	A	401	PLP	C4A-C4-C5	-4.32	116.48	120.94
2	B	401	PLP	C4A-C4-C5	-4.21	116.60	120.94
2	A	401	PLP	O4P-C5A-C5	3.80	116.48	109.36
2	B	401	PLP	C3-C4-C5	3.77	123.12	118.59
3	A	402	SEP	O3P-P-OG	3.77	116.50	106.67
3	B	402	SEP	O2P-P-OG	3.33	115.35	106.67
2	B	401	PLP	O4P-C5A-C5	3.32	115.57	109.36
3	A	402	SEP	OG-P-O1P	-2.90	98.59	106.44
3	A	402	SEP	O2P-P-OG	2.89	114.20	106.67
2	B	401	PLP	C5-C6-N1	-2.85	119.19	123.83
2	B	401	PLP	O3P-P-O2P	2.78	118.23	107.80
2	B	401	PLP	C6-N1-C2	2.56	123.85	119.20
2	A	401	PLP	C4-C3-C2	-2.48	116.18	119.89
2	B	401	PLP	O3P-P-O1P	-2.41	101.43	110.83
2	A	401	PLP	O3-C3-C4	2.41	124.40	118.10
3	B	402	SEP	OXT-C-O	-2.41	118.61	124.08
2	B	401	PLP	O2P-P-O4P	2.35	112.78	106.67
3	B	402	SEP	O3P-P-OG	2.11	112.18	106.67

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	401	PLP	C5A-O4P-P-O3P
2	B	401	PLP	C5A-O4P-P-O3P
3	A	402	SEP	O-C-CA-CB
3	A	402	SEP	CB-OG-P-O1P
3	A	402	SEP	CB-OG-P-O2P
3	A	402	SEP	CB-OG-P-O3P
3	B	402	SEP	O-C-CA-CB
4	A	403	MPD	O2-C2-C3-C4
3	A	402	SEP	OXT-C-CA-N
3	B	402	SEP	OXT-C-CA-N

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	402	SEP	2	0
4	A	403	MPD	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	382/389 (98%)	-0.37	2 (0%) 87 89	16, 32, 51, 64	4 (1%)
1	B	383/389 (98%)	-0.36	2 (0%) 87 89	20, 32, 50, 64	5 (1%)
All	All	765/778 (98%)	-0.37	4 (0%) 87 89	16, 32, 51, 64	9 (1%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	302	GLY	2.9
1	B	289[A]	GLN	2.5
1	A	14[A]	ASP	2.2
1	A	64	CYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	SEP	B	402	11/11	0.94	0.09	30,32,55,56	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	MPD	A	403	8/8	0.94	0.08	32,38,40,42	0
3	SEP	A	402	11/11	0.95	0.09	32,35,54,57	0
2	PLP	A	401	15/16	0.96	0.06	23,29,32,33	0
2	PLP	B	401	15/16	0.98	0.05	22,25,29,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.