



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 07:09 PM UTC

PDB ID : 3VTY / pdb_00003vty
Title : Crystal structure of MamA
Authors : Zeytuni, N.; Baran, D.; Davidov, G.; Zarivach, R.
Deposited on : 2012-06-08
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	184	0.2% 85% 9% 5%
1	B	184	0.2% 86% 8% 5%
1	C	184	3% 82% 12% . .
1	D	184	2% 84% 7% . 7%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 6229 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MamA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	175	Total	C	N	O	S	0	4	0
			1404	907	225	268	4			
1	B	175	Total	C	N	O	S	0	3	0
			1399	900	224	271	4			
1	C	178	Total	C	N	O	S	0	7	0
			1447	937	233	273	4			
1	D	171	Total	C	N	O	S	0	3	0
			1369	883	220	262	4			

- Molecule 2 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cl	0	0
			1	1		
2	B	2	Total	Cl	0	0
			2	2		
2	C	1	Total	Cl	0	0
			1	1		
2	D	2	Total	Cl	0	0
			2	2		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	136	Total	O	0	0
			136	136		
3	B	112	Total	O	0	0
			112	112		
3	C	177	Total	O	0	0
			177	177		
3	D	179	Total	O	0	0
			179	179		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: MamA

Chain A: 



- Molecule 1: MamA

Chain B: 



- Molecule 1: MamA

Chain C: 



- Molecule 1: MamA

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.70Å 101.31Å 139.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	81.99 – 2.00 81.99 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.1 (81.99-2.00) 99.1 (81.99-2.00)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.37 (at 2.01Å)	Xtriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.163 , 0.219 0.168 , 0.223	Depositor DCC
R_{free} test set	2542 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	18.3	Xtriage
Anisotropy	0.115	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 51.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6229	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 26.25 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.7257e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	1.33	5/1438 (0.3%)	1.09	3/1929 (0.2%)
1	B	1.27	1/1430 (0.1%)	1.04	1/1920 (0.1%)
1	C	1.32	4/1490 (0.3%)	1.07	0/1997
1	D	1.36	3/1400 (0.2%)	1.12	0/1878
All	All	1.32	13/5758 (0.2%)	1.08	4/7724 (0.1%)

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	167	GLU	CB-CG	-6.64	1.32	1.52
1	C	193	LYS	CG-CD	-5.84	1.34	1.52
1	A	146	ALA	CA-CB	5.64	1.62	1.53
1	B	59	ASP	CA-C	5.57	1.60	1.52
1	C	146	ALA	CA-CB	5.55	1.61	1.53
1	D	139	ILE	CB-CG2	5.44	1.70	1.52
1	C	55	ARG	C-O	-5.35	1.18	1.24
1	A	182	ALA	CA-CB	5.34	1.61	1.53
1	D	130	ALA	CA-CB	5.22	1.61	1.53
1	D	143	THR	CA-CB	5.20	1.62	1.53
1	C	153	GLY	N-CA	-5.19	1.39	1.45
1	A	75	PRO	C-O	-5.07	1.17	1.24
1	A	181	ARG	C-O	-5.03	1.17	1.24

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	167	GLU	CB-CA-C	-5.66	102.00	110.88
1	A	74	ASP	CA-C-N	5.23	124.89	119.56
1	A	74	ASP	C-N-CA	5.23	124.89	119.56
1	B	70	VAL	CB-CA-C	-5.02	105.54	111.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1404	0	1451	13	0
1	B	1399	0	1428	14	0
1	C	1447	0	1514	36	0
1	D	1369	0	1401	19	0
2	A	1	0	0	0	0
2	B	2	0	0	0	0
2	C	1	0	0	0	0
2	D	2	0	0	0	0
3	A	136	0	0	0	0
3	B	112	0	0	2	0
3	C	177	0	0	5	0
3	D	179	0	0	3	0
All	All	6229	0	5794	68	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:78:VAL:CG2	1:C:143[A]:THR:HG21	1.80	1.12
1:C:196[A]:ARG:HH11	1:C:196[A]:ARG:CG	1.71	1.04
1:C:196[A]:ARG:HG3	1:C:196[A]:ARG:NH1	1.60	1.01
1:B:78:VAL:HG21	1:C:143[A]:THR:HG21	1.42	1.00
1:C:48:MET:HE1	1:C:83[B]:LYS:HD2	1.51	0.91
1:C:196[A]:ARG:HH11	1:C:196[A]:ARG:HG3	0.78	0.89
1:D:46:ILE:O	1:D:46:ILE:HG22	1.74	0.87
1:B:78:VAL:HG23	1:C:143[A]:THR:HG21	1.55	0.86
1:A:152:LEU:HD23	1:D:46:ILE:HD11	1.66	0.77
1:C:48:MET:HE1	1:C:83[B]:LYS:CD	2.17	0.75
1:C:173[A]:ILE:HD12	1:C:183:TYR:CZ	2.22	0.74

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:42:GLU:N	1:C:42:GLU:OE1	2.24	0.71
1:B:136[B]:GLN:HE21	1:B:136[B]:GLN:HA	1.56	0.70
1:C:198:GLU:OE1	1:C:201[A]:LYS:NZ	2.22	0.70
1:C:173[A]:ILE:HD12	1:C:183:TYR:CE2	2.28	0.68
1:C:48:MET:CE	1:C:83[B]:LYS:CE	2.73	0.67
1:B:44:THR:O	1:B:48:MET:HG3	1.97	0.65
1:D:46:ILE:O	1:D:46:ILE:CG2	2.43	0.65
1:A:152:LEU:HD23	1:D:46:ILE:CD1	2.27	0.63
1:D:94:PRO:HG3	1:D:124:MET:HE2	1.80	0.63
1:C:48:MET:HE1	1:C:83[B]:LYS:CE	2.29	0.63
1:B:193:LYS:HB3	1:B:195:LEU:HD13	1.81	0.62
1:A:167:GLU:HG2	1:D:46:ILE:HG23	1.81	0.62
1:C:49:ASP:HA	3:C:457:HOH:O	2.00	0.61
1:A:132[A]:ILE:HD11	1:A:155:VAL:CG1	2.30	0.60
1:A:210:GLU:OE1	1:B:180:ILE:HG22	2.00	0.60
1:D:57:LYS:NZ	3:D:486:HOH:O	2.34	0.60
1:D:50:ILE:HD13	3:D:546:HOH:O	2.02	0.58
1:B:193:LYS:CB	1:B:195:LEU:HD13	2.33	0.58
1:B:48:MET:HE1	1:B:79:GLU:CD	2.28	0.58
1:C:180:ILE:HG22	1:D:210[A]:GLU:OE2	2.02	0.58
1:C:196[A]:ARG:CG	1:C:196[A]:ARG:NH1	2.42	0.57
1:C:128[A]:LYS:HE3	1:C:158:SER:OG	2.05	0.57
1:C:48:MET:CE	1:C:83[B]:LYS:HE3	2.35	0.57
1:C:173[A]:ILE:CD1	1:C:183:TYR:CE2	2.88	0.56
1:A:42:GLU:HA	1:A:42:GLU:OE1	2.06	0.55
1:C:83[B]:LYS:HE2	3:C:483:HOH:O	2.07	0.55
1:B:72:LYS:NZ	3:B:483:HOH:O	2.38	0.55
1:D:50:ILE:HD12	1:D:50:ILE:N	2.22	0.54
1:B:78:VAL:HG23	1:C:143[A]:THR:CG2	2.31	0.53
1:C:142:ASN:HD21	1:C:144:VAL:HG22	1.74	0.52
1:A:152:LEU:CD2	1:D:46:ILE:CD1	2.88	0.52
1:A:132[A]:ILE:HD11	1:A:155:VAL:HG11	1.91	0.51
1:A:132[A]:ILE:CD1	1:A:155:VAL:HG11	2.41	0.51
1:D:50:ILE:N	1:D:50:ILE:CD1	2.74	0.51
1:C:48:MET:CE	1:C:83[B]:LYS:NZ	2.75	0.50
1:B:78:VAL:CG2	1:C:143[B]:THR:HG21	2.42	0.49
1:C:218:LEU:O	1:C:219:ALA:HB3	2.12	0.49
1:C:197:ASP:O	1:C:201[A]:LYS:HG3	2.12	0.49
1:D:46:ILE:O	1:D:47:TYR:HD2	1.95	0.49
1:C:48:MET:HE2	1:C:83[B]:LYS:HE3	1.96	0.48
1:A:132[A]:ILE:HD12	1:A:152:LEU:HD12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167[B]:GLU:HG3	3:B:474:HOH:O	2.14	0.47
1:C:49:ASP:CA	3:C:457:HOH:O	2.60	0.46
1:D:46:ILE:O	1:D:47:TYR:CD2	2.68	0.46
1:D:196:ARG:NH1	3:D:494:HOH:O	2.48	0.46
1:B:60:PHE:O	1:B:64:ILE:HG13	2.16	0.45
1:A:167:GLU:HB3	1:D:46:ILE:HG23	1.99	0.45
1:C:198:GLU:OE1	1:C:201[A]:LYS:CE	2.67	0.43
1:C:213:LYS:HZ1	1:D:213:LYS:HZ2	1.66	0.43
1:C:196[B]:ARG:NH1	3:C:492:HOH:O	2.51	0.43
1:D:215:LYS:O	1:D:216:TYR:CD2	2.72	0.42
1:C:213:LYS:NZ	1:D:213:LYS:NZ	2.68	0.42
1:C:143[B]:THR:OG1	3:C:572:HOH:O	2.08	0.41
1:A:104:PHE:CD1	1:A:104:PHE:C	2.97	0.41
1:C:217:GLU:O	1:C:218:LEU:HD12	2.20	0.41
1:C:218:LEU:O	1:C:219:ALA:CB	2.69	0.40
1:A:105:VAL:O	1:A:109:THR:HG22	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	177/184 (96%)	175 (99%)	2 (1%)	0	100	100
1	B	176/184 (96%)	172 (98%)	3 (2%)	1 (1%)	21	17
1	C	183/184 (100%)	179 (98%)	4 (2%)	0	100	100
1	D	172/184 (94%)	170 (99%)	2 (1%)	0	100	100
All	All	708/736 (96%)	696 (98%)	11 (2%)	1 (0%)	48	46

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	126	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	146/149 (98%)	138 (94%)	8 (6%)	19	17
1	B	145/149 (97%)	143 (99%)	2 (1%)	59	66
1	C	151/149 (101%)	141 (93%)	10 (7%)	15	12
1	D	141/149 (95%)	135 (96%)	6 (4%)	26	25
All	All	583/596 (98%)	557 (96%)	26 (4%)	28	23

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	GLU
1	A	53	LYS
1	A	86[A]	LYS
1	A	86[B]	LYS
1	A	167	GLU
1	A	209	LYS
1	A	212	LYS
1	A	215	LYS
1	B	64	ILE
1	B	205	LYS
1	C	42	GLU
1	C	45	THR
1	C	83[A]	LYS
1	C	83[B]	LYS
1	C	143[A]	THR
1	C	143[B]	THR
1	C	196[A]	ARG
1	C	196[B]	ARG
1	C	205	LYS
1	C	216	TYR
1	D	46	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	47	TYR
1	D	50	ILE
1	D	196	ARG
1	D	201	LYS
1	D	205	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	122	ASN
1	C	142	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	175/184 (95%)	-0.49	2 (1%) 78 77	7, 18, 33, 55	7 (4%)
1	B	175/184 (95%)	-0.30	2 (1%) 78 77	11, 22, 39, 64	3 (1%)
1	C	178/184 (96%)	-0.57	5 (2%) 55 54	7, 14, 31, 63	10 (5%)
1	D	171/184 (92%)	-0.51	3 (1%) 67 67	6, 16, 37, 57	5 (2%)
All	All	699/736 (94%)	-0.47	12 (1%) 69 68	6, 17, 37, 64	25 (3%)

All (12) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	216	TYR	10.8
1	A	216	TYR	4.5
1	C	219	ALA	4.1
1	D	47	TYR	4.0
1	D	46	ILE	3.7
1	A	53	LYS	3.3
1	C	218	LEU	3.2
1	B	216	TYR	3.1
1	C	49	ASP	2.8
1	D	216	TYR	2.8
1	C	53	LYS	2.2
1	B	214	ALA	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	CL	B	302	1/1	0.98	0.08	29,29,29,29	0
2	CL	B	301	1/1	0.99	0.06	15,15,15,15	0
2	CL	A	301	1/1	0.99	0.05	24,24,24,24	0
2	CL	C	301	1/1	0.99	0.06	11,11,11,11	0
2	CL	D	301	1/1	0.99	0.04	11,11,11,11	0
2	CL	D	302	1/1	0.99	0.05	24,24,24,24	0

6.5 Other polymers [i](#)

There are no such residues in this entry.