



Full wwPDB X-ray Structure Validation Report ⓘ

Jul 6, 2026 – 06:23 PM JST

PDB ID : 9VTL / pdb_00009vtl
Title : Crystal Structure of the SARS-CoV-2 (COVID-19) main protease with inhibitor AD06cn
Authors : Kong, W.P.; Yin, W.; Wong, K.Y.
Deposited on : 2025-07-10
Resolution : 1.97 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.015 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.97 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1506 (1.98-1.98)
Clashscore	190562	1534 (1.98-1.98)
Ramachandran outliers	187476	1518 (1.98-1.98)
Sidechain outliers	187428	1518 (1.98-1.98)
RSRZ outliers	180081	1506 (1.98-1.98)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	306	
2	B	4	

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 5223 atoms, of which 2410 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 3C-like proteinase nsp5.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	300	Total	C	H	N	O	S	77	21	0
			4781	1523	2378	405	451	24			

- Molecule 2 is a protein called inhibitor AD06cn.

Mol	Chain	Residues	Atoms							ZeroOcc	AltConf	Trace
2	B	4	Total	C	Cl	H	N	O	S	1	0	0
			66	22	1	32	6	4	1			


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	375	Total	O	0	0
			375	375		
3	B	1	Total	O	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 3C-like proteinase nsp5



- Molecule 2: inhibitor AD06cn



4 Data and refinement statistics

Property	Value
Space group	I 1 2 1
Cell constants a, b, c, α , β , γ	51.62Å 80.47Å 89.15Å 90.00° 97.17° 90.00°
Resolution (Å)	24.29 – 1.97 24.29 – 1.97
% Data completeness (in resolution range)	98.5 (24.29-1.97) 87.0 (24.29-1.97)
R_{merge}	0.15
R_{sym}	(Not available)
$\langle I/\sigma(I) \rangle$ ¹	1.17 (at 1.96Å)
Refinement program	REFMAC 5.8.0430 (refmacat 0.4.105), REFMAC 5.8.0430 (refmacat 0.4.105)
R, R_{free}	0.216 , 0.265 0.215 , 0.258
R_{free} test set	1243 reflections (4.83%)
Wilson B-factor (Å ²)	17.4
Anisotropy	0.541
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 54.7
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$
Estimated twinning fraction	No twinning to report.
F_o, F_c correlation	0.92
Total number of atoms	5223
Average B, all atoms (Å ²)	27.0

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 79I, A1L8H

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.64	0/2532	1.18	6/3437 (0.2%)
2	B	0.58	0/14	1.70	0/18
All	All	0.64	0/2546	1.19	6/3455 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	CYS	CB-CA-C	-10.79	87.67	109.79
1	A	154	TYR	CB-CA-C	8.30	126.93	110.42
1	A	60	ARG	CB-CA-C	-7.20	97.03	110.36
1	A	103	PHE	CA-CB-CG	6.63	120.43	113.80
1	A	60	ARG	N-CA-CB	5.69	119.23	110.46
1	A	127	GLN	N-CA-CB	-5.57	101.72	110.69

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	60	ARG	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2403	2378	2290	33	0
2	B	34	32	19	7	0
3	A	375	0	0	9	1
3	B	1	0	0	0	0
All	All	2813	2410	2309	34	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:MET:HE2	2:B:3:LEU:HG	1.39	1.03
1:A:191:ALA:HA	2:B:1:A1L8H:CL1	2.03	0.95
1:A:49:MET:CE	2:B:3:LEU:HG	2.17	0.74
1:A:142:ASN:OD1	3:A:401:HOH:O	2.07	0.71
1:A:225:THR:OG1	1:A:269[A]:LYS:NZ	2.29	0.66
1:A:286[B]:LEU:HD12	1:A:286[B]:LEU:C	2.22	0.65
1:A:45:THR:HG21	3:A:698:HOH:O	2.06	0.55
1:A:286[B]:LEU:HD12	1:A:286[B]:LEU:O	2.09	0.53
1:A:132:PRO:HD2	1:A:197:ASP:OD1	2.09	0.52
1:A:63:ASN:OD1	1:A:80:HIS:ND1	2.39	0.52
1:A:279:ARG:HD2	3:A:651:HOH:O	2.10	0.52
1:A:41:HIS:ND1	2:B:3:LEU:HD12	2.26	0.51
1:A:40:ARG:HD3	1:A:85:CYS:HA	1.94	0.49
1:A:154:TYR:CD1	1:A:154:TYR:N	2.77	0.49
1:A:263:ASP:OD2	3:A:402:HOH:O	2.19	0.49
1:A:5:LYS:HE3	1:A:291:PHE:CZ	2.48	0.49
1:A:154:TYR:OH	3:A:403:HOH:O	2.20	0.48
1:A:190:THR:HG23	1:A:191:ALA:O	2.13	0.48
1:A:82[A]:MET:HE3	1:A:82[A]:MET:HB2	1.72	0.45
1:A:72:ASN:O	1:A:73[A]:VAL:C	2.59	0.45
1:A:19:GLN:OE1	1:A:26:THR:HG21	2.17	0.44
1:A:73[B]:VAL:O	1:A:73[B]:VAL:CG2	2.66	0.44
1:A:31:TRP:CD2	1:A:95:ASN:HB2	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:234:ALA:HB3	3:A:428:HOH:O	2.18	0.43
1:A:191:ALA:CA	2:B:1:A1L8H:CL1	2.92	0.42
1:A:10:SER:OG	1:A:14:GLU:OE2	2.30	0.42
1:A:113:SER:O	1:A:149:GLY:HA2	2.19	0.42
1:A:190:THR:O	2:B:1:A1L8H:C22	2.68	0.42
1:A:100:LYS:HB3	3:A:691:HOH:O	2.19	0.42
1:A:31:TRP:CE2	1:A:95:ASN:HB2	2.55	0.41
1:A:190:THR:O	2:B:1:A1L8H:S1	2.78	0.41
1:A:286[B]:LEU:C	1:A:286[B]:LEU:CD1	2.91	0.41
1:A:88:LYS:NZ	3:A:409:HOH:O	2.33	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:667:HOH:O	3:A:712:HOH:O[2_655]	2.14	0.06

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	319/306 (104%)	311 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	275/263 (105%)	265 (96%)	10 (4%)	31	20
2	B	2/2 (100%)	2 (100%)	0	100	100
All	All	277/265 (104%)	267 (96%)	10 (4%)	32	20

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	47	GLU
1	A	55	GLU
1	A	59	ILE
1	A	62	SER
1	A	72	ASN
1	A	74[A]	GLN
1	A	74[B]	GLN
1	A	190	THR
1	A	196	THR
1	A	227	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	41	HIS
1	A	51	ASN
1	A	72	ASN
1	A	119	ASN
1	A	274	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	300/306 (98%)	0.23	12 (4%)	42 52	6, 21, 53, 81	12 (4%)
2	B	2/4 (50%)	2.85	1 (50%)	0 0	47, 47, 47, 55	0
All	All	302/310 (97%)	0.25	13 (4%)	40 49	6, 22, 53, 81	12 (3%)

All (13) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	TYR	5.1
2	B	2	VAL	4.2
1	A	50	LEU	3.7
1	A	191	ALA	3.2
1	A	46	SER	2.9
1	A	189	GLN	2.9
1	A	49	MET	2.7
1	A	190	THR	2.7
1	A	188	ARG	2.5
1	A	73[A]	VAL	2.4
1	A	196	THR	2.4
1	A	72	ASN	2.2
1	A	277	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.