



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 06:08 AM UTC

PDB ID : 2VUU / pdb_00002vuu
Title : Crystal structure of NADP-bound NmrA-AreA zinc finger complex
Authors : Kotaka, M.; Johnson, C.; Lamb, H.K.; Hawkins, A.R.; Ren, J.; Stammers, D.K.
Deposited on : 2008-05-30
Resolution : 2.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3866 (2.80-2.80)
Clashscore	190562	4276 (2.80-2.80)
Ramachandran outliers	187476	4196 (2.80-2.80)
Sidechain outliers	187428	4198 (2.80-2.80)
RSRZ outliers	180081	3869 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	352	52% 34% • 10%
1	B	352	45% 42% • 9%
1	C	352	52% 35% • 10%
1	D	352	50% 37% • • 9%
1	E	352	57% 30% • 10%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	352	 50% 37% • 10%
1	G	352	 55% 33% • 9%
1	H	352	 49% 38% • 9%
2	I	43	 58% 37% • •
2	J	43	 63% 33% • •
2	K	43	 44% 47% 7% •
2	L	43	 30% 53% 12% • •
2	M	43	 70% 26% • •
2	N	43	 58% 40% •
2	O	43	 60% 30% 5% 5%
2	P	43	 53% 42% • •

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 23588 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NITROGEN METABOLITE REPRESSION REGULATOR NMRA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	318	Total	C	N	O	S	0	0	0
			2530	1637	428	457	8			
1	B	319	Total	C	N	O	S	0	0	0
			2541	1643	432	458	8			
1	C	318	Total	C	N	O	S	0	0	0
			2530	1637	428	457	8			
1	D	319	Total	C	N	O	S	0	0	0
			2541	1643	432	458	8			
1	E	318	Total	C	N	O	S	0	0	0
			2530	1637	428	457	8			
1	F	318	Total	C	N	O	S	0	0	0
			2530	1637	428	457	8			
1	G	319	Total	C	N	O	S	0	0	0
			2541	1643	432	458	8			
1	H	319	Total	C	N	O	S	0	0	0
			2539	1642	430	459	8			


There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	238	ARG	LEU	conflict	UNP O59919
B	238	ARG	LEU	conflict	UNP O59919
C	238	ARG	LEU	conflict	UNP O59919
D	238	ARG	LEU	conflict	UNP O59919
E	238	ARG	LEU	conflict	UNP O59919
F	238	ARG	LEU	conflict	UNP O59919
G	238	ARG	LEU	conflict	UNP O59919
H	238	ARG	LEU	conflict	UNP O59919

- Molecule 2 is a protein called NITROGEN REGULATORY PROTEIN AREA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	I	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	J	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	K	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	L	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	M	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	N	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			
2	O	41	Total	C	N	O	S	0	0	0
			318	200	59	55	4			
2	P	42	Total	C	N	O	S	0	0	0
			326	206	60	56	4			

- Molecule 3 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NAP) (formula: C₂₁H₂₈N₇O₁₇P₃).



Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	D	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	E	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	F	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	G	1	Total	C	N	O	P	0	0
			48	21	7	17	3		
3	H	1	Total	C	N	O	P	0	0
			48	21	7	17	3		

- Molecule 4 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	I	1	Total	Zn	0	0
			1	1		
4	J	1	Total	Zn	0	0
			1	1		
4	K	1	Total	Zn	0	0
			1	1		
4	L	1	Total	Zn	0	0
			1	1		
4	M	1	Total	Zn	0	0
			1	1		
4	N	1	Total	Zn	0	0
			1	1		
4	O	1	Total	Zn	0	0
			1	1		
4	P	1	Total	Zn	0	0
			1	1		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	32	Total	O	0	0
			32	32		
5	B	26	Total	O	0	0
			26	26		
5	C	39	Total	O	0	0
			39	39		
5	D	36	Total	O	0	0
			36	36		

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	E	42	Total 42	O 42	0	0
5	F	33	Total 33	O 33	0	0
5	G	41	Total 41	O 41	0	0
5	H	33	Total 33	O 33	0	0
5	I	4	Total 4	O 4	0	0
5	J	3	Total 3	O 3	0	0
5	K	4	Total 4	O 4	0	0
5	L	4	Total 4	O 4	0	0
5	M	5	Total 5	O 5	0	0
5	N	3	Total 3	O 3	0	0
5	O	6	Total 6	O 6	0	0
5	P	3	Total 3	O 3	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA




• Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA




• Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain C: 




• Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain D: 




• Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain E: 





- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain F: 50% 37% 10%




- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA


Chain G:  55% 33% 9%



- Molecule 1: NITROGEN METABOLITE REPRESSION REGULATOR NMRA

Chain H: 49% 38% 9%





• Molecule 2: NITROGEN REGULATORY PROTEIN AREA

Chain I: 58% 37% ..



• Molecule 2: NITROGEN REGULATORY PROTEIN AREA

Chain J: 63% 33% ..



• Molecule 2: NITROGEN REGULATORY PROTEIN AREA

Chain K: 44% 47% 7% .




• Molecule 2: NITROGEN REGULATORY PROTEIN AREA

Chain L: 30% 53% 12% ..



• Molecule 2: NITROGEN REGULATORY PROTEIN AREA

Chain M: 70% 26% ..



• Molecule 2: NITROGEN REGULATORY PROTEIN AREA


Chain N: 58% 40% .



● Molecule 2: NITROGEN REGULATORY PROTEIN AREA



● Molecule 2: NITROGEN REGULATORY PROTEIN AREA



4 Data and refinement statistics

Property	Value	Source
Space group	H 3	Depositor
Cell constants a, b, c, α , β , γ	231.73Å 231.73Å 223.45Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	29.86 – 2.80 29.86 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.9 (29.86-2.80) 100.0 (29.86-2.80)	Depositor EDS
R_{merge}	0.14	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 2.80Å)	Xtriage
Refinement program	CNS 1.1	Depositor
R, R_{free}	0.232 , 0.287 0.232 , 0.287	Depositor DCC
R_{free} test set	5547 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	50.8	Xtriage
Anisotropy	0.101	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 27.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	0.459 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	23588	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.37 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.1182e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.52	0/2606	1.10	18/3557 (0.5%)
1	B	0.51	0/2617	1.10	20/3571 (0.6%)
1	C	0.51	0/2606	1.07	17/3557 (0.5%)
1	D	0.54	0/2617	1.06	19/3571 (0.5%)
1	E	0.52	0/2606	1.09	17/3557 (0.5%)
1	F	0.54	0/2606	1.13	24/3557 (0.7%)
1	G	0.51	0/2617	1.11	17/3571 (0.5%)
1	H	0.51	0/2615	1.10	26/3569 (0.7%)
2	I	0.49	0/334	0.94	1/456 (0.2%)
2	J	0.52	0/334	1.04	2/456 (0.4%)
2	K	0.51	0/334	0.95	0/456
2	L	0.65	1/334 (0.3%)	1.11	3/456 (0.7%)
2	M	0.54	0/334	0.90	0/456
2	N	0.46	0/334	0.94	0/456
2	O	0.50	0/326	0.98	0/445
2	P	0.47	0/334	0.94	0/456
All	All	0.52	1/23554 (0.0%)	1.08	164/32147 (0.5%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	688	PRO	CA-C	5.17	1.59	1.52

The worst 5 of 164 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	F	156	ASN	N-CA-C	-10.25	99.32	112.23
1	H	156	ASN	N-CA-C	-10.25	100.18	112.89
1	A	156	ASN	N-CA-C	-10.02	100.47	112.89
1	G	276	TYR	N-CA-C	-9.92	99.49	111.69
1	E	156	ASN	N-CA-C	-9.91	100.60	112.89

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2530	0	2476	104	0
1	B	2541	0	2489	114	0
1	C	2530	0	2476	113	0
1	D	2541	0	2489	111	0
1	E	2530	0	2476	88	0
1	F	2530	0	2476	106	0
1	G	2541	0	2489	103	0
1	H	2539	0	2484	117	0
2	I	326	0	326	22	0
2	J	326	0	326	12	0
2	K	326	0	327	20	0
2	L	326	0	326	30	0
2	M	326	0	326	21	0
2	N	326	0	326	14	0
2	O	318	0	315	17	0
2	P	326	0	326	21	0
3	A	48	0	25	2	0
3	B	48	0	25	1	0
3	C	48	0	25	1	0
3	D	48	0	25	2	0
3	E	48	0	25	0	0
3	F	48	0	25	1	0
3	G	48	0	25	0	0
3	H	48	0	25	3	0
4	I	1	0	0	0	0
4	J	1	0	0	0	0
4	K	1	0	0	0	0
4	L	1	0	0	0	0
4	M	1	0	0	0	0
4	N	1	0	0	0	0
4	O	1	0	0	0	0
4	P	1	0	0	0	0
5	A	32	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	26	0	0	0	0
5	C	39	0	0	2	0
5	D	36	0	0	3	0
5	E	42	0	0	4	0
5	F	33	0	0	1	0
5	G	41	0	0	4	0
5	H	33	0	0	4	0
5	I	4	0	0	1	0
5	J	3	0	0	0	0
5	K	4	0	0	2	0
5	L	4	0	0	0	0
5	M	5	0	0	0	0
5	N	3	0	0	0	0
5	O	6	0	0	0	0
5	P	3	0	0	0	0
All	All	23588	0	22653	986	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 986 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:M:686:ASN:HB2	2:M:690:GLN:HB2	1.20	1.11
1:H:167:MET:HE1	1:H:240:VAL:HG11	1.28	1.09
1:D:239:ARG:HH11	1:D:239:ARG:HB3	1.20	1.06
1:F:82:THR:HG22	3:F:1353:NAP:H51N	1.36	1.03
1:C:82:THR:HG22	3:C:1353:NAP:H51N	1.40	1.02

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	314/352 (89%)	288 (92%)	23 (7%)	3 (1%)	12	38
1	B	315/352 (90%)	296 (94%)	18 (6%)	1 (0%)	36	66
1	C	314/352 (89%)	286 (91%)	23 (7%)	5 (2%)	7	27
1	D	315/352 (90%)	286 (91%)	27 (9%)	2 (1%)	21	51
1	E	314/352 (89%)	288 (92%)	25 (8%)	1 (0%)	36	66
1	F	314/352 (89%)	286 (91%)	27 (9%)	1 (0%)	36	66
1	G	315/352 (90%)	284 (90%)	29 (9%)	2 (1%)	21	51
1	H	315/352 (90%)	288 (91%)	23 (7%)	4 (1%)	9	31
2	I	40/43 (93%)	36 (90%)	3 (8%)	1 (2%)	4	16
2	J	40/43 (93%)	35 (88%)	5 (12%)	0	100	100
2	K	40/43 (93%)	34 (85%)	3 (8%)	3 (8%)	1	2
2	L	40/43 (93%)	34 (85%)	4 (10%)	2 (5%)	1	5
2	M	40/43 (93%)	37 (92%)	3 (8%)	0	100	100
2	N	40/43 (93%)	33 (82%)	6 (15%)	1 (2%)	4	16
2	O	39/43 (91%)	33 (85%)	5 (13%)	1 (3%)	4	15
2	P	40/43 (93%)	37 (92%)	3 (8%)	0	100	100
All	All	2835/3160 (90%)	2581 (91%)	227 (8%)	27 (1%)	12	38

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	350	TRP
2	I	711	SER
2	L	689	GLU
1	A	87	ASP
1	A	256	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	266/284 (94%)	254 (96%)	12 (4%)	24	58
1	B	267/284 (94%)	256 (96%)	11 (4%)	27	62
1	C	266/284 (94%)	258 (97%)	8 (3%)	36	72
1	D	267/284 (94%)	258 (97%)	9 (3%)	32	68
1	E	266/284 (94%)	259 (97%)	7 (3%)	40	75
1	F	266/284 (94%)	256 (96%)	10 (4%)	29	64
1	G	267/284 (94%)	260 (97%)	7 (3%)	40	75
1	H	267/284 (94%)	264 (99%)	3 (1%)	65	87
2	I	38/39 (97%)	36 (95%)	2 (5%)	20	52
2	J	38/39 (97%)	37 (97%)	1 (3%)	40	75
2	K	38/39 (97%)	36 (95%)	2 (5%)	20	52
2	L	38/39 (97%)	35 (92%)	3 (8%)	11	35
2	M	38/39 (97%)	36 (95%)	2 (5%)	20	52
2	N	38/39 (97%)	37 (97%)	1 (3%)	40	75
2	O	37/39 (95%)	35 (95%)	2 (5%)	20	51
2	P	38/39 (97%)	37 (97%)	1 (3%)	40	75
All	All	2435/2584 (94%)	2354 (97%)	81 (3%)	33	69

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	51	PRO
2	K	694	CYS
1	G	117	SER
1	H	209	GLN
2	M	690	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 103 such sidechains are listed below:

Mol	Chain	Res	Type
1	G	57	GLN
1	H	62	ASN
2	O	675	ASN
1	G	84	GLN
1	G	214	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 16 ligands modelled in this entry, 8 are monoatomic - leaving 8 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAP	E	1353	-	50,52,52	2.25	12 (24%)	71,80,80	1.32	7 (9%)
3	NAP	F	1353	-	50,52,52	2.27	11 (22%)	71,80,80	1.32	7 (9%)
3	NAP	G	1353	-	50,52,52	2.33	14 (28%)	71,80,80	1.29	7 (9%)
3	NAP	D	1353	-	50,52,52	2.28	11 (22%)	71,80,80	1.32	7 (9%)
3	NAP	A	1353	-	50,52,52	2.29	10 (20%)	71,80,80	1.29	7 (9%)
3	NAP	B	1353	-	50,52,52	2.31	9 (18%)	71,80,80	1.28	7 (9%)
3	NAP	H	1353	-	50,52,52	2.35	10 (20%)	71,80,80	1.27	8 (11%)
3	NAP	C	1353	-	50,52,52	2.25	11 (22%)	71,80,80	1.31	7 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAP	E	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	F	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	G	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	D	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	A	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	B	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	H	1353	-	-	5/35/67/67	0/5/5/5
3	NAP	C	1353	-	-	5/35/67/67	0/5/5/5

The worst 5 of 88 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	1353	NAP	C2N-N1N	11.43	1.47	1.35
3	G	1353	NAP	C2N-N1N	11.30	1.47	1.35
3	F	1353	NAP	C2N-N1N	11.25	1.47	1.35
3	D	1353	NAP	C2N-N1N	11.22	1.47	1.35
3	H	1353	NAP	C2N-N1N	11.05	1.47	1.35

The worst 5 of 57 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	G	1353	NAP	O4B-C4B-C5B	5.69	127.55	109.33
3	H	1353	NAP	O4B-C4B-C5B	5.58	127.20	109.33
3	F	1353	NAP	O4B-C4B-C5B	5.55	127.11	109.33
3	B	1353	NAP	O4B-C4B-C5B	5.53	127.05	109.33
3	A	1353	NAP	O4B-C4B-C5B	5.43	126.73	109.33

There are no chirality outliers.

5 of 40 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1353	NAP	O4D-C1D-N1N-C2N
3	B	1353	NAP	O4D-C1D-N1N-C2N
3	C	1353	NAP	O4D-C1D-N1N-C2N
3	D	1353	NAP	O4D-C1D-N1N-C2N
3	E	1353	NAP	O4D-C1D-N1N-C2N


There are no ring outliers.


6 monomers are involved in 10 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	F	1353	NAP	1	0
3	D	1353	NAP	2	0
3	A	1353	NAP	2	0
3	B	1353	NAP	1	0
3	H	1353	NAP	3	0
3	C	1353	NAP	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.




















5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	318/352 (90%)	-1.73	0 100 100	22, 47, 74, 91	0
1	B	319/352 (90%)	-1.68	0 100 100	21, 49, 78, 99	0
1	C	318/352 (90%)	-1.74	0 100 100	23, 45, 73, 90	0
1	D	319/352 (90%)	-1.74	0 100 100	20, 46, 74, 91	0
1	E	318/352 (90%)	-1.76	0 100 100	18, 43, 71, 83	0
1	F	318/352 (90%)	-1.75	0 100 100	23, 43, 69, 81	0
1	G	319/352 (90%)	-1.72	0 100 100	25, 44, 71, 99	0
1	H	319/352 (90%)	-1.76	0 100 100	21, 43, 73, 85	0
2	I	42/43 (97%)	-1.48	0 100 100	30, 60, 121, 126	0
2	J	42/43 (97%)	-1.43	0 100 100	35, 60, 110, 121	0
2	K	42/43 (97%)	-1.51	0 100 100	35, 62, 121, 130	0
2	L	42/43 (97%)	-1.56	0 100 100	29, 57, 114, 139	0
2	M	42/43 (97%)	-1.52	0 100 100	22, 53, 111, 134	0
2	N	42/43 (97%)	-1.50	0 100 100	30, 57, 116, 130	0
2	O	41/43 (95%)	-1.54	0 100 100	36, 54, 108, 124	0
2	P	42/43 (97%)	-1.47	0 100 100	33, 55, 111, 125	0
All	All	2883/3160 (91%)	-1.71	0 100 100	18, 46, 78, 139	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	NAP	A	1353	48/48	0.99	0.03	36,54,98,105	0
3	NAP	B	1353	48/48	0.99	0.04	25,55,87,91	0
3	NAP	C	1353	48/48	0.99	0.03	31,57,96,101	0
3	NAP	D	1353	48/48	0.99	0.04	25,53,74,85	0
3	NAP	E	1353	48/48	0.99	0.04	20,48,80,87	0
3	NAP	F	1353	48/48	0.99	0.03	25,49,80,87	0
3	NAP	G	1353	48/48	0.99	0.03	27,51,78,99	0
3	NAP	H	1353	48/48	1.00	0.03	22,61,106,110	0
4	ZN	I	1713	1/1	1.00	0.01	58,58,58,58	0
4	ZN	J	1713	1/1	1.00	0.01	57,57,57,57	0
4	ZN	K	1713	1/1	1.00	0.02	66,66,66,66	0
4	ZN	L	1713	1/1	1.00	0.01	58,58,58,58	0
4	ZN	M	1712	1/1	1.00	0.00	48,48,48,48	0
4	ZN	N	1713	1/1	1.00	0.00	53,53,53,53	0
4	ZN	O	1712	1/1	1.00	0.01	52,52,52,52	0
4	ZN	P	1713	1/1	1.00	0.01	47,47,47,47	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around NAP A 1353:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP B 1353:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP C 1353:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP D 1353:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP E 1353:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around NAP F 1353:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around NAP G 1353:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around NAP H 1353:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.