



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 08:02 PM UTC

PDB ID : 3VV6 / pdb_00003vv6
Title : Crystal Structure of beta secetase in complex with 2-amino-3-methyl-6-((1S, 2R)-2-phenylcyclopropyl)pyrimidin-4(3H)-one
Authors : Yonezawa, S.; Yamamoto, T.; Yamakawa, H.; Muto, C.; Hosono, M.; Hattori, K.; Higashino, K.; Sakagami, M.; Togame, H.; Tanaka, Y.; Nakano, T.; Takemoto, H.; Arisawa, M.; Shuto, S.
Deposited on : 2012-07-17
Resolution : 2.05 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.05 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2260 (2.04-2.04)
Clashscore	190562	2333 (2.04-2.04)
Ramachandran outliers	187476	2318 (2.04-2.04)
Sidechain outliers	187428	2318 (2.04-2.04)
RSRZ outliers	180081	2260 (2.04-2.04)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	416	

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 3030 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Beta-secretase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	360	Total	C	N	O	S	0	0	0
			2773	1778	458	523	14			


There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-22	GLY	-	expression tag	UNP P56817
A	-21	SER	-	expression tag	UNP P56817
A	-20	HIS	-	expression tag	UNP P56817
A	-19	MET	-	expression tag	UNP P56817

- Molecule 2 is IODIDE ION (CCD ID: IOD) (formula: I).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	5	Total	I	0	0
			5	5		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		
3	A	1	Total	C	O	0	0
			6	3	3		

- Molecule 4 is 2-amino-3-methyl-6-[(1S,2R)-2-phenylcyclopropyl]pyrimidin-4(3H)-one (CCD ID: B00) (formula: C₁₄H₁₅N₃O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			18	14	3	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	210	Total	O	0	0
			210	210		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Beta-secretase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants a, b, c, α , β , γ	102.49Å 102.49Å 170.16Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	20.00 – 2.05 20.00 – 2.05	Depositor EDS
% Data completeness (in resolution range)	99.4 (20.00-2.05) 99.2 (20.00-2.05)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.28 (at 2.05Å)	Xtriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.249 , 0.280 0.250 , 0.281	Depositor DCC
R_{free} test set	1706 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.2	Xtriage
Anisotropy	0.095	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 45.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	3030	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.68% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, GOL, B00

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.80	1/2838 (0.0%)	0.89	6/3857 (0.2%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	212	ASP	C-O	-5.08	1.17	1.23

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	69	VAL	N-CA-C	6.57	114.65	108.15
1	A	204	VAL	CB-CA-C	5.99	116.52	110.65
1	A	208	ILE	O-C-N	5.48	129.01	123.20
1	A	236	LEU	CA-C-N	-5.04	114.58	119.78
1	A	236	LEU	C-N-CA	-5.04	114.58	119.78
1	A	178	GLY	N-CA-C	5.01	117.86	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2773	0	2642	57	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	5	0	0	2	0
3	A	24	0	32	2	0
4	A	18	0	15	2	0
5	A	210	0	0	1	0
All	All	3030	0	2689	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:502:IOD:I	5:A:1203:HOH:O	2.28	1.21
1:A:209:ASN:HD21	1:A:281:PRO:HA	1.11	1.15
1:A:5:ASN:HD21	1:A:174:MET:H	1.14	0.92
1:A:209:ASN:HD21	1:A:281:PRO:CA	1.84	0.90
1:A:25:GLN:HE22	1:A:53:GLN:H	1.17	0.89
1:A:303:GLN:H	1:A:303:GLN:HE21	1.16	0.88
1:A:209:ASN:OD1	1:A:281:PRO:HB3	1.77	0.83
1:A:148:ASN:HD21	1:A:347:ARG:HE	1.31	0.78
1:A:209:ASN:ND2	1:A:281:PRO:HA	1.96	0.78
1:A:204:VAL:HG13	1:A:376:THR:HG21	1.68	0.76
1:A:209:ASN:ND2	1:A:281:PRO:CA	2.55	0.66
1:A:238:LYS:CB	2:A:504:IOD:I	3.14	0.66
1:A:55:GLN:HE21	1:A:55:GLN:H	1.45	0.65
1:A:195:ARG:HG2	1:A:197:TRP:CD1	2.32	0.64
1:A:301:LEU:N	1:A:301:LEU:HD12	2.15	0.62
1:A:181:HIS:HB3	3:A:508:GOL:H12	1.80	0.61
1:A:244:ALA:O	1:A:248:ILE:HG13	2.01	0.61
1:A:204:VAL:CG1	1:A:376:THR:HG21	2.29	0.60
1:A:278:ASN:OD1	1:A:280:PHE:HB2	2.06	0.56
1:A:174:MET:HE3	1:A:176:ILE:HD11	1.89	0.54
1:A:236:LEU:O	1:A:326:GLN:HA	2.08	0.54
1:A:188:LEU:HD23	1:A:355:ALA:HB2	1.91	0.53
1:A:213:LEU:N	1:A:213:LEU:HD23	2.23	0.53
1:A:241:PHE:CD2	1:A:326:GLN:HG2	2.44	0.53
1:A:209:ASN:CG	1:A:281:PRO:HB3	2.32	0.53
1:A:208:ILE:HG23	1:A:281:PRO:HB2	1.91	0.53
1:A:279:ILE:HG12	1:A:279:ILE:O	2.09	0.52
1:A:184:TYR:CE1	3:A:508:GOL:H32	2.45	0.52
1:A:209:ASN:ND2	1:A:282:VAL:H	2.07	0.52

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:149:LEU:C	1:A:149:LEU:HD12	2.36	0.51
1:A:303:GLN:HE21	1:A:303:GLN:N	1.97	0.51
1:A:68:TYR:CD2	1:A:70:PRO:HD3	2.46	0.51
1:A:241:PHE:CZ	1:A:245:VAL:HG21	2.45	0.50
1:A:235:ARG:O	1:A:331:THR:HA	2.11	0.50
1:A:180:ASP:HB3	1:A:183:LEU:HD22	1.94	0.50
1:A:209:ASN:ND2	1:A:281:PRO:CB	2.74	0.50
1:A:304:GLN:O	1:A:336:VAL:HG22	2.14	0.48
1:A:254:THR:HB	1:A:279:ILE:HD12	1.95	0.48
1:A:59:THR:HG21	1:A:84:LEU:HD22	1.95	0.48
1:A:208:ILE:C	1:A:210:GLY:H	2.22	0.47
1:A:2:MET:HG2	1:A:89:HIS:O	2.15	0.47
1:A:209:ASN:CG	1:A:281:PRO:CB	2.88	0.46
1:A:234:LEU:O	1:A:324:ILE:HA	2.16	0.46
1:A:302:PRO:O	1:A:306:LEU:HB2	2.15	0.46
1:A:301:LEU:HD13	1:A:304:GLN:OE1	2.16	0.46
1:A:1:GLU:HG2	1:A:2:MET:HG3	1.98	0.45
1:A:148:ASN:ND2	1:A:347:ARG:HE	2.06	0.45
1:A:290:GLU:H	1:A:294:GLN:HE22	1.64	0.45
1:A:118:ILE:HD12	4:A:510:B00:C13	2.47	0.44
1:A:224:LYS:O	1:A:331:THR:N	2.49	0.44
4:A:510:B00:H6	4:A:510:B00:H8	1.80	0.43
1:A:125:GLU:OE1	1:A:195:ARG:NH1	2.52	0.43
1:A:301:LEU:N	1:A:301:LEU:CD1	2.82	0.42
1:A:197:TRP:CD1	1:A:197:TRP:N	2.88	0.42
1:A:205:ARG:HB3	1:A:286:TYR:HB2	2.01	0.42
1:A:199:TYR:HB3	1:A:352:ILE:HD11	2.02	0.41
1:A:233:ASN:HB3	1:A:323:ALA:O	2.20	0.41
1:A:197:TRP:CZ2	1:A:224:LYS:HD2	2.56	0.41
1:A:55:GLN:H	1:A:55:GLN:NE2	2.15	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	350/416 (84%)	340 (97%)	8 (2%)	2 (1%)	21	13

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	209	ASN
1	A	210	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	292/356 (82%)	264 (90%)	28 (10%)	8	3

All (28) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	30	LEU
1	A	55	GLN
1	A	56	LEU
1	A	63	LEU
1	A	75	LYS
1	A	84	LEU
1	A	128	ARG
1	A	149	LEU
1	A	154	LEU
1	A	183	LEU
1	A	204	VAL
1	A	208	ILE
1	A	213	LEU
1	A	224	LYS
1	A	252	SER
1	A	254	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	263	LEU
1	A	267	LEU
1	A	278	ASN
1	A	279	ILE
1	A	287	LEU
1	A	301	LEU
1	A	303	GLN
1	A	306	LEU
1	A	307	ARG
1	A	377	LEU
1	A	380	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (15) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	25	GLN
1	A	28	ASN
1	A	55	GLN
1	A	89	HIS
1	A	92	ASN
1	A	111	ASN
1	A	114	ASN
1	A	148	ASN
1	A	209	ASN
1	A	233	ASN
1	A	294	GLN
1	A	303	GLN
1	A	326	GLN
1	A	385	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 5 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	GOL	A	508	-	5,5,5	0.38	0	5,5,5	0.31	0
4	B00	A	510	-	19,20,20	1.47	2 (10%)	24,29,29	2.35	9 (37%)
3	GOL	A	509	-	5,5,5	0.34	0	5,5,5	0.36	0
3	GOL	A	506	-	5,5,5	0.50	0	5,5,5	0.35	0
3	GOL	A	507	-	5,5,5	0.39	0	5,5,5	0.26	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	A	508	-	-	2/4/4/4	-
4	B00	A	510	-	-	0/4/13/13	0/3/3/3
3	GOL	A	509	-	-	4/4/4/4	-
3	GOL	A	506	-	-	0/4/4/4	-
3	GOL	A	507	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	510	B00	C4-N2	3.27	1.43	1.37
4	A	510	B00	C2-C3	2.91	1.40	1.36

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	510	B00	C14-C6-C3	-6.21	112.01	119.64
4	A	510	B00	C6-C3-C2	-4.37	120.53	125.42
4	A	510	B00	C2-C1-N2	4.16	119.27	115.36
4	A	510	B00	C14-C7-C8	-4.04	114.70	122.27
4	A	510	B00	C2-C3-N1	-3.54	119.52	125.07
4	A	510	B00	N3-C4-N2	2.91	121.13	118.79
4	A	510	B00	O-C1-C2	-2.27	120.01	125.61
4	A	510	B00	C8-C7-C6	-2.15	116.48	120.67
4	A	510	B00	C4-N2-C1	-2.14	118.97	121.07

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	507	GOL	O1-C1-C2-C3
3	A	509	GOL	O1-C1-C2-C3
3	A	509	GOL	C1-C2-C3-O3
3	A	507	GOL	O1-C1-C2-O2
3	A	509	GOL	O1-C1-C2-O2
3	A	509	GOL	O2-C2-C3-O3
3	A	507	GOL	O2-C2-C3-O3
3	A	507	GOL	C1-C2-C3-O3
3	A	508	GOL	O2-C2-C3-O3
3	A	508	GOL	C1-C2-C3-O3

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	508	GOL	2	0
4	A	510	B00	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	360/416 (86%)	1.01	87 (24%) 2 1	14, 28, 50, 57	0

All (87) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	319	CYS	9.7
1	A	268	VAL	6.4
1	A	365	PHE	5.9
1	A	245	VAL	5.8
1	A	280	PHE	5.6
1	A	322	PHE	5.6
1	A	209	ASN	5.4
1	A	262	TRP	5.1
1	A	236	LEU	5.0
1	A	264	GLY	4.9
1	A	302	PRO	4.8
1	A	252	SER	4.6
1	A	222	TYR	4.4
1	A	279	ILE	4.4
1	A	260	GLY	4.4
1	A	257	PHE	4.3
1	A	301	LEU	4.2
1	A	384	TYR	4.1
1	A	250	ALA	4.1
1	A	306	LEU	3.9
1	A	320	TYR	3.9
1	A	328	SER	3.9
1	A	253	SER	3.9
1	A	259	ASP	3.8
1	A	267	LEU	3.8
1	A	305	TYR	3.8
1	A	244	ALA	3.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	269	CYS	3.8
1	A	282	VAL	3.7
1	A	261	PHE	3.7
1	A	324	ILE	3.7
1	A	214	LYS	3.7
1	A	318	ASP	3.6
1	A	308	PRO	3.6
1	A	238	LYS	3.6
1	A	265	GLU	3.5
1	A	215	MET	3.5
1	A	251	ALA	3.5
1	A	211	GLN	3.5
1	A	254	THR	3.4
1	A	243	ALA	3.4
1	A	383	GLY	3.4
1	A	241	PHE	3.4
1	A	367	THR	3.4
1	A	72	THR	3.3
1	A	237	PRO	3.3
1	A	307	ARG	3.2
1	A	263	LEU	3.1
1	A	363	ASP	3.1
1	A	216	ASP	3.0
1	A	240	VAL	3.0
1	A	278	ASN	2.9
1	A	213	LEU	2.9
1	A	155	CYS	2.9
1	A	95	VAL	2.9
1	A	73	GLN	2.8
1	A	309	VAL	2.8
1	A	303	GLN	2.8
1	A	327	SER	2.7
1	A	234	LEU	2.7
1	A	248	ILE	2.7
1	A	361	VAL	2.6
1	A	323	ALA	2.6
1	A	281	PRO	2.6
1	A	325	SER	2.6
1	A	304	GLN	2.6
1	A	71	TYR	2.5
1	A	69	VAL	2.5
1	A	329	THR	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	212	ASP	2.4
1	A	366	ARG	2.3
1	A	-2	SER	2.3
1	A	321	LYS	2.3
1	A	145	HIS	2.3
1	A	-3	GLY	2.3
1	A	235	ARG	2.2
1	A	156	GLY	2.2
1	A	219	GLU	2.2
1	A	242	GLU	2.2
1	A	326	GLN	2.2
1	A	233	ASN	2.1
1	A	362	HIS	2.1
1	A	381	ASP	2.1
1	A	247	SER	2.1
1	A	385	ASN	2.1
1	A	300	ILE	2.0
1	A	210	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	GOL	A	509	6/6	0.72	0.15	58,59,59,59	0
3	GOL	A	507	6/6	0.76	0.17	50,51,53,55	0
3	GOL	A	508	6/6	0.78	0.19	48,49,49,51	0
4	B00	A	510	18/18	0.83	0.14	19,36,47,47	0
3	GOL	A	506	6/6	0.91	0.11	25,32,34,36	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	IOD	A	505	1/1	0.94	0.12	101,101,101,101	0
2	IOD	A	503	1/1	0.95	0.18	61,61,61,61	0
2	IOD	A	502	1/1	0.97	0.11	50,50,50,50	0
2	IOD	A	504	1/1	0.97	0.26	54,54,54,54	0
2	IOD	A	501	1/1	1.00	0.03	25,25,25,25	0

6.5 Other polymers [i](#)

There are no such residues in this entry.