



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 26, 2026 – 09:09 AM EDT

PDB ID : 7VVD / pdb_00007vvd
Title : Crystal Structure of the Kv7.1 C-terminal Domain in Complex with Calmodulin disease mutation Q135P
Authors : Chen, L.
Deposited on : 2021-11-05
Resolution : 3.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION


The reported resolution of this entry is 3.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2351 (3.18-3.10)
Clashscore	190562	2452 (3.18-3.10)
Ramachandran outliers	187476	2324 (3.18-3.10)
Sidechain outliers	187428	2324 (3.18-3.10)
RSRZ outliers	180081	2351 (3.18-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	69	
1	D	69	
2	C	149	
2	E	149	

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 2966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium voltage-gated channel subfamily KQT member 1, Potassium voltage-gated channel subfamily KQT member 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	64	Total	C	N	O	S	0	0	0
			500	316	98	84	2			
1	D	60	Total	C	N	O	S	0	0	0
			455	290	87	76	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	534	ILE	-	expression tag	UNP P51787
A	535	GLY	-	expression tag	UNP P51787
A	536	SER	-	expression tag	UNP P51787
A	537	GLY	-	expression tag	UNP P51787
D	534	ILE	-	expression tag	UNP P51787
D	535	GLY	-	expression tag	UNP P51787
D	536	SER	-	expression tag	UNP P51787
D	537	GLY	-	expression tag	UNP P51787

- Molecule 2 is a protein called Calmodulin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	145	Total	C	N	O	S	0	0	0
			1062	653	171	230	8			
2	E	140	Total	C	N	O	S	0	0	0
			943	579	155	201	8			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	135	PRO	GLN	engineered mutation	UNP P0DP23
E	135	PRO	GLN	engineered mutation	UNP P0DP23

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	C	2	Total 2	Ca 2	0	0
3	E	2	Total 2	Ca 2	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	C	1	Total 1	O 1	0	0
4	E	1	Total 1	O 1	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1, Potassium voltage-gated channel subfamily KQT member 1

Chain A: 




- Molecule 1: Potassium voltage-gated channel subfamily KQT member 1, Potassium voltage-gated channel subfamily KQT member 1

Chain D: 



- Molecule 2: Calmodulin-1

Chain C: 



- Molecule 2: Calmodulin-1

Chain E: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	40.51Å 86.27Å 126.12Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.79 – 3.13 37.79 – 3.13	Depositor EDS
% Data completeness (in resolution range)	91.1 (37.79-3.13) 87.9 (37.79-3.13)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.28 (at 3.12Å)	Xtriage
Refinement program	PHENIX 1.14_3247	Depositor
R, R_{free}	0.243 , 0.274 0.243 , 0.274	Depositor DCC
R_{free} test set	752 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å ²)	66.8	Xtriage
Anisotropy	0.217	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 33.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	2966	wwPDB-VP
Average B, all atoms (Å ²)	57.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.19% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.14	0/512	0.37	0/694
1	D	0.64	2/466 (0.4%)	0.42	0/632
2	C	0.43	1/1074 (0.1%)	0.45	1/1456 (0.1%)
2	E	0.48	2/952 (0.2%)	0.47	1/1293 (0.1%)
All	All	0.45	5/3004 (0.2%)	0.44	2/4075 (0.0%)

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	D	376	GLN	C-O	-6.61	1.16	1.24
2	C	114	GLU	C-O	-6.06	1.15	1.24
2	E	115	LYS	CA-C	-5.47	1.45	1.52
2	E	115	LYS	C-O	-5.36	1.17	1.24
1	D	375	ILE	C-O	-5.02	1.18	1.24

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	115	LYS	N-CA-C	8.34	128.56	110.80
2	E	116	LEU	N-CA-C	5.61	122.74	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	500	0	465	7	0
1	D	455	0	412	6	0
2	C	1062	0	911	13	0
2	E	943	0	742	20	0
3	C	2	0	0	0	0
3	E	2	0	0	0	0
4	C	1	0	0	0	0
4	E	1	0	0	0	0
All	All	2966	0	2530	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:117:THR:HG22	2:E:120:GLU:H	1.09	1.07
2:E:117:THR:CG2	2:E:120:GLU:H	1.85	0.89
2:E:117:THR:HG22	2:E:120:GLU:N	1.92	0.84
2:C:82:GLU:HG2	2:C:146:THR:HG21	1.67	0.77
1:D:373:SER:O	1:D:377:THR:OG1	2.07	0.72
2:C:44:THR:HG22	2:C:46:ALA:H	1.54	0.71
2:E:117:THR:HG23	2:E:119:GLU:H	1.56	0.69
2:E:117:THR:HG23	2:E:119:GLU:N	2.13	0.62
1:A:376:GLN:HE21	2:C:109:MET:HA	1.66	0.60
2:C:60:ASN:ND2	2:C:62:THR:OG1	2.35	0.59
2:C:55:VAL:HG12	2:C:71:MET:HG3	1.86	0.57
2:E:109:MET:HG3	2:E:124:MET:HE1	1.87	0.56
1:A:373:SER:OG	2:C:114:GLU:OE2	2.21	0.53
1:D:389:SER:OG	2:E:123:GLU:OE2	2.24	0.51
2:C:106:ARG:NH2	2:C:122:ASP:OD1	2.40	0.51
2:E:55:VAL:HG13	2:E:71:MET:HB2	1.93	0.51
1:D:512:ALA:O	1:D:516:VAL:HG23	2.12	0.50
2:E:55:VAL:HG22	2:E:71:MET:HG3	1.95	0.49
1:D:394:ILE:HG22	1:D:395:TYR:H	1.77	0.48
2:E:5:THR:HG22	2:E:7:GLU:H	1.78	0.48
1:A:512:ALA:O	1:A:516:VAL:HG23	2.14	0.48
2:E:31:GLU:O	2:E:35:VAL:HG23	2.13	0.48
2:E:5:THR:HG22	2:E:7:GLU:N	2.28	0.48
2:E:101:SER:HA	2:E:135:PRO:HB3	1.96	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:C:104:GLU:O	2:C:108:VAL:HG22	2.15	0.47
2:C:22:ASP:OD1	2:C:22:ASP:N	2.36	0.47
1:D:520:MET:HE2	2:E:51:MET:HB3	1.97	0.47
1:A:523:PHE:HE2	2:C:71:MET:HG2	1.82	0.45
2:E:71:MET:HE3	2:E:71:MET:HB3	1.88	0.43
2:E:78:ASP:OD1	2:E:79:THR:N	2.52	0.43
1:A:372:ALA:O	1:A:376:GLN:HG3	2.18	0.43
1:D:379:TRP:NE1	2:E:124:MET:HG3	2.34	0.42
2:E:117:THR:HG22	2:E:120:GLU:HB2	2.01	0.42
1:A:523:PHE:HB3	2:C:54:GLU:HG2	2.00	0.42
1:A:521:GLN:HG2	2:C:51:MET:HE2	2.02	0.42
2:E:139:GLU:HA	2:E:142:VAL:HG12	2.01	0.41
2:C:43:PRO:HG2	2:C:48:LEU:HD11	2.01	0.41
2:E:19:PHE:CD1	2:E:35:VAL:HG21	2.56	0.41

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	62/69 (90%)	60 (97%)	2 (3%)	0	100	100
1	D	56/69 (81%)	55 (98%)	1 (2%)	0	100	100
2	C	141/149 (95%)	132 (94%)	7 (5%)	2 (1%)	9	31
2	E	134/149 (90%)	127 (95%)	5 (4%)	2 (2%)	8	29
All	All	393/436 (90%)	374 (95%)	15 (4%)	4 (1%)	12	38

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	C	77	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	E	116	LEU
2	C	129	ASP
2	E	132	GLY

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	44/58 (76%)	44 (100%)	0	100	100
1	D	38/58 (66%)	37 (97%)	1 (3%)	40	64
2	C	101/127 (80%)	101 (100%)	0	100	100
2	E	76/127 (60%)	75 (99%)	1 (1%)	61	73
All	All	259/370 (70%)	257 (99%)	2 (1%)	73	78

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	377	THR
2	E	117	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	521	GLN
2	C	111	ASN
1	D	510	HIS
1	D	521	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	64/69 (92%)	-0.16	0	100 100	40, 48, 64, 70	0
1	D	60/69 (86%)	0.22	0	100 100	45, 60, 75, 82	0
2	C	145/149 (97%)	-0.10	1 (0%)	84 68	42, 52, 75, 82	0
2	E	140/149 (93%)	0.08	1 (0%)	84 68	45, 61, 82, 95	0
All	All	409/436 (93%)	0.00	2 (0%)	87 73	40, 55, 79, 95	0

All (2) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	E	127	GLU	2.7
2	C	76	MET	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	C	201	1/1	0.96	0.04	55,55,55,55	0
3	CA	C	202	1/1	0.98	0.04	55,55,55,55	0
3	CA	E	201	1/1	0.99	0.02	56,56,56,56	0
3	CA	E	202	1/1	0.99	0.09	55,55,55,55	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around CA C 202:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around CA E 201:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers ⓘ

There are no such residues in this entry.