



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:21 AM EDT

PDB ID : 2VWS / pdb_00002vws
Title : Crystal structure of YfaU, a metal ion dependent class II aldolase from Escherichia coli K12
Authors : Rea, D.; Rakus, J.F.; Gerlt, J.A.; Fulop, V.; Bugg, T.D.H.; Roper, D.I.
Deposited on : 2008-06-26
Resolution : 1.39 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.39 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2563 (1.40-1.40)
Clashscore	190562	2660 (1.40-1.40)
Ramachandran outliers	187476	2611 (1.40-1.40)
Sidechain outliers	187428	2610 (1.40-1.40)
RSRZ outliers	180081	2561 (1.40-1.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	267	21% 85% 9% ••
1	B	267	28% 87% 9% ••
1	C	267	18% 90% 6% •

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	PO4	C	1257	-	X	-	-

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 6845 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	0	0
			1951	1233	338	371	9			
1	B	256	Total	C	N	O	S	0	1	0
			1952	1233	338	371	10			
1	C	256	Total	C	N	O	S	0	0	0
			1951	1233	338	371	9			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	C	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	C	1	Total	C	O	0	0
			6	3	3		
3	C	1	Total	C	O	0	0
			6	3	3		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	320	Total	O	0	0
			320	320		
4	B	321	Total	O	0	0
			321	321		
4	C	318	Total	O	0	0
			318	318		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



• Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



• Molecule 1: YFAU, 2-KETO-3-DEOXY SUGAR ALDOLASE



4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, α , β , γ	105.20Å 136.80Å 123.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	59.76 – 1.39 59.76 – 1.39	Depositor EDS
% Data completeness (in resolution range)	99.5 (59.76-1.39) 99.5 (59.76-1.39)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.64 (at 1.39Å)	Xtriage
Refinement program	REFMAC 5.1.9999	Depositor
R, R_{free}	0.171 , 0.191 0.171 , 0.190	Depositor DCC
R_{free} test set	7041 reflections (3.99%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.305	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 41.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	6845	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.45% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PO4, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.82	0/1988	0.92	2/2706 (0.1%)
1	B	0.72	0/1994	0.88	1/2714 (0.0%)
1	C	0.77	0/1988	0.91	2/2706 (0.1%)
All	All	0.77	0/5970	0.90	5/8126 (0.1%)

There are no bond length outliers.

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	192	HIS	N-CA-C	7.65	126.72	109.81
1	C	188	ASP	N-CA-C	6.28	118.01	111.03
1	B	142	VAL	N-CA-C	5.35	115.55	110.42
1	C	179	ASP	N-CA-C	-5.12	105.59	111.07
1	A	255	LYS	N-CA-C	5.12	115.05	108.34

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1951	0	1950	19	0
1	B	1952	0	1951	17	0
1	C	1951	0	1950	8	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	A	10	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
3	C	12	0	16	0	0
4	A	320	0	0	2	0
4	B	321	0	0	5	0
4	C	318	0	0	4	0
All	All	6845	0	5867	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:105:ALA:HB1	1:B:170:ILE:HD11	1.40	1.01
1:A:105:ALA:HB1	1:A:170:ILE:HD11	1.56	0.87
1:A:122:VAL:HG23	4:C:2071:HOH:O	1.81	0.79
1:B:181:SER:OG	1:B:190:ALA:HA	1.86	0.76
1:B:179:ASP:HB2	4:B:2230:HOH:O	1.85	0.75
1:A:222:MET:HE3	1:A:226:CYS:SG	2.35	0.67
1:B:13:LEU:HD12	4:B:2027:HOH:O	1.95	0.67
1:B:222:MET:HE2	1:B:226[B]:CYS:SG	2.37	0.65
1:B:80:LYS:HG2	4:B:2131:HOH:O	2.02	0.60
1:A:187:PRO:HD2	1:A:189:ASN:HD21	1.68	0.58
1:C:158:LEU:HD11	1:C:180:LEU:HD11	1.85	0.58
1:B:153:GLU:HA	1:B:153:GLU:OE1	2.04	0.57
1:A:241:MET:HG2	4:B:2180:HOH:O	2.05	0.56
1:A:153:GLU:HB3	4:A:2221:HOH:O	2.04	0.56
1:B:222:MET:CE	1:B:226[B]:CYS:SG	2.95	0.54
1:A:189:ASN:ND2	1:A:189:ASN:H	2.06	0.54
1:B:80:LYS:HB2	1:B:81:PRO:HD3	1.90	0.54
1:C:158:LEU:CD1	1:C:180:LEU:HD11	2.39	0.53
1:C:205:ARG:HG3	4:C:2253:HOH:O	2.09	0.52
1:A:102:ALA:HB2	1:A:163:GLU:HB3	1.93	0.51
1:B:47:GLY:O	1:B:76:VAL:HG22	2.10	0.51
1:C:152:VAL:CG2	1:C:175:ILE:HG12	2.41	0.50
1:A:115:PRO:HG2	4:C:2071:HOH:O	2.09	0.50
1:B:105:ALA:CB	1:B:170:ILE:HD11	2.28	0.49
1:B:124:ALA:HB1	1:B:129:ALA:HB3	1.96	0.48
1:C:47:GLY:O	1:C:76:VAL:HG22	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:124:ALA:HB1	1:A:129:ALA:HB3	1.96	0.48
1:A:176:GLY:HA3	4:A:2223:HOH:O	2.14	0.48
1:A:182:ALA:HA	1:A:187:PRO:HA	1.95	0.48
1:A:152:VAL:HG13	1:A:157:ALA:HB1	1.95	0.47
1:A:3:ALA:HB1	1:A:136:GLU:HG2	1.96	0.46
1:C:80:LYS:HE3	1:C:119:GLU:HB3	1.98	0.46
1:C:76:VAL:HG23	4:C:2114:HOH:O	2.18	0.43
1:A:187:PRO:O	1:A:188:ASP:HB2	2.18	0.43
1:A:189:ASN:ND2	1:A:189:ASN:N	2.66	0.43
1:B:181:SER:HB3	1:B:186:TYR:O	2.19	0.43
1:B:80:LYS:HB2	1:B:81:PRO:CD	2.49	0.43
1:B:249:GLN:NE2	4:B:2307:HOH:O	2.40	0.42
1:A:158:LEU:HD11	1:A:198:ILE:HG21	2.01	0.42
1:A:44:LEU:HD13	1:A:44:LEU:C	2.45	0.41
1:B:110:SER:OG	1:B:119:GLU:HG3	2.20	0.41
1:B:131:ARG:O	1:B:134:ARG:HD3	2.21	0.41
1:A:152:VAL:HG13	1:A:157:ALA:CB	2.51	0.41
1:C:152:VAL:HG23	1:C:175:ILE:HG12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	254/267 (95%)	250 (98%)	2 (1%)	2 (1%)	16	3
1	B	255/267 (96%)	251 (98%)	4 (2%)	0	100	100
1	C	254/267 (95%)	246 (97%)	5 (2%)	3 (1%)	10	1
All	All	763/801 (95%)	747 (98%)	11 (1%)	5 (1%)	18	4

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	188	ASP
1	C	190	ALA
1	C	123	GLY
1	C	191	GLY
1	A	192	HIS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	203/211 (96%)	198 (98%)	5 (2%)	42	11
1	B	204/211 (97%)	199 (98%)	5 (2%)	42	11
1	C	203/211 (96%)	202 (100%)	1 (0%)	81	61
All	All	610/633 (96%)	599 (98%)	11 (2%)	51	20

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	4	LEU
1	A	73	ILE
1	A	136	GLU
1	A	155	LYS
1	A	189	ASN
1	B	119	GLU
1	B	125	SER
1	B	155	LYS
1	B	179	ASP
1	B	188	ASP
1	C	225	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	189	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	225	GLN
1	B	62	GLN
1	B	189	ASN
1	B	225	GLN
1	B	249	GLN
1	C	62	GLN
1	C	70	GLN
1	C	196	GLN
1	C	225	GLN
1	C	249	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	GOL	C	1259	-	5,5,5	0.44	0	5,5,5	0.46	0
3	GOL	C	1258	-	5,5,5	0.20	0	5,5,5	0.71	0
2	PO4	B	1257	-	4,4,4	2.05	1 (25%)	6,6,6	2.42	2 (33%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	C	1257	-	4,4,4	1.85	1 (25%)	6,6,6	2.68	3 (50%)
2	PO4	A	1257	-	4,4,4	1.67	1 (25%)	6,6,6	2.13	2 (33%)
2	PO4	A	1258	-	4,4,4	0.84	0	6,6,6	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	C	1258	-	-	2/4/4/4	-
3	GOL	C	1259	-	-	0/4/4/4	-

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1257	PO4	P-O1	3.91	1.59	1.50
2	C	1257	PO4	P-O1	3.46	1.58	1.50
2	A	1257	PO4	P-O1	3.30	1.58	1.50

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	1257	PO4	O2-P-O1	-4.26	95.88	110.95
2	C	1257	PO4	O4-P-O1	4.17	125.71	110.95
2	B	1257	PO4	O4-P-O1	4.04	125.23	110.95
2	B	1257	PO4	O2-P-O1	-3.74	97.73	110.95
2	A	1257	PO4	O4-P-O1	3.47	123.22	110.95
2	A	1257	PO4	O2-P-O1	-3.35	99.09	110.95
2	C	1257	PO4	O3-P-O2	2.38	115.30	107.91

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	1258	GOL	C1-C2-C3-O3
3	C	1258	GOL	O2-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	256/267 (95%)	1.48	57 (22%) 2 2	13, 19, 31, 50	0
1	B	256/267 (95%)	1.48	75 (29%) 1 1	13, 20, 34, 40	1 (0%)
1	C	256/267 (95%)	1.27	47 (18%) 3 3	14, 19, 29, 53	0
All	All	768/801 (95%)	1.41	179 (23%) 2 1	13, 20, 33, 53	1 (0%)

All (179) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	190	ALA	15.6
1	C	190	ALA	9.2
1	A	189	ASN	9.0
1	B	117	TYR	8.2
1	C	189	ASN	7.9
1	A	191	GLY	7.6
1	C	187	PRO	7.4
1	A	187	PRO	7.3
1	A	116	PRO	6.5
1	A	186	TYR	6.2
1	A	126	VAL	6.1
1	C	191	GLY	6.0
1	A	117	TYR	6.0
1	B	3	ALA	5.9
1	A	192	HIS	5.8
1	A	125	SER	5.7
1	A	188	ASP	5.6
1	B	256	SER	5.6
1	C	256	SER	5.4
1	A	3	ALA	5.1
1	C	126	VAL	4.8
1	B	116	PRO	4.6
1	B	187	PRO	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	188	ASP	4.4
1	A	256	SER	4.4
1	A	156	THR	4.2
1	B	2	ASN	4.2
1	C	229	TRP	4.2
1	A	2	ASN	4.1
1	B	4	LEU	4.1
1	A	1	MET	4.1
1	A	16	GLY	4.0
1	A	183	SER	3.9
1	A	115	PRO	3.8
1	B	126	VAL	3.7
1	B	255	LYS	3.7
1	A	184	LEU	3.7
1	B	209	ALA	3.7
1	B	125	SER	3.6
1	C	122	VAL	3.6
1	B	122	VAL	3.5
1	A	124	ALA	3.4
1	B	124	ALA	3.4
1	C	180	LEU	3.4
1	C	117	TYR	3.4
1	B	1	MET	3.3
1	A	193	PRO	3.2
1	C	2	ASN	3.2
1	B	144	ASP	3.2
1	C	221	ASP	3.2
1	A	198	ILE	3.1
1	B	118	GLY	3.1
1	B	14	ARG	3.1
1	A	119	GLU	3.1
1	B	6	SER	3.1
1	B	140	ALA	3.1
1	A	208	ALA	3.0
1	C	228	ALA	3.0
1	A	136	GLU	3.0
1	B	30	TYR	3.0
1	B	114	TYR	3.0
1	A	195	VAL	3.0
1	B	143	ASN	3.0
1	A	185	GLY	3.0
1	B	139	MET	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	155	LYS	2.9
1	C	199	ILE	2.9
1	B	137	ASN	2.9
1	B	170	ILE	2.8
1	C	226	CYS	2.8
1	A	180	LEU	2.8
1	B	57	LEU	2.8
1	B	153	GLU	2.8
1	C	238	VAL	2.7
1	C	116	PRO	2.7
1	C	186	TYR	2.7
1	B	115	PRO	2.7
1	A	166	ASP	2.7
1	B	32	ALA	2.7
1	A	221	ASP	2.7
1	B	135	ILE	2.6
1	B	158	LEU	2.6
1	B	28	THR	2.6
1	C	3	ALA	2.6
1	B	188	ASP	2.6
1	B	119	GLU	2.6
1	A	4	LEU	2.6
1	C	240	THR	2.6
1	B	36	ALA	2.6
1	B	50	ALA	2.6
1	B	190	ALA	2.6
1	A	181	SER	2.6
1	A	255	LYS	2.6
1	C	255	LYS	2.6
1	C	1	MET	2.6
1	B	24	LEU	2.5
1	C	205	ARG	2.5
1	B	141	GLN	2.5
1	B	142	VAL	2.5
1	B	218	VAL	2.5
1	C	152	VAL	2.5
1	C	115	PRO	2.5
1	A	153	GLU	2.5
1	A	194	GLU	2.4
1	B	208	ALA	2.4
1	B	138	TYR	2.4
1	C	82	LEU	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	148	LEU	2.4
1	B	191	GLY	2.4
1	C	195	VAL	2.4
1	B	179	ASP	2.4
1	B	198	ILE	2.4
1	A	162	ASP	2.4
1	C	144	ASP	2.4
1	C	4	LEU	2.4
1	B	197	ARG	2.3
1	C	54	ILE	2.3
1	B	40	TYR	2.3
1	A	122	VAL	2.3
1	B	130	ALA	2.3
1	C	192	HIS	2.3
1	B	64	VAL	2.3
1	C	11	GLU	2.3
1	A	175	ILE	2.3
1	B	20	ILE	2.3
1	B	22	LEU	2.3
1	B	168	GLU	2.3
1	B	186	TYR	2.3
1	A	64	VAL	2.3
1	C	86	VAL	2.3
1	C	14	ARG	2.2
1	A	226	CYS	2.2
1	B	189	ASN	2.2
1	C	153	GLU	2.2
1	A	158	LEU	2.2
1	C	179	ASP	2.2
1	C	203	ILE	2.2
1	A	6	SER	2.2
1	B	35	ALA	2.2
1	B	63	ALA	2.2
1	B	228	ALA	2.2
1	A	154	SER	2.2
1	A	5	LEU	2.2
1	B	34	ILE	2.2
1	B	192	HIS	2.2
1	B	18	VAL	2.2
1	B	162	ASP	2.2
1	A	170	ILE	2.1
1	B	67	TYR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	210	GLY	2.1
1	C	124	ALA	2.1
1	B	154	SER	2.1
1	A	89	ILE	2.1
1	A	102	ALA	2.1
1	B	29	ALA	2.1
1	B	58	TYR	2.1
1	C	243	TYR	2.1
1	A	76	VAL	2.1
1	A	107	GLN	2.1
1	B	236	VAL	2.1
1	B	51	PRO	2.1
1	C	123	GLY	2.1
1	A	137	ASN	2.1
1	B	61	LEU	2.1
1	B	215	PHE	2.1
1	A	54	ILE	2.1
1	C	73	ILE	2.1
1	A	18	VAL	2.1
1	B	152	VAL	2.1
1	C	137	ASN	2.0
1	C	39	GLY	2.0
1	B	53	THR	2.0
1	A	182	ALA	2.0
1	A	209	ALA	2.0
1	B	217	ALA	2.0
1	A	135	ILE	2.0
1	C	38	SER	2.0
1	B	156	THR	2.0
1	C	53	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	PO4	A	1258	5/5	0.55	0.23	65,66,67,68	0
2	PO4	A	1257	5/5	0.72	0.21	22,26,29,30	0
2	PO4	C	1257	5/5	0.74	0.19	22,24,30,31	0
2	PO4	B	1257	5/5	0.79	0.17	21,25,29,32	0
3	GOL	C	1258	6/6	0.85	0.19	25,36,38,39	0
3	GOL	C	1259	6/6	0.88	0.14	22,24,25,26	0

6.5 Other polymers [i](#)

There are no such residues in this entry.