



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 05:31 AM UTC

PDB ID : 7VWK / pdb_00007vwk
Title : The product template domain of AviM
Authors : Feng, Y.; Yang, X.; Zheng, J.
Deposited on : 2021-11-10
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	7% 90% 10%
1	B	260	8% 80% 12% 8%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 3743 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Polyketide synthase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	260	Total	C	N	O	S	0	1	0
			1928	1203	351	370	4			
1	B	240	Total	C	N	O	S	0	0	0
			1776	1110	319	343	4			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	27	Total	O	0	0
			27	27		
2	B	12	Total	O	0	0
			12	12		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Polyketide synthase



• Molecule 1: Polyketide synthase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.68Å 58.71Å 82.50Å 90.00° 96.28° 90.00°	Depositor
Resolution (Å)	50.00 – 2.00 50.00 – 2.00	Depositor EDS
% Data completeness (in resolution range)	97.9 (50.00-2.00) 98.0 (50.00-2.00)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.62 (at 2.00Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.202 , 0.253 0.212 , 0.257	Depositor DCC
R_{free} test set	1657 reflections (4.95%)	wwPDB-VP
Wilson B-factor (Å ²)	39.0	Xtriage
Anisotropy	0.064	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 36.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	3743	wwPDB-VP
Average B, all atoms (Å ²)	50.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.70% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.80	0/1970	0.63	0/2702
1	B	0.71	0/1812	0.60	0/2483
All	All	0.76	0/3782	0.61	0/5185

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1928	0	1911	19	0
1	B	1776	0	1755	26	0
2	A	27	0	0	0	0
2	B	12	0	0	0	0
All	All	3743	0	3666	41	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (41) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:934:TRP:HZ2	1:B:972:GLU:HG3	1.34	0.92

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:935:HIS:HE1	1:A:997:GLN:HE21	1.33	0.76
1:B:1143:VAL:HG13	1:B:1153:VAL:CG1	2.17	0.75
1:B:934:TRP:CZ2	1:B:972:GLU:HG3	2.22	0.73
1:B:935:HIS:HE1	1:B:997:GLN:HE21	1.37	0.73
1:B:915:ASP:OD1	1:B:1098:SER:HB2	1.89	0.73
1:A:935:HIS:CE1	1:A:997:GLN:HE21	2.07	0.72
1:B:1143:VAL:HG13	1:B:1153:VAL:HG11	1.71	0.71
1:A:1143:VAL:HG13	1:A:1153:VAL:HG11	1.74	0.69
1:B:915:ASP:CG	1:B:1098:SER:HB2	2.17	0.69
1:B:935:HIS:CE1	1:B:997:GLN:HE21	2.11	0.68
1:A:1143:VAL:HG13	1:A:1153:VAL:CG1	2.22	0.68
1:A:1110:ALA:HA	1:A:1143:VAL:HG11	1.76	0.68
1:A:983:THR:HG22	1:A:1125:HIS:HD2	1.59	0.67
1:B:977:ARG:NH2	1:B:1003:GLU:OE2	2.31	0.63
1:B:932:ARG:HE	1:B:1000:ARG:NH1	2.01	0.58
1:B:1110:ALA:HA	1:B:1143:VAL:HG11	1.85	0.58
1:A:983:THR:HG22	1:A:1125:HIS:CD2	2.41	0.56
1:A:931:LEU:HD21	1:B:926:VAL:HG11	1.89	0.54
1:B:956:ILE:HG12	1:B:989:LEU:CD2	2.37	0.54
1:B:1143:VAL:CG1	1:B:1153:VAL:HG11	2.38	0.52
1:A:1115:VAL:HG12	1:A:1116:GLY:N	2.25	0.52
1:A:922:THR:HG23	1:B:1018:PRO:HB2	1.93	0.51
1:B:938:LEU:HD23	1:B:990:THR:HG21	1.93	0.50
1:A:931:LEU:HD22	1:B:931:LEU:HD22	1.93	0.50
1:A:1143:VAL:CG1	1:A:1153:VAL:HG11	2.42	0.49
1:A:916:SER:O	1:A:917:HIS:HB2	2.14	0.47
1:B:1100:ALA:HB3	1:B:1101:PRO:HD3	1.97	0.45
1:A:951:LEU:O	1:A:952:ASN:C	2.60	0.45
1:A:1126:VAL:HG22	1:A:1171:LEU:HD23	1.99	0.45
1:B:1113:ALA:HB2	1:B:1143:VAL:HG12	2.00	0.44
1:A:926:VAL:HG11	1:B:931:LEU:HD21	2.01	0.42
1:B:1110:ALA:N	1:B:1111:PRO:HD2	2.34	0.42
1:A:1110:ALA:N	1:A:1111:PRO:HD2	2.34	0.42
1:A:1100:ALA:HB3	1:A:1101:PRO:HD3	2.01	0.42
1:B:1126:VAL:HG22	1:B:1171:LEU:HD23	2.01	0.42
1:B:1051:LEU:HB3	1:B:1081:SER:OG	2.20	0.41
1:B:956:ILE:HG12	1:B:989:LEU:HD23	2.03	0.41
1:B:943:ARG:HD2	1:B:943:ARG:HA	1.72	0.41
1:A:969:ALA:O	1:A:1000:ARG:HD3	2.20	0.41
1:B:1120:LEU:HD23	1:B:1120:LEU:HA	1.93	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	259/260 (100%)	257 (99%)	2 (1%)	0	100	100
1	B	234/260 (90%)	228 (97%)	6 (3%)	0	100	100
All	All	493/520 (95%)	485 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	201/200 (100%)	199 (99%)	2 (1%)	68	75
1	B	185/200 (92%)	183 (99%)	2 (1%)	65	73
All	All	386/400 (96%)	382 (99%)	4 (1%)	68	75

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	974	GLU
1	A	975	GLU
1	B	931	LEU
1	B	952	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	935	HIS
1	A	952	ASN
1	A	993	GLN
1	A	1125	HIS
1	B	935	HIS
1	B	993	GLN
1	B	1125	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	260/260 (100%)	0.40	17 (6%)	25 23	16, 41, 73, 120	1 (0%)
1	B	240/260 (92%)	0.85	21 (8%)	15 14	28, 54, 98, 134	0
All	All	500/520 (96%)	0.62	38 (7%)	20 18	16, 46, 88, 134	1 (0%)

All (38) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	1072	GLY	5.3
1	B	951	LEU	4.9
1	A	1058	LEU	4.2
1	A	1068	VAL	3.9
1	B	945	TYR	3.7
1	A	1069	PRO	3.6
1	A	1071	THR	3.2
1	B	1054	ALA	3.1
1	B	954	VAL	3.0
1	B	1080	LEU	3.0
1	B	943	ARG	2.8
1	B	1117	LEU	2.7
1	B	1051	LEU	2.6
1	B	1092	LEU	2.6
1	A	949	HIS	2.6
1	A	915	ASP	2.6
1	B	956	ILE	2.5
1	B	937	THR	2.5
1	B	1109	ILE	2.5
1	A	1084	LEU	2.4
1	A	942	SER	2.4
1	B	1053	PRO	2.4
1	A	948	SER	2.3
1	A	1117	LEU	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	1162	GLY	2.3
1	A	944	PRO	2.3
1	A	971	ALA	2.2
1	A	1162	GLY	2.2
1	B	1039	ALA	2.2
1	A	951	LEU	2.2
1	B	1116	GLY	2.2
1	B	1036	ALA	2.2
1	B	1040	ALA	2.2
1	B	1071	THR	2.2
1	A	973	GLY	2.1
1	B	1043	LEU	2.1
1	A	1140	THR	2.1
1	B	944	PRO	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.