



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 12:35 AM UTC

PDB ID : 3VX0 / pdb_00003vx0
Title : Crystal Structure of alpha-amylase from Aspergillus oryzae
Authors : Sugahara, M.
Deposited on : 2012-09-06
Resolution : 1.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	478	7% 81% 16% ..

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 4301 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Alpha-amylase A type-1/2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	475	Total	C	N	O	S	0	0	0
			3680	2333	594	735	18			

- Molecule 2 is 2-acetamido-2-deoxy-beta-D-glucopyranose (CCD ID: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			14	8	1	5		

- Molecule 3 is GADOLINIUM ATOM (CCD ID: GD) (formula: Gd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	6	Total	Gd	0	0
			6	6		

- Molecule 4 is CALCIUM ION (CCD ID: CA) (formula: Ca).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total 1	Ca 1	0	0

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	600	Total 600	O 600	0	0

i

- Molecule 1: Alpha-amylase A type-1/2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	48.52Å 65.62Å 130.24Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	18.97 – 1.50 18.97 – 1.50	Depositor EDS
% Data completeness (in resolution range)	(Not available) (18.97-1.50) 98.2 (18.97-1.50)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.74 (at 1.50Å)	Xtriage
Refinement program	CNS	Depositor
R, R_{free}	0.192 , 0.199 0.190 , 0.198	Depositor DCC
R_{free} test set	3336 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	13.6	Xtriage
Anisotropy	0.087	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 35.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4301	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GD, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.54	1/3775 (0.0%)	1.00	6/5156 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	216	TRP	NE1-CE2	11.65	1.50	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	11	ILE	N-CA-C	8.12	119.97	108.36
1	A	121	ASN	N-CA-C	6.71	118.60	111.28
1	A	213	LYS	N-CA-C	5.82	118.37	111.33
1	A	132	VAL	N-CA-C	5.51	116.37	108.93
1	A	1	ALA	O-C-N	-5.43	114.31	123.00
1	A	204	ARG	CD-NE-CZ	-5.07	117.31	124.40

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	1	ALA	Mainchain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3680	0	3473	68	2
2	A	14	0	13	0	0
3	A	6	0	0	0	0
4	A	1	0	0	0	0
5	A	600	0	0	9	2
All	All	4301	0	3486	68	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:433:GLN:NE2	1:A:442:THR:HB	1.63	1.11
1:A:433:GLN:HE22	1:A:442:THR:CB	1.75	0.98
1:A:433:GLN:HE22	1:A:442:THR:HB	0.82	0.97
1:A:92:GLU:OE1	5:A:1270:HOH:O	1.96	0.82
1:A:296:HIS:H	1:A:296:HIS:CD2	2.04	0.75
1:A:433:GLN:NE2	1:A:442:THR:CB	2.48	0.64
1:A:293:VAL:HG21	1:A:314:VAL:HG21	1.80	0.64
1:A:296:HIS:H	1:A:296:HIS:HD2	1.45	0.62
1:A:319:ILE:HD12	1:A:365:ALA:HB3	1.82	0.60
1:A:357:GLU:OE1	5:A:1605:HOH:O	2.15	0.60
1:A:357:GLU:CD	5:A:1605:HOH:O	2.45	0.60
1:A:397:ARG:HD3	1:A:397:ARG:C	2.28	0.59
1:A:351:GLY:O	1:A:352:TYR:HB2	2.02	0.58
1:A:212:GLN:H	1:A:212:GLN:HE21	1.53	0.57
1:A:280:LYS:NZ	5:A:1560:HOH:O	2.37	0.56
1:A:69:LEU:HD12	1:A:69:LEU:O	2.05	0.56
1:A:446:GLY:HA2	5:A:1412:HOH:O	2.07	0.55
1:A:361:LEU:C	1:A:361:LEU:HD13	2.32	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:LEU:HD12	1:A:69:LEU:C	2.34	0.53
1:A:6:TRP:CD1	1:A:226:TYR:HB3	2.45	0.52
1:A:1:ALA:N	1:A:201:ASP:O	2.43	0.52
1:A:63:THR:HB	1:A:64:PRO:HD2	1.92	0.52
1:A:252:TYR:N	1:A:253:PRO:CD	2.74	0.51
1:A:182:VAL:HG13	1:A:183:VAL:N	2.27	0.49
1:A:61:TRP:C	1:A:61:TRP:CD1	2.90	0.49
1:A:134:TYR:O	1:A:141:SER:HA	2.12	0.49
1:A:85:GLN:HA	1:A:123:MET:HE3	1.94	0.49
1:A:48:LYS:NZ	5:A:1496:HOH:O	2.44	0.49
1:A:319:ILE:CD1	1:A:365:ALA:HB3	2.41	0.48
1:A:384:ASN:HD21	1:A:396:MET:HB2	1.79	0.48
1:A:214:ASP:OD1	1:A:214:ASP:C	2.57	0.48
1:A:48:LYS:HA	1:A:48:LYS:HD2	1.67	0.48
1:A:308:ILE:HG13	1:A:308:ILE:O	2.14	0.47
1:A:67:ALA:HB3	1:A:86:ASP:HB3	1.97	0.47
1:A:432:GLN:NE2	5:A:1429:HOH:O	2.47	0.47
1:A:220:ASN:C	1:A:220:ASN:HD22	2.23	0.46
1:A:37:TYR:CE1	1:A:79:TYR:HA	2.51	0.46
1:A:134:TYR:CD1	1:A:143:GLN:HB2	2.51	0.45
1:A:82:TYR:OH	1:A:296:HIS:HE1	2.00	0.45
1:A:411:ASN:OD1	1:A:411:ASN:C	2.59	0.45
1:A:293:VAL:HB	1:A:318:ILE:HD11	1.99	0.45
1:A:474:ILE:O	1:A:475:CYS:HB2	2.16	0.45
1:A:166:LEU:HB3	1:A:173:LEU:HD12	1.99	0.45
1:A:212:GLN:H	1:A:212:GLN:NE2	2.15	0.45
1:A:276:ILE:O	1:A:280:LYS:HB2	2.17	0.44
1:A:343:ASN:OD1	1:A:344:ARG:HD2	2.18	0.43
1:A:134:TYR:CE1	1:A:143:GLN:HB2	2.54	0.43
1:A:433:GLN:NE2	1:A:442:THR:CG2	2.81	0.43
1:A:265:THR:HA	1:A:411:ASN:O	2.19	0.43
1:A:307:ASP:HB3	1:A:310:LEU:HD12	2.01	0.42
1:A:432:GLN:O	1:A:445:VAL:HG23	2.19	0.42
1:A:397:ARG:NH2	1:A:401:ASP:OD1	2.41	0.42
1:A:48:LYS:HE2	1:A:352:TYR:CG	2.53	0.42
1:A:182:VAL:CG1	1:A:183:VAL:N	2.83	0.42
1:A:55:MET:HG3	1:A:359:TYR:CE2	2.55	0.42
1:A:388:TYR:CG	1:A:389:LYS:N	2.87	0.42
1:A:137:PHE:HB2	1:A:140:PHE:O	2.20	0.41
1:A:236:PRO:O	1:A:240:CYS:HB2	2.20	0.41
1:A:280:LYS:HG3	5:A:1322:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:408:ILE:N	1:A:408:ILE:HD12	2.36	0.41
1:A:340:ASP:OD1	1:A:341:PRO:HA	2.20	0.41
1:A:272:LEU:O	1:A:276:ILE:HG13	2.21	0.41
1:A:293:VAL:HG12	1:A:318:ILE:HD11	2.03	0.41
1:A:375:LYS:NZ	5:A:1452:HOH:O	2.54	0.40
1:A:319:ILE:HD12	1:A:365:ALA:CB	2.48	0.40
1:A:321:ASN:ND2	1:A:322:ASP:H	2.19	0.40
1:A:63:THR:HB	1:A:64:PRO:CD	2.50	0.40
1:A:296:HIS:CD2	1:A:296:HIS:N	2.75	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:156:GLU:OE1	5:A:1614:HOH:O[3_655]	1.87	0.33
1:A:448:ASP:OD2	5:A:1605:HOH:O[4_565]	2.10	0.10

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	473/478 (99%)	458 (97%)	14 (3%)	1 (0%)	43 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	352	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	396/399 (99%)	385 (97%)	11 (3%)	38 11

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	TRP
1	A	212	GLN
1	A	232	LEU
1	A	280	LYS
1	A	283	CYS
1	A	296	HIS
1	A	308	ILE
1	A	375	LYS
1	A	389	LYS
1	A	467	GLU
1	A	473	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (13) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	9	GLN
1	A	143	GLN
1	A	212	GLN
1	A	296	HIS
1	A	321	ASN
1	A	331	GLN
1	A	333	GLN
1	A	339	ASN
1	A	370	ASN
1	A	384	ASN
1	A	432	GLN
1	A	433	GLN
1	A	450	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 8 ligands modelled in this entry, 7 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	A	1001	1	14,14,15	0.66	0	17,19,21	1.20	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	A	1001	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1001	NAG	C1-O5-C5	3.26	116.56	112.19
2	A	1001	NAG	C6-C5-C4	-2.11	107.83	113.02

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1001	NAG	O5-C5-C6-O6
2	A	1001	NAG	C4-C5-C6-O6

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	475/478 (99%)	0.29	33 (6%) 23 25	8, 15, 31, 47	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	150	CYS	4.3
1	A	449	GLY	3.7
1	A	474	ILE	3.5
1	A	283	CYS	3.4
1	A	227	CYS	3.3
1	A	475	CYS	3.2
1	A	467	GLU	3.2
1	A	375	LYS	3.1
1	A	433	GLN	3.0
1	A	440	CYS	2.9
1	A	445	VAL	2.9
1	A	471	GLY	2.9
1	A	1	ALA	2.7
1	A	470	ALA	2.7
1	A	417	ASP	2.7
1	A	447	SER	2.6
1	A	429	THR	2.5
1	A	443	VAL	2.5
1	A	389	LYS	2.4
1	A	152	ILE	2.3
1	A	469	LEU	2.3
1	A	446	GLY	2.3
1	A	151	PHE	2.3
1	A	421	LEU	2.3
1	A	448	ASP	2.2
1	A	391	ASP	2.2
1	A	182	VAL	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	352	TYR	2.1
1	A	308	ILE	2.1
1	A	473	LYS	2.1
1	A	149	PHE	2.0
1	A	155	TYR	2.0
1	A	428	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	NAG	A	1001	14/15	0.63	0.28	54,70,79,79	0
3	GD	A	1003	1/1	0.81	0.14	35,35,35,35	1
3	GD	A	1004	1/1	0.94	0.06	23,23,23,23	1
3	GD	A	1007	1/1	0.94	0.05	14,14,14,14	1
3	GD	A	1002	1/1	0.96	0.05	15,15,15,15	1
3	GD	A	1005	1/1	0.97	0.05	18,18,18,18	1
3	GD	A	1006	1/1	0.98	0.04	16,16,16,16	1
4	CA	A	1008	1/1	0.99	0.02	9,9,9,9	0

6.5 Other polymers [i](#)

There are no such residues in this entry.