



wwPDB EM Validation Summary Report ⓘ

Mar 8, 2026 – 07:59 AM UTC

PDB ID : 7VYE / pdb_00007vye
EMDB ID : EMD-32202
Title : Membrane arm of deactive state CI from Q10-NADH dataset
Authors : Gu, J.K.; Yang, M.J.
Deposited on : 2021-11-14
Resolution : 3.10 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:



EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49

1 Overall quality at a glance

The following experimental techniques were used to determine the structure:
ELECTRON MICROSCOPY

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.













Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14724 (2.60 - 3.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	Q	40	
2	S	70	
3	U	83	
4	V	140	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
5	W	113	
6	X	88	
7	Y	70	
8	Z	84	
9	a	140	
10	b	126	
11	c	156	
12	d	175	
13	e	107	
14	f	42	
15	g	121	
16	h	105	
17	i	347	
18	j	113	
19	k	98	
20	l	603	
21	m	175	
22	n	56	
23	o	128	
24	p	178	
25	r	459	
26	s	318	
27	u	171	
28	v	126	
29	w	320	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
30	PEE	r	501	-	-	X	-
31	CDL	l	701	-	-	X	-

2 Entry composition

There are 35 unique types of molecules in this entry. The entry contains 38846 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Q	40	Total	C	N	O	S	0	0
			333	217	56	59	1		

- Molecule 2 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	S	70	Total	C	N	O	S	0	0
			567	364	104	94	5		

- Molecule 3 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	U	83	Total	C	N	O	S	0	0
			643	417	110	115	1		

- Molecule 4 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 11.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	V	140	Total	C	N	O	S	0	0
			1015	648	171	190	6		

- Molecule 5 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	W	113	Total	C	N	O	S	0	0
			945	612	160	165	8		

- Molecule 6 is a protein called Acyl carrier protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	X	88	Total	C	N	O	S	0	0
			699	449	103	142	5		

- Molecule 7 is a protein called NADH-ubiquinone oxidoreductase AGGG subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	Y	70	Total	C	N	O	S	0	0
			600	393	98	108	1		

- Molecule 8 is a protein called NADH-ubiquinone oxidoreductase B12 subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	Z	84	Total	C	N	O	S	0	0
			674	437	116	120	1		

- Molecule 9 is a protein called NADH-ubiquinone oxidoreductase SGDH subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	a	140	Total	C	N	O	S	0	0
			1165	762	199	201	3		

- Molecule 10 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	b	103	Total	C	N	O	S	0	0
			879	573	158	147	1		

- Molecule 11 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	c	156	Total	C	N	O	S	0	0
			1315	853	213	241	8		

- Molecule 12 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	d	175	Total	C	N	O	S	0	0
			1461	916	265	272	8		

- Molecule 13 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit

11, mitochondrial.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	e	107	Total	C	N	O	S	0	0
			890	568	145	173	4		

- Molecule 14 is a protein called NADH-ubiquinone oxidoreductase KFYI subunit.

Mol	Chain	Residues	Atoms				AltConf	Trace
14	f	42	Total	C	N	O	0	0
			342	225	58	59		

- Molecule 15 is a protein called NADH dehydrogenase [ubiquinone] 1 subunit C2.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	g	121	Total	C	N	O	S	0	0
			1000	650	173	171	6		

- Molecule 16 is a protein called NADH dehydrogenase [ubiquinone] iron-sulfur protein 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
16	h	105	Total	C	N	O	S	0	0
			867	550	161	150	6		

- Molecule 17 is a protein called NADH-ubiquinone oxidoreductase chain 2.

Mol	Chain	Residues	Atoms					AltConf	Trace
17	i	347	Total	C	N	O	S	0	0
			2710	1782	420	462	46		

- Molecule 18 is a protein called NADH-ubiquinone oxidoreductase chain 3.

Mol	Chain	Residues	Atoms					AltConf	Trace
18	j	99	Total	C	N	O	S	0	0
			800	545	118	132	5		

- Molecule 19 is a protein called NADH-ubiquinone oxidoreductase chain 4L.

Mol	Chain	Residues	Atoms					AltConf	Trace
19	k	98	Total	C	N	O	S	0	0
			748	493	113	128	14		

- Molecule 20 is a protein called NADH-ubiquinone oxidoreductase chain 5.

Mol	Chain	Residues	Atoms					AltConf	Trace
20	l	603	Total	C	N	O	S	0	0
			4785	3173	741	820	51		

- Molecule 21 is a protein called NADH-ubiquinone oxidoreductase chain 6.

Mol	Chain	Residues	Atoms					AltConf	Trace
21	m	129	Total	C	N	O	S	0	0
			951	637	138	168	8		

- Molecule 22 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
22	n	56	Total	C	N	O	S	0	0
			479	311	88	79	1		

- Molecule 23 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
23	o	128	Total	C	N	O	S	0	0
			1062	691	182	189			

- Molecule 24 is a protein called NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9.

Mol	Chain	Residues	Atoms					AltConf	Trace
24	p	178	Total	C	N	O	S	0	0
			1534	982	279	265	8		

- Molecule 25 is a protein called NADH-ubiquinone oxidoreductase chain 4.

Mol	Chain	Residues	Atoms					AltConf	Trace
25	r	459	Total	C	N	O	S	0	0
			3631	2412	572	609	38		

- Molecule 26 is a protein called NADH-ubiquinone oxidoreductase chain 1.

Mol	Chain	Residues	Atoms					AltConf	Trace
26	s	303	Total	C	N	O	S	0	0
			2394	1607	369	397	21		

- Molecule 27 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex sub-

unit 8.

Mol	Chain	Residues	Atoms					AltConf	Trace
27	u	171	Total	C	N	O	S	0	0
			1398	887	250	251	10		

- Molecule 28 is a protein called NADH-ubiquinone oxidoreductase B18 subunit.

Mol	Chain	Residues	Atoms					AltConf	Trace
28	v	124	Total	C	N	O	S	0	0
			1012	633	188	182	9		


There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
v	1	MYR	-	acetylation	UNP F1SCH1

- Molecule 29 is a protein called NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial.


Mol	Chain	Residues	Atoms					AltConf	Trace
29	w	320	Total	C	N	O	S	0	0
			2583	1646	437	491	9		

- Molecule 30 is 1,2-dioleoyl-sn-glycero-3-phosphoethanolamine (CCD ID: PEE) (formula: $C_{41}H_{78}NO_8P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					AltConf
30	U	1	Total	C	N	O	P	0
			51	41	1	8	1	
30	V	1	Total	C	N	O	P	0
			40	30	1	8	1	
30	b	1	Total	C	N	O	P	0
			46	36	1	8	1	
30	i	1	Total	C	N	O	P	0
			47	37	1	8	1	
30	l	1	Total	C	N	O	P	0
			46	36	1	8	1	
30	r	1	Total	C	N	O	P	0
			51	41	1	8	1	
30	s	1	Total	C	N	O	P	0
			41	31	1	8	1	
30	s	1	Total	C	N	O	P	0
			51	41	1	8	1	

- Molecule 31 is CARDIOLIPIN (CCD ID: CDL) (formula: $C_{81}H_{156}O_{17}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				AltConf
31	V	1	Total	C	O	P	0
			94	75	17	2	
31	a	1	Total	C	O	P	0
			91	72	17	2	
31	g	1	Total	C	O	P	0
			97	78	17	2	

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				AltConf
31	i	1	Total	C	O	P	0
			66	47	17	2	
31	l	1	Total	C	O	P	0
			99	80	17	2	
31	l	1	Total	C	O	P	0
			100	81	17	2	
31	o	1	Total	C	O	P	0
			68	49	17	2	
31	u	1	Total	C	O	P	0
			78	59	17	2	

- Molecule 32 is S-[2-({N-[(2R)-2-hydroxy-3,3-dimethyl-4-(phosphonooxy)butanoyl]-beta-alanyl}amino)ethyl] dodecanethioate (CCD ID: 8Q1) (formula: C₂₃H₄₅N₂O₈PS) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms						AltConf
32	X	1	Total	C	N	O	P	S	0
			35	23	2	8	1	1	

- Molecule 33 is (9R,11S)-9-({[(1S)-1-HYDROXYHEXADECYL]OXY}METHYL)-2,2-DIMETHYL-5,7,10-TRIOXA-2LAMBDA 5 -AZA-6LAMBDA 5 -PHOSPHAOCTACOSANE-6,6,11-TRIOL (CCD ID: PLX) (formula: C₄₂H₈₉NO₈P) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
33	a	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	g	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	j	1	Total	C	N	O	P	0
			52	42	1	8	1	
33	r	1	Total	C	N	O	P	0
			52	42	1	8	1	

- Molecule 34 is Coenzyme Q10, (2Z,6E,10Z,14E,18E,22E,26Z)-isomer (CCD ID: UQ) (formula: C₅₉H₉₀O₄) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
34	s	1	Total	C	O	0
			28	24	4	

- Molecule 35 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					AltConf
35	w	1	Total	C	N	O	P	0
			27	10	5	10	2	


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NADH dehydrogenase [ubiquinone] iron-sulfur protein 2




- Molecule 2: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 1




- Molecule 3: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 3



- Molecule 4: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 11




- Molecule 5: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 13







- Molecule 6: Acyl carrier protein





- Molecule 7: NADH-ubiquinone oxidoreductase AGGG subunit




- Molecule 8: NADH-ubiquinone oxidoreductase B12 subunit




- Molecule 9: NADH-ubiquinone oxidoreductase SGD subunit




- Molecule 10: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 6





- Molecule 11: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 8, mitochondrial









- Molecule 12: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10




- Molecule 13: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 11, mitochondrial





- Molecule 14: NADH-ubiquinone oxidoreductase KFYI subunit




- Molecule 15: NADH dehydrogenase [ubiquinone] 1 subunit C2




- Molecule 16: NADH dehydrogenase [ubiquinone] iron-sulfur protein 5




- Molecule 17: NADH-ubiquinone oxidoreductase chain 2




- Molecule 18: NADH-ubiquinone oxidoreductase chain 3




- Molecule 19: NADH-ubiquinone oxidoreductase chain 4L




- Molecule 20: NADH-ubiquinone oxidoreductase chain 5








- Molecule 21: NADH-ubiquinone oxidoreductase chain 6





- Molecule 22: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 1



- Molecule 23: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 4




- Molecule 24: NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 9




- Molecule 25: NADH-ubiquinone oxidoreductase chain 4








• Molecule 26: NADH-ubiquinone oxidoreductase chain 1



• Molecule 27: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 8




• Molecule 28: NADH-ubiquinone oxidoreductase B18 subunit



• Molecule 29: NADH dehydrogenase [ubiquinone] 1 alpha subcomplex subunit 10, mitochondrial





4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	104881	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1300	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K3 (6k x 4k)	Depositor
Maximum map value	0.157	Depositor
Minimum map value	-0.086	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.005	Depositor
Recommended contour level	0.0306	Depositor
Map size (Å)	333.002, 333.002, 333.002	wwPDB
Map dimensions	310, 310, 310	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	1.0742, 1.0742, 1.0742	Depositor

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PEE, 8Q1, UQ, CDL, PLX, ADP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	Q	0.15	0/350	0.33	0/483
2	S	0.21	0/582	0.42	0/783
3	U	0.16	0/664	0.31	0/912
4	V	0.22	0/1036	0.53	2/1404 (0.1%)
5	W	0.23	0/969	0.41	0/1307
6	X	0.17	0/711	0.37	0/963
7	Y	0.15	0/626	0.34	0/857
8	Z	0.16	0/695	0.35	0/939
9	a	0.23	0/1199	0.62	2/1623 (0.1%)
10	b	0.21	0/906	0.54	1/1232 (0.1%)
11	c	0.23	0/1371	0.52	2/1875 (0.1%)
12	d	0.16	0/1494	0.29	0/2015
13	e	0.21	0/916	0.54	2/1246 (0.2%)
14	f	0.15	0/350	0.30	0/473
15	g	0.18	0/1031	0.32	0/1394
16	h	0.22	0/889	0.51	0/1190
17	i	0.21	0/2773	0.39	0/3768
18	j	0.28	0/819	0.48	0/1117
19	k	0.22	0/759	0.39	0/1029
20	l	0.20	0/4914	0.39	0/6683
21	m	0.21	0/973	0.41	0/1320
22	n	0.22	0/491	0.59	1/663 (0.2%)
23	o	0.17	0/1092	0.33	0/1481
24	p	0.18	0/1590	0.38	1/2155 (0.0%)
25	r	0.23	0/3723	0.42	0/5078
26	s	0.25	0/2464	0.53	2/3369 (0.1%)
27	u	0.23	0/1436	0.52	1/1938 (0.1%)
28	v	0.17	0/1036	0.48	0/1393
29	w	0.20	0/2643	0.45	1/3580 (0.0%)
All	All	0.21	0/38502	0.44	15/52270 (0.0%)

There are no bond length outliers.

The worst 5 of 15 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
9	a	94	GLY	N-CA-C	10.19	123.28	111.36
9	a	92	PHE	N-CA-C	8.66	120.80	111.36
11	c	75	TYR	CA-C-N	8.26	128.19	119.85
11	c	75	TYR	C-N-CA	8.26	128.19	119.85
26	s	87	VAL	CA-C-N	8.07	127.65	118.85

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Q	333	0	301	7	0
2	S	567	0	565	7	0
3	U	643	0	642	9	0
4	V	1015	0	1016	25	0
5	W	945	0	931	27	0
6	X	699	0	674	19	0
7	Y	600	0	539	12	0
8	Z	674	0	643	12	0
9	a	1165	0	1174	38	0
10	b	879	0	899	34	0
11	c	1315	0	1208	35	0
12	d	1461	0	1429	42	0
13	e	890	0	837	19	0
14	f	342	0	341	6	0
15	g	1000	0	994	14	0
16	h	867	0	871	16	0
17	i	2710	0	2874	78	0
18	j	800	0	855	38	0
19	k	748	0	799	23	0
20	l	4785	0	4933	117	0
21	m	951	0	962	38	0
22	n	479	0	486	14	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
23	o	1062	0	1072	9	0
24	p	1534	0	1470	30	0
25	r	3631	0	3839	93	0
26	s	2394	0	2508	76	0
27	u	1398	0	1378	35	0
28	v	1012	0	947	21	0
29	w	2583	0	2539	63	0
30	U	51	0	82	6	0
30	V	40	0	54	6	0
30	b	46	0	69	18	0
30	i	47	0	71	10	0
30	l	46	0	69	8	0
30	r	51	0	82	21	0
30	s	92	0	141	10	0
31	V	94	0	138	19	0
31	a	91	0	132	16	0
31	g	97	0	147	8	0
31	i	66	0	76	16	0
31	l	199	0	307	46	0
31	o	68	0	83	14	0
31	u	78	0	103	13	0
32	X	35	0	0	2	0
33	a	52	0	88	9	0
33	g	52	0	88	1	0
33	j	52	0	88	2	0
33	r	52	0	88	7	0
34	s	28	0	31	5	0
35	w	27	0	11	4	0
All	All	38846	0	39674	928	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

The worst 5 of 928 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:a:96:ALA:HB2	12:d:61:TYR:CE2	1.52	1.41
31:V:202:CDL:H601	31:V:202:CDL:H732	1.24	1.18
30:i:402:PEE:H36	30:i:402:PEE:H28	1.25	1.17
25:r:201:MET:CE	30:r:501:PEE:H78	1.75	1.15
31:o:201:CDL:H112	31:o:201:CDL:H342	1.26	1.13

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	Q	38/40 (95%)	37 (97%)	1 (3%)	0	100	100
2	S	68/70 (97%)	63 (93%)	5 (7%)	0	100	100
3	U	81/83 (98%)	79 (98%)	2 (2%)	0	100	100
4	V	138/140 (99%)	132 (96%)	5 (4%)	1 (1%)	18	49
5	W	111/113 (98%)	109 (98%)	2 (2%)	0	100	100
6	X	86/88 (98%)	83 (96%)	3 (4%)	0	100	100
7	Y	68/70 (97%)	64 (94%)	4 (6%)	0	100	100
8	Z	82/84 (98%)	78 (95%)	4 (5%)	0	100	100
9	a	138/140 (99%)	136 (99%)	1 (1%)	1 (1%)	18	49
10	b	99/126 (79%)	95 (96%)	4 (4%)	0	100	100
11	c	154/156 (99%)	145 (94%)	8 (5%)	1 (1%)	21	52
12	d	173/175 (99%)	172 (99%)	1 (1%)	0	100	100
13	e	105/107 (98%)	102 (97%)	3 (3%)	0	100	100
14	f	40/42 (95%)	39 (98%)	1 (2%)	0	100	100
15	g	119/121 (98%)	114 (96%)	5 (4%)	0	100	100
16	h	103/105 (98%)	98 (95%)	5 (5%)	0	100	100
17	i	345/347 (99%)	328 (95%)	15 (4%)	2 (1%)	21	52
18	j	95/113 (84%)	89 (94%)	6 (6%)	0	100	100
19	k	96/98 (98%)	87 (91%)	9 (9%)	0	100	100
20	l	601/603 (100%)	575 (96%)	26 (4%)	0	100	100
21	m	125/175 (71%)	112 (90%)	13 (10%)	0	100	100
22	n	54/56 (96%)	54 (100%)	0	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
23	o	126/128 (98%)	120 (95%)	6 (5%)	0	100	100
24	p	176/178 (99%)	163 (93%)	13 (7%)	0	100	100
25	r	457/459 (100%)	441 (96%)	16 (4%)	0	100	100
26	s	299/318 (94%)	286 (96%)	12 (4%)	1 (0%)	36	67
27	u	169/171 (99%)	164 (97%)	5 (3%)	0	100	100
28	v	122/126 (97%)	115 (94%)	7 (6%)	0	100	100
29	w	318/320 (99%)	303 (95%)	15 (5%)	0	100	100
All	All	4586/4752 (96%)	4383 (96%)	197 (4%)	6 (0%)	49	78

5 of 6 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
17	i	93	VAL
11	c	81	ARG
9	a	95	GLU
17	i	92	PRO
4	V	46	PRO

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	Q	34/34 (100%)	34 (100%)	0	100	100
2	S	58/58 (100%)	58 (100%)	0	100	100
3	U	69/69 (100%)	69 (100%)	0	100	100
4	V	100/101 (99%)	100 (100%)	0	100	100
5	W	98/99 (99%)	98 (100%)	0	100	100
6	X	77/81 (95%)	77 (100%)	0	100	100
7	Y	63/63 (100%)	63 (100%)	0	100	100
8	Z	65/65 (100%)	65 (100%)	0	100	100
9	a	122/122 (100%)	120 (98%)	2 (2%)	55	75

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
10	b	98/119 (82%)	98 (100%)	0	100	100
11	c	141/141 (100%)	140 (99%)	1 (1%)	76	82
12	d	155/155 (100%)	155 (100%)	0	100	100
13	e	99/99 (100%)	99 (100%)	0	100	100
14	f	35/38 (92%)	35 (100%)	0	100	100
15	g	108/108 (100%)	108 (100%)	0	100	100
16	h	93/93 (100%)	91 (98%)	2 (2%)	45	71
17	i	311/311 (100%)	311 (100%)	0	100	100
18	j	88/99 (89%)	87 (99%)	1 (1%)	65	78
19	k	85/85 (100%)	85 (100%)	0	100	100
20	l	537/537 (100%)	537 (100%)	0	100	100
21	m	99/141 (70%)	99 (100%)	0	100	100
22	n	53/53 (100%)	53 (100%)	0	100	100
23	o	113/113 (100%)	113 (100%)	0	100	100
24	p	159/159 (100%)	159 (100%)	0	100	100
25	r	410/410 (100%)	410 (100%)	0	100	100
26	s	263/275 (96%)	262 (100%)	1 (0%)	84	86
27	u	153/153 (100%)	151 (99%)	2 (1%)	61	77
28	v	101/112 (90%)	101 (100%)	0	100	100
29	w	281/283 (99%)	279 (99%)	2 (1%)	76	82
All	All	4068/4176 (97%)	4057 (100%)	11 (0%)	84	87

5 of 11 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
27	u	88	CYS
27	u	100	CYS
29	w	246	LEU
29	w	214	ILE
16	h	102	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 42 such sidechains are listed below:

Mol	Chain	Res	Type
25	r	192	ASN
27	u	163	HIS
25	r	304	GLN
26	s	317	GLN
28	v	85	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

23 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
31	CDL	g	202	-	96,96,99	1.11	8 (8%)	102,108,111	0.89	4 (3%)
32	8Q1	X	201	-	32,34,34	1.62	6 (18%)	39,43,43	1.61	6 (15%)
33	PLX	a	202	-	51,51,51	0.60	0	53,59,59	0.70	0
33	PLX	r	502	-	51,51,51	1.20	5 (9%)	53,59,59	0.63	1 (1%)
34	UQ	s	403	-	28,28,63	3.30	7 (25%)	36,37,79	2.77	11 (30%)
31	CDL	a	201	-	90,90,99	0.97	4 (4%)	96,102,111	1.07	6 (6%)
31	CDL	u	201	-	77,77,99	1.02	4 (5%)	83,89,111	1.16	7 (8%)
33	PLX	j	201	-	51,51,51	1.21	4 (7%)	53,59,59	0.61	1 (1%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
31	CDL	o	201	-	67,67,99	1.11	4 (5%)	73,79,111	1.20	6 (8%)
31	CDL	i	401	-	65,65,99	1.14	4 (6%)	71,77,111	1.21	5 (7%)
30	PEE	i	402	-	46,46,50	1.22	6 (13%)	49,51,55	1.02	3 (6%)
31	CDL	l	701	-	98,98,99	0.92	4 (4%)	104,110,111	1.17	9 (8%)
30	PEE	s	401	-	40,40,50	1.16	5 (12%)	43,45,55	1.05	2 (4%)
33	PLX	g	201	-	51,51,51	1.18	3 (5%)	53,59,59	0.67	1 (1%)
31	CDL	l	702	-	99,99,99	1.11	8 (8%)	105,111,111	0.84	4 (3%)
30	PEE	s	402	-	50,50,50	1.18	6 (12%)	53,55,55	0.98	2 (3%)
30	PEE	U	101	-	50,50,50	1.17	6 (12%)	53,55,55	0.96	2 (3%)
31	CDL	V	202	-	93,93,99	0.95	4 (4%)	99,105,111	1.08	5 (5%)
30	PEE	l	703	-	45,45,50	1.24	6 (13%)	48,50,55	1.02	2 (4%)
30	PEE	b	201	-	45,45,50	1.24	6 (13%)	48,50,55	1.02	2 (4%)
35	ADP	w	401	-	28,29,29	3.16	9 (32%)	43,45,45	1.91	9 (20%)
30	PEE	r	501	-	50,50,50	1.17	6 (12%)	53,55,55	1.02	2 (3%)
30	PEE	V	201	-	39,39,50	1.33	6 (15%)	42,44,55	1.10	3 (7%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
31	CDL	g	202	-	-	63/107/107/110	-
32	8Q1	X	201	-	-	17/41/41/41	-
33	PLX	a	202	-	-	15/55/55/55	-
33	PLX	r	502	-	-	38/55/55/55	-
34	UQ	s	403	-	-	9/21/45/87	0/1/1/1
31	CDL	a	201	-	-	26/101/101/110	-
31	CDL	u	201	-	-	28/88/88/110	-
33	PLX	j	201	-	-	24/55/55/55	-
31	CDL	o	201	-	-	33/78/78/110	-
31	CDL	i	401	-	-	30/76/76/110	-
30	PEE	i	402	-	-	22/50/50/54	-
31	CDL	l	701	-	-	39/109/109/110	-
30	PEE	s	401	-	-	18/44/44/54	-
33	PLX	g	201	-	-	23/55/55/55	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
31	CDL	l	702	-	-	65/110/110/110	-
30	PEE	s	402	-	-	22/54/54/54	-
30	PEE	U	101	-	-	24/54/54/54	-
31	CDL	V	202	-	-	37/104/104/110	-
30	PEE	l	703	-	-	20/49/49/54	-
30	PEE	b	201	-	-	26/49/49/54	-
35	ADP	w	401	-	-	3/16/32/32	0/3/3/3
30	PEE	r	501	-	-	21/54/54/54	-
30	PEE	V	201	-	-	21/43/43/54	-

The worst 5 of 121 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
34	s	403	UQ	C13-C14	9.59	1.55	1.33
34	s	403	UQ	C8-C9	9.12	1.54	1.33
35	w	401	ADP	C3'-C4'	-8.95	1.30	1.53
34	s	403	UQ	C18-C19	7.94	1.56	1.32
35	w	401	ADP	O4'-C4'	7.65	1.62	1.45

The worst 5 of 93 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
34	s	403	UQ	C7-C8-C9	-9.66	110.19	126.83
34	s	403	UQ	C12-C13-C14	-6.19	113.45	127.62
32	X	201	8Q1	C6-C1-S44	5.98	120.53	113.40
35	w	401	ADP	C5-C4-N3	-5.45	119.21	126.72
31	l	701	CDL	OA6-CA5-C11	4.86	121.99	111.48

There are no chirality outliers.

5 of 624 torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
30	U	101	PEE	C1-O3P-P-O1P
30	U	101	PEE	C4-O4P-P-O3P
30	U	101	PEE	C4-O4P-P-O2P
30	U	101	PEE	C4-O4P-P-O1P
30	V	201	PEE	C11-C10-O2-C2


There are no ring outliers.


23 monomers are involved in 226 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
31	g	202	CDL	8	0
32	X	201	8Q1	2	0
33	a	202	PLX	9	0
33	r	502	PLX	7	0
34	s	403	UQ	5	0
31	a	201	CDL	16	0
31	u	201	CDL	13	0
33	j	201	PLX	2	0
31	o	201	CDL	14	0
31	i	401	CDL	16	0
30	i	402	PEE	10	0
31	l	701	CDL	39	0
30	s	401	PEE	8	0
33	g	201	PLX	1	0
31	l	702	CDL	7	0
30	s	402	PEE	2	0
30	U	101	PEE	6	0
31	V	202	CDL	19	0
30	l	703	PEE	8	0
30	b	201	PEE	18	0
35	w	401	ADP	4	0
30	r	501	PEE	21	0
30	V	201	PEE	6	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
























5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-32202. These allow visual inspection of the internal detail of the map and identification of artifacts.


No raw map or half-maps were deposited for this entry and therefore no images, graphs, etc. pertaining to the raw map can be shown.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X



Y




Z

The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 155



Y Index: 155




Z Index: 155


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 204



Y Index: 131




Z Index: 119


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) [i](#)


6.4.1 Primary map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.

6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.0306. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.


6.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 177 nm³; this corresponds to an approximate mass of 160 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ



*Reported resolution corresponds to spatial frequency of 0.323 Å⁻¹


8 Fourier-Shell correlation

This section was not generated. No FSC curve or half-maps provided.


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-32202 and PDB model 7VYE. Per-residue inclusion information can be found in section [3](#) on page [14](#).


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.0306 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.0306).

9.4 Atom inclusion [i](#)



At the recommended contour level, 80% of all backbone atoms, 74% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.0306) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	 0.7420	 0.5700
Q	 0.5950	 0.5550
S	 0.8170	 0.5770
U	 0.7120	 0.5550
V	 0.4550	 0.5080
W	 0.7870	 0.5860
X	 0.7270	 0.5500
Y	 0.6100	 0.5110
Z	 0.5610	 0.4790
a	 0.7680	 0.5850
b	 0.6630	 0.5280
c	 0.7610	 0.5690
d	 0.7390	 0.5590
e	 0.7010	 0.5560
f	 0.6650	 0.5450
g	 0.7750	 0.5850
h	 0.7670	 0.5780
i	 0.8250	 0.6030
j	 0.6210	 0.5370
k	 0.6970	 0.5690
l	 0.7830	 0.5870
m	 0.6940	 0.5530
n	 0.6490	 0.5300
o	 0.7070	 0.5660
p	 0.7810	 0.5720
r	 0.8480	 0.6060
s	 0.7630	 0.5800
u	 0.7450	 0.5760
v	 0.6430	 0.5080
w	 0.7230	 0.5640

