



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 5, 2026 – 08:42 AM UTC

PDB ID : 2VZO / pdb_00002vzo
Title : Crystal structure of Amycolatopsis orientalis exo-chitosanase CsxA
Authors : Lammerts van Bueren, A.; Ghinet, M.G.; Gregg, K.; Fleury, A.; Brzezinski, R.; Boraston, A.B.
Deposited on : 2008-08-05
Resolution : 2.24 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3416 (2.26-2.22)
Clashscore	190562	3556 (2.26-2.22)
Ramachandran outliers	187476	3500 (2.26-2.22)
Sidechain outliers	187428	3501 (2.26-2.22)
RSRZ outliers	180081	3415 (2.26-2.22)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	1032	
1	B	1032	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	CD	A	1903	-	-	-	X

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13644 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXO-BETA-D-GLUCOSAMINIDASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	851	Total	C	N	O	S	9	0	1
			6510	4090	1126	1277	17			
1	B	849	Total	C	N	O	S	0	0	1
			6488	4077	1123	1271	17			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	469	GLU	ASP	engineered mutation	UNP Q56F26
A	750	ASN	TRP	conflict	UNP Q56F26
B	469	GLU	ASP	engineered mutation	UNP Q56F26
B	750	ASN	TRP	conflict	UNP Q56F26

- Molecule 2 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	7	Total	Cd	0	0
			7	7		
2	B	6	Total	Cd	0	0
			6	6		

- Molecule 3 is ACETATE ION (CCD ID: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			6	3	3		
4	A	1	Total	C	O	0	0
			6	3	3		


Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			6	3	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	301	Total	O	0	0
			301	301		
5	B	306	Total	O	0	0
			306	306		



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	86.40Å 122.03Å 91.60Å 90.00° 90.52° 90.00°	Depositor
Resolution (Å)	20.00 – 2.24 20.00 – 2.24	Depositor EDS
% Data completeness (in resolution range)	99.2 (20.00-2.24) 99.0 (20.00-2.24)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	5.24 (at 2.24Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.175 , 0.223 0.175 , 0.222	Depositor DCC
R_{free} test set	4531 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	28.9	Xtriage
Anisotropy	0.267	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 32.8	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.022 for h,-k,-l	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	13644	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.73% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, ACT, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.46	9/6673 (0.1%)	1.11	22/9105 (0.2%)
1	B	1.01	3/6650 (0.0%)	1.05	16/9074 (0.2%)
All	All	1.25	12/13323 (0.1%)	1.08	38/18179 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	2	3
1	B	1	3
All	All	3	6

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	431	GLU	CA-CB	86.30	2.99	1.53
1	A	468	SER	CA-C	7.96	1.56	1.52
1	A	430	GLU	CB-CG	7.67	1.75	1.52
1	B	300	VAL	CA-CB	7.31	1.62	1.54
1	A	739	VAL	CA-CB	6.11	1.61	1.54

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	431	GLU	N-CA-CB	-28.23	62.79	110.49
1	A	431	GLU	CB-CA-C	-21.38	67.88	110.42
1	B	99	LYS	N-CA-C	-14.26	96.36	114.04
1	A	99	LYS	N-CA-C	-12.16	98.58	113.41

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	597	TYR	N-CA-C	11.46	135.21	110.80

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
1	A	469	GLU	CA
1	A	597	TYR	CA
1	B	597	TYR	CA

5 of 6 planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	468	SER	Peptide
1	A	596	ARG	Peptide
1	A	98	GLY	Peptide
1	B	468	SER	Peptide
1	B	98	GLY	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6510	0	6287	80	0
1	B	6488	0	6263	73	0
2	A	7	0	0	0	0
2	B	6	0	0	0	0
3	A	4	0	3	0	0
3	B	4	0	3	0	0
4	A	12	0	16	0	0
4	B	6	0	8	0	0
5	A	301	0	0	9	1
5	B	306	0	0	11	1
All	All	13644	0	12580	153	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 153 close contacts within the same asymmetric unit are listed below, sorted by

their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:109:MET:CE	1:B:368:PHE:HE1	1.50	1.22
1:B:109:MET:CE	1:B:368:PHE:CE1	2.29	1.14
1:A:608:ARG:HG3	1:A:889:TRP:CZ3	1.96	0.99
1:A:773:THR:HG23	1:A:788:ALA:HB3	1.45	0.98
1:B:895:THR:HG22	5:B:2306:HOH:O	1.63	0.97

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:2044:HOH:O	5:B:2099:HOH:O[1_655]	2.15	0.05

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	849/1032 (82%)	823 (97%)	22 (3%)	4 (0%)	24	23
1	B	845/1032 (82%)	824 (98%)	18 (2%)	3 (0%)	30	30
All	All	1694/2064 (82%)	1647 (97%)	40 (2%)	7 (0%)	30	30

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	469	GLU
1	A	597	TYR
1	B	469	GLU
1	B	597	TYR
1	A	431	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	698/834 (84%)	675 (97%)	23 (3%)	33	39
1	B	695/834 (83%)	668 (96%)	27 (4%)	28	33
All	All	1393/1668 (84%)	1343 (96%)	50 (4%)	31	36

5 of 50 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	247	LEU
1	B	597	TYR
1	B	895	THR
1	B	299	ASN
1	B	356	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 33 such sidechains are listed below:

Mol	Chain	Res	Type
1	B	750	ASN
1	B	786	GLN
1	B	894	GLN
1	A	796	ASN
1	A	786	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 13 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	GOL	A	1907	-	5,5,5	0.38	0	5,5,5	1.11	1 (20%)
3	ACT	B	1906	-	3,3,3	1.27	0	3,3,3	1.41	0
4	GOL	A	1908	-	5,5,5	0.42	0	5,5,5	0.56	0
3	ACT	A	1906	-	3,3,3	1.33	0	3,3,3	0.78	0
4	GOL	B	1907	-	5,5,5	0.50	0	5,5,5	1.42	1 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1907	-	-	3/4/4/4	-
4	GOL	B	1907	-	-	3/4/4/4	-
4	GOL	A	1908	-	-	4/4/4/4	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	1907	GOL	O2-C2-C3	2.42	119.19	109.18
4	A	1907	GOL	O2-C2-C3	2.04	117.64	109.18

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1907	GOL	O1-C1-C2-O2
4	A	1907	GOL	O1-C1-C2-C3
4	B	1907	GOL	O1-C1-C2-C3
4	A	1908	GOL	O1-C1-C2-O2
4	A	1907	GOL	C1-C2-C3-O3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	851/1032 (82%)	-0.59	3 (0%) 88 90	16, 25, 35, 52	2 (0%)
1	B	849/1032 (82%)	-0.59	3 (0%) 88 90	16, 25, 35, 51	0
All	All	1700/2064 (82%)	-0.59	6 (0%) 88 90	16, 25, 35, 52	2 (0%)

The worst 5 of 6 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	430	GLU	3.5
1	A	431	GLU	3.2
1	B	49	GLY	3.0
1	A	49	GLY	2.3
1	A	432	LYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	CD	A	1903	1/1	0.45	0.42	295,295,295,295	0
3	ACT	A	1906	4/4	0.61	0.18	32,32,33,33	0
3	ACT	B	1906	4/4	0.65	0.15	24,25,26,27	0
4	GOL	B	1907	6/6	0.71	0.15	38,39,40,41	0
4	GOL	A	1907	6/6	0.81	0.14	36,39,40,41	0
4	GOL	A	1908	6/6	0.86	0.11	39,43,46,47	0
2	CD	B	1905	1/1	0.93	0.15	102,102,102,102	0
2	CD	B	1902	1/1	0.95	0.05	48,48,48,48	0
2	CD	B	1904	1/1	0.95	0.07	100,100,100,100	0
2	CD	A	1904	1/1	0.96	0.07	105,105,105,105	0
2	CD	A	1902	1/1	0.96	0.04	53,53,53,53	0
2	CD	A	1900	1/1	0.96	0.04	47,47,47,47	0
2	CD	A	1905	1/1	0.98	0.06	76,76,76,76	0
2	CD	B	1900	1/1	0.99	0.02	25,25,25,25	0
2	CD	B	1903	1/1	1.00	0.01	34,34,34,34	0
2	CD	A	1901	1/1	1.00	0.01	37,37,37,37	0
2	CD	B	1901	1/1	1.00	0.02	25,25,25,25	0
2	CD	A	1909	1/1	1.00	0.01	28,28,28,28	0

6.5 Other polymers ⓘ

There are no such residues in this entry.