



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 9, 2026 – 08:07 PM UTC

PDB ID : 1W0M / pdb_00001w0m
Title : Triosephosphate isomerase from Thermoproteus tenax
Authors : Walden, H.; Taylor, G.; Lorentzen, E.; Pohl, E.; Lilie, H.; Schramm, A.; Knura, T.; Stubbe, K.; Tjaden, B.; Hensel, R.
Deposited on : 2004-06-08
Resolution : 2.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	NOT EXECUTED
EDS	:	NOT EXECUTED
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.50 Å.



Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	6492 (2.50-2.50)
Ramachandran outliers	187476	6378 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	226	 83% 16% .
1	B	226	 76% 21% .
1	C	226	 81% 18% .
1	D	226	 77% 22% .
1	E	226	 78% 19% .
1	F	226	 82% 16% .
1	G	226	 71% 27% ..
1	H	226	 77% 21% .

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 13752 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called TRIOSEPHOSPHATE ISOMERASE.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			
1	B	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			
1	C	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			
1	D	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			
1	E	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			
1	F	225	Total	C	N	O	S	0	0	1
			1621	1024	290	305	2			
1	G	224	Total	C	N	O	S	0	0	1
			1612	1019	289	302	2			
1	H	226	Total	C	N	O	S	0	0	1
			1629	1030	291	306	2			

- Molecule 2 is PHOSPHATE ION (CCD ID: PO4) (formula: O₄P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	P	0	0
			5	4	1		
2	B	1	Total	O	P	0	0
			5	4	1		
2	C	1	Total	O	P	0	0
			5	4	1		
2	D	1	Total	O	P	0	0
			5	4	1		
2	E	1	Total	O	P	0	0
			5	4	1		
2	F	1	Total	O	P	0	0
			5	4	1		
2	G	1	Total	O	P	0	0
			5	4	1		
2	H	1	Total	O	P	0	0
			5	4	1		

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	102	Total	O	0	0
			102	102		
3	B	97	Total	O	0	0
			97	97		
3	C	139	Total	O	0	0
			139	139		
3	D	110	Total	O	0	0
			110	110		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	E	89	Total 89	O 89	0	0
3	F	62	Total 62	O 62	0	0
3	G	47	Total 47	O 47	0	0
3	H	59	Total 59	O 59	0	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


• Molecule 1: TRIOSEPHOSPHATE ISOMERASE




• Molecule 1: TRIOSEPHOSPHATE ISOMERASE




• Molecule 1: TRIOSEPHOSPHATE ISOMERASE




• Molecule 1: TRIOSEPHOSPHATE ISOMERASE






• Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain E: 78% 19%




• Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain F: 82% 16%



• Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain G: 71% 27%



• Molecule 1: TRIOSEPHOSPHATE ISOMERASE

Chain H: 77% 21%



4 Data and refinement statistics

Xtriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, α , β , γ	186.89Å 186.89Å 287.76Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.50	Depositor
% Data completeness (in resolution range)	96.5 (30.00-2.50)	Depositor
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.197 , 0.226	Depositor
Estimated twinning fraction	No twinning to report.	Xtriage
Total number of atoms	13752	wwPDB-VP
Average B, all atoms (Å ²)	37.0	wwPDB-VP

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.67	1/1653 (0.1%)	1.00	0/2250
1	B	0.67	1/1653 (0.1%)	1.05	7/2250 (0.3%)
1	C	0.73	1/1653 (0.1%)	1.07	5/2250 (0.2%)
1	D	0.71	1/1653 (0.1%)	1.06	6/2250 (0.3%)
1	E	0.65	2/1653 (0.1%)	1.22	11/2250 (0.5%)
1	F	0.64	1/1645 (0.1%)	1.03	2/2239 (0.1%)
1	G	0.55	1/1636 (0.1%)	0.99	1/2227 (0.0%)
1	H	0.57	1/1653 (0.1%)	1.03	5/2250 (0.2%)
All	All	0.65	9/13199 (0.1%)	1.06	37/17966 (0.2%)

The worst 5 of 9 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	1	MET	SD-CE	-6.95	1.62	1.79
1	G	223	SER	C-N	-5.78	1.25	1.33
1	B	225	LEU	C-N	-5.61	1.25	1.33
1	A	225	LEU	C-N	-5.30	1.25	1.33
1	D	225	LEU	C-N	-5.29	1.25	1.33

The worst 5 of 37 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	E	3	LEU	CA-C-N	-19.69	98.57	119.83
1	E	3	LEU	C-N-CA	-19.69	98.57	119.83
1	E	3	LEU	N-CA-C	7.93	127.33	109.81
1	G	116	ASP	N-CA-C	-6.78	101.39	110.55
1	C	210	ASP	CA-C-N	6.58	126.08	119.24

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1629	0	1702	50	0
1	B	1629	0	1702	52	0
1	C	1629	0	1702	46	0
1	D	1629	0	1702	48	0
1	E	1629	0	1702	43	0
1	F	1621	0	1691	39	0
1	G	1612	0	1685	62	0
1	H	1629	0	1702	46	0
2	A	5	0	0	0	0
2	B	5	0	0	0	0
2	C	5	0	0	0	0
2	D	5	0	0	0	0
2	E	5	0	0	0	0
2	F	5	0	0	0	0
2	G	5	0	0	0	0
2	H	5	0	0	0	0
3	A	102	0	0	2	0
3	B	97	0	0	6	0
3	C	139	0	0	6	0
3	D	110	0	0	6	0
3	E	89	0	0	2	0
3	F	62	0	0	2	0
3	G	47	0	0	4	0
3	H	59	0	0	5	0
All	All	13752	0	13588	363	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 14.

The worst 5 of 363 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LEU:HB3	1:A:4:PRO:HD3	1.29	1.14
1:A:3:LEU:CB	1:A:4:PRO:HD3	1.83	1.07
1:B:3:LEU:CB	1:B:4:PRO:HD3	1.92	0.99

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:93:HIS:HD2	1:H:95:GLU:H	1.13	0.96
1:H:117:VAL:H	1:H:136:HIS:HD2	1.14	0.96

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/226 (99%)	215 (96%)	7 (3%)	2 (1%)	14	27
1	B	224/226 (99%)	213 (95%)	9 (4%)	2 (1%)	14	27
1	C	224/226 (99%)	216 (96%)	6 (3%)	2 (1%)	14	27
1	D	224/226 (99%)	215 (96%)	7 (3%)	2 (1%)	14	27
1	E	224/226 (99%)	214 (96%)	8 (4%)	2 (1%)	14	27
1	F	223/226 (99%)	209 (94%)	12 (5%)	2 (1%)	14	27
1	G	222/226 (98%)	206 (93%)	13 (6%)	3 (1%)	9	17
1	H	224/226 (99%)	214 (96%)	8 (4%)	2 (1%)	14	27
All	All	1789/1808 (99%)	1702 (95%)	70 (4%)	17 (1%)	12	24

5 of 17 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	3	LEU
1	B	3	LEU
1	C	3	LEU
1	D	3	LEU
1	E	225	LEU

5.3.2 Protein sidechains [i](#)

There are no protein residues with a non-rotameric sidechain to report in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	PO4	D	1226	-	4,4,4	1.39	0	6,6,6	0.47	0
2	PO4	A	1226	-	4,4,4	1.40	0	6,6,6	0.44	0
2	PO4	G	1224	-	4,4,4	1.60	0	6,6,6	0.49	0
2	PO4	C	1226	-	4,4,4	1.45	0	6,6,6	0.47	0
2	PO4	F	1225	-	4,4,4	1.35	0	6,6,6	0.46	0
2	PO4	H	1226	-	4,4,4	1.63	0	6,6,6	0.41	0
2	PO4	B	1226	-	4,4,4	1.48	0	6,6,6	0.46	0
2	PO4	E	1226	-	4,4,4	1.36	0	6,6,6	0.49	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates [i](#)

EDS was not executed - this section is therefore empty.

6.4 Ligands [i](#)

EDS was not executed - this section is therefore empty.

6.5 Other polymers [i](#)

EDS was not executed - this section is therefore empty.