



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 15, 2026 – 01:42 AM UTC

PDB ID : 5W0B / pdb_00005w0b
Title : Structure of human TUT7 catalytic module (CM)
Authors : Faehnle, C.R.; Walleshauser, J.; Joshua-Tor, L.
Deposited on : 2017-05-30
Resolution : 2.61 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.61 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4951 (2.64-2.60)
Clashscore	190562	5303 (2.64-2.60)
Ramachandran outliers	187476	5217 (2.64-2.60)
Sidechain outliers	187428	5217 (2.64-2.60)
RSRZ outliers	180081	4950 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	389	14% 82% • 14%
1	B	389	7% 85% • 10%
1	C	389	12% 84% 6% 9%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16803 atoms, of which 8381 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Terminal uridylyltransferase 7.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	334	Total	C	H	N	O	S	0	0	0
			5373	1717	2702	454	481	19			
1	B	349	Total	C	H	N	O	S	0	0	0
			5659	1835	2832	466	512	14			
1	C	353	Total	C	H	N	O	S	0	0	0
			5710	1857	2847	473	519	14			

There are 21 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	977	GLY	-	expression tag	UNP Q5VYS8
A	978	ALA	-	expression tag	UNP Q5VYS8
A	979	GLY	-	expression tag	UNP Q5VYS8
A	980	ALA	-	expression tag	UNP Q5VYS8
A	981	GLY	-	expression tag	UNP Q5VYS8
A	982	SER	-	expression tag	UNP Q5VYS8
A	1060	ALA	ASP	engineered mutation	UNP Q5VYS8
B	977	GLY	-	expression tag	UNP Q5VYS8
B	978	ALA	-	expression tag	UNP Q5VYS8
B	979	GLY	-	expression tag	UNP Q5VYS8
B	980	ALA	-	expression tag	UNP Q5VYS8
B	981	GLY	-	expression tag	UNP Q5VYS8
B	982	SER	-	expression tag	UNP Q5VYS8
B	1060	ALA	ASP	engineered mutation	UNP Q5VYS8
C	977	GLY	-	expression tag	UNP Q5VYS8
C	978	ALA	-	expression tag	UNP Q5VYS8
C	979	GLY	-	expression tag	UNP Q5VYS8
C	980	ALA	-	expression tag	UNP Q5VYS8
C	981	GLY	-	expression tag	UNP Q5VYS8
C	982	SER	-	expression tag	UNP Q5VYS8
C	1060	ALA	ASP	engineered mutation	UNP Q5VYS8

- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is SULFATE ION (CCD ID: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	O	S	0	0
			5	4	1		
3	A	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	B	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		
3	C	1	Total	O	S	0	0
			5	4	1		

- Molecule 4 is IODIDE ION (CCD ID: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total	I	0	0
			1	1		
4	C	2	Total	I	0	0
			2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	5	Total 5	O 5	0	0
5	B	2	Total 2	O 2	0	0
5	C	15	Total 15	O 15	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Terminal uridylyltransferase 7



• Molecule 1: Terminal uridylyltransferase 7



• Molecule 1: Terminal uridylyltransferase 7



ASP
CYS
PRO
MET
ARG
ARG
LYS

4 Data and refinement statistics

Property	Value	Source
Space group	P 61	Depositor
Cell constants a, b, c, α , β , γ	141.23Å 141.23Å 174.47Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	61.16 – 2.61 61.16 – 2.61	Depositor EDS
% Data completeness (in resolution range)	99.9 (61.16-2.61) 99.9 (61.16-2.61)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.28 (at 2.62Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.231 , 0.255 0.239 , 0.260	Depositor DCC
R_{free} test set	3007 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	66.4	Xtriage
Anisotropy	0.348	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 55.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.037 for h,-h-k,-l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	16803	wwPDB-VP
Average B, all atoms (Å ²)	95.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.34% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, ZN, IOD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.13	0/2728	0.31	0/3682
1	B	0.12	0/2896	0.30	0/3921
1	C	0.14	0/2932	0.34	0/3965
All	All	0.13	0/8556	0.32	0/11568

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2671	2702	2699	10	0
1	B	2827	2832	2838	11	0
1	C	2863	2847	2872	15	0
2	A	1	0	0	0	0
3	A	10	0	0	1	0
3	B	15	0	0	1	0
3	C	10	0	0	0	0
4	B	1	0	0	0	0
4	C	2	0	0	0	0
5	A	5	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	B	2	0	0	0	0
5	C	15	0	0	0	0
All	All	8422	8381	8409	34	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (34) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:1306:ASN:OD1	1:B:1309:ARG:NH2	2.13	0.81
1:C:1306:ASN:OD1	1:C:1309:ARG:NH2	2.35	0.60
1:A:986:GLN:OE1	1:A:987:LEU:N	2.36	0.58
1:C:1037:PHE:O	1:C:1040:THR:OG1	2.19	0.53
1:A:1349:ILE:HD12	1:B:1032:PHE:HA	1.88	0.53
1:B:1046:GLY:H	1:B:1131:THR:HG21	1.75	0.52
1:A:1349:ILE:CD1	1:B:1032:PHE:HA	2.40	0.51
1:A:1191:VAL:HG11	1:A:1260:ARG:HA	1.92	0.51
1:C:1121:SER:OG	1:C:1124:ASN:OD1	2.13	0.50
1:A:1318:PHE:CD1	1:A:1319:PRO:HA	2.47	0.48
1:A:1152:LYS:NZ	3:A:1402:SO4:O2	2.39	0.47
1:B:1039:GLY:O	1:B:1065:ILE:HA	2.15	0.46
1:C:1100:THR:HA	1:C:1105:PRO:HB3	1.98	0.46
1:B:1130:ASN:OD1	1:B:1131:THR:N	2.48	0.46
1:C:1106:ILE:HG22	1:C:1121:SER:HB3	1.98	0.46
1:C:1074:LEU:O	1:C:1076:CYS:N	2.50	0.44
1:A:1344:ASP:OD1	1:A:1345:ARG:N	2.51	0.43
1:C:988:GLU:O	1:C:1308:ARG:NH2	2.39	0.43
1:C:1081:GLU:OE2	1:C:1105:PRO:HG2	2.18	0.43
1:C:1340:LEU:HD23	1:C:1341:ALA:O	2.18	0.43
1:B:1088:ARG:NH2	1:B:1096:ILE:O	2.52	0.43
1:A:1347:CYS:HB3	1:A:1350:CYS:SG	2.59	0.43
1:C:1264:LEU:HD21	1:C:1269:LYS:HD2	2.01	0.43
1:B:1152:LYS:NZ	3:B:1402:SO4:O2	2.49	0.42
1:C:1074:LEU:HD11	1:C:1080:ILE:HG12	2.01	0.42
1:C:1022:ARG:NH2	1:C:1057:SER:O	2.53	0.42
1:C:1075:ASP:C	1:C:1077:VAL:H	2.28	0.42
1:B:1108:LYS:NZ	1:B:1119:ASP:OD1	2.51	0.42
1:C:1252:LYS:HD3	1:C:1252:LYS:O	2.20	0.42
1:C:1124:ASN:O	1:C:1128:LEU:HD12	2.20	0.41
1:A:1347:CYS:SG	1:A:1349:ILE:HG23	2.60	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:1111:HIS:CE1	1:A:1113:ARG:HB2	2.56	0.41
1:B:1104:VAL:O	1:B:1106:ILE:HG23	2.21	0.40
1:B:1127:ALA:O	1:B:1131:THR:HG23	2.20	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	326/389 (84%)	318 (98%)	8 (2%)	0	100	100
1	B	343/389 (88%)	338 (98%)	5 (2%)	0	100	100
1	C	347/389 (89%)	333 (96%)	13 (4%)	1 (0%)	36	56
All	All	1016/1167 (87%)	989 (97%)	26 (3%)	1 (0%)	48	69

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	1075	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	300/348 (86%)	299 (100%)	1 (0%)	86	94

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	316/348 (91%)	315 (100%)	1 (0%)	86	94
1	C	319/348 (92%)	318 (100%)	1 (0%)	86	94
All	All	935/1044 (90%)	932 (100%)	3 (0%)	86	94

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1349	ILE
1	B	996	LYS
1	C	1320	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	1028	ASN
1	C	1254	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 4 are monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	SO4	C	1402	-	4,4,4	0.22	0	6,6,6	0.07	0
3	SO4	B	1402	-	4,4,4	0.23	0	6,6,6	0.06	0
3	SO4	A	1403	-	4,4,4	0.24	0	6,6,6	0.07	0
3	SO4	B	1401	-	4,4,4	0.22	0	6,6,6	0.08	0
3	SO4	C	1401	-	4,4,4	0.25	0	6,6,6	0.08	0
3	SO4	B	1403	-	4,4,4	0.24	0	6,6,6	0.10	0
3	SO4	A	1402	-	4,4,4	0.23	0	6,6,6	0.08	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	1402	SO4	1	0
3	A	1402	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	334/389 (85%)	0.89	54 (16%) 4 3	61, 101, 173, 249	0
1	B	349/389 (89%)	0.45	26 (7%) 20 17	47, 83, 139, 166	0
1	C	353/389 (90%)	0.59	48 (13%) 7 5	52, 78, 152, 191	0
All	All	1036/1167 (88%)	0.64	128 (12%) 8 6	47, 87, 158, 249	0

All (128) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	1084	ALA	6.9
1	C	1104	VAL	6.8
1	A	1340	LEU	6.4
1	A	1342	PRO	6.3
1	B	1275	TYR	6.2
1	C	1080	ILE	5.7
1	A	1361	PRO	5.6
1	C	1081	GLU	5.4
1	A	1356	PHE	5.4
1	A	1341	ALA	5.0
1	C	1083	LEU	5.0
1	C	1069	GLU	4.8
1	C	1075	ASP	4.7
1	C	1077	VAL	4.6
1	A	1113	ARG	4.5
1	C	1338	GLY	4.4
1	C	1340	LEU	4.4
1	C	1124	ASN	4.3
1	C	1076	CYS	4.2
1	C	1073	GLY	4.2
1	C	1078	ARG	4.2
1	A	1344	ASP	4.1
1	A	1349	ILE	4.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1125	THR	4.1
1	C	1275	TYR	4.0
1	C	1339	GLU	3.7
1	B	1201	LYS	3.7
1	C	1337	GLU	3.6
1	A	989	PRO	3.6
1	B	1276	ILE	3.6
1	A	1350	CYS	3.5
1	A	1345	ARG	3.5
1	B	985	ILE	3.5
1	C	1199	GLU	3.5
1	C	1315	VAL	3.4
1	C	1322	TYR	3.4
1	B	1271	TRP	3.3
1	A	1343	ASN	3.3
1	A	1314	PRO	3.3
1	C	1126	LEU	3.3
1	A	1346	CYS	3.3
1	C	1074	LEU	3.2
1	C	1099	ILE	3.2
1	B	1125	THR	3.2
1	C	1079	THR	3.2
1	B	1266	THR	3.2
1	A	1292	LEU	3.1
1	C	1295	LYS	3.0
1	A	1318	PHE	3.0
1	A	1293	SER	2.9
1	A	1195	ILE	2.9
1	C	1318	PHE	2.9
1	A	1323	PRO	2.9
1	B	1126	LEU	2.9
1	A	1319	PRO	2.9
1	B	1124	ASN	2.9
1	A	993	LEU	2.9
1	B	1315	VAL	2.8
1	A	1276	ILE	2.8
1	B	1223	TYR	2.8
1	C	1248	GLU	2.8
1	C	1113	ARG	2.8
1	A	1192	LEU	2.8
1	C	1082	GLU	2.8
1	A	1325	LYS	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	C	1334	VAL	2.8
1	A	1051	GLY	2.7
1	A	1322	TYR	2.7
1	B	1016	ILE	2.7
1	A	1320	LYS	2.7
1	A	1265	THR	2.7
1	C	1320	LYS	2.7
1	A	1321	ASP	2.7
1	B	1082	GLU	2.6
1	B	1316	LYS	2.6
1	C	1317	GLY	2.6
1	C	986	GLN	2.6
1	A	1248	GLU	2.6
1	A	1348	ARG	2.6
1	B	1088	ARG	2.5
1	A	1114	SER	2.5
1	B	1108	LYS	2.5
1	B	1128	LEU	2.5
1	C	1252	LYS	2.5
1	B	1318	PHE	2.5
1	A	1228	GLY	2.5
1	A	1357	MET	2.5
1	C	1336	THR	2.5
1	A	1263	LEU	2.4
1	C	987	LEU	2.4
1	C	1270	GLN	2.4
1	C	1267	PHE	2.4
1	A	1266	THR	2.4
1	A	1254	HIS	2.4
1	B	1195	ILE	2.4
1	A	1353	ILE	2.3
1	A	1190	PRO	2.3
1	B	1273	SER	2.3
1	A	991	PRO	2.3
1	C	1085	ARG	2.3
1	C	1323	PRO	2.3
1	A	1112	LEU	2.3
1	B	1113	ARG	2.3
1	C	1319	PRO	2.3
1	A	1264	LEU	2.3
1	B	1292	LEU	2.3
1	A	1056	GLN	2.2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	1334	VAL	2.2
1	A	1358	LYS	2.2
1	B	1073	GLY	2.2
1	C	993	LEU	2.2
1	A	1071	ALA	2.2
1	A	1347	CYS	2.2
1	B	1319	PRO	2.2
1	A	1101	THR	2.2
1	A	1191	VAL	2.2
1	C	1122	LEU	2.1
1	A	1354	GLY	2.1
1	B	1067	GLY	2.1
1	C	1341	ALA	2.1
1	A	1277	VAL	2.1
1	C	1086	VAL	2.1
1	A	1260	ARG	2.0
1	B	1293	SER	2.0
1	C	1342	PRO	2.0
1	A	987	LEU	2.0
1	A	1035	GLN	2.0
1	C	1309	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1403	5/5	0.60	0.16	139,139,139,140	0
3	SO4	C	1401	5/5	0.87	0.12	111,111,112,113	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	SO4	A	1402	5/5	0.91	0.10	94,96,96,97	0
2	ZN	A	1401	1/1	0.92	0.08	107,107,107,107	0
3	SO4	C	1402	5/5	0.92	0.16	90,91,92,94	0
3	SO4	B	1401	5/5	0.93	0.10	85,86,86,86	0
3	SO4	B	1402	5/5	0.94	0.09	80,83,86,87	0
3	SO4	B	1403	5/5	0.94	0.09	91,93,96,98	0
4	IOD	B	1404	1/1	0.97	0.14	152,152,152,152	0
4	IOD	C	1404	1/1	0.98	0.23	144,144,144,144	0
4	IOD	C	1403	1/1	0.99	0.11	103,103,103,103	0

6.5 Other polymers [i](#)

There are no such residues in this entry.