



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:25 AM EDT

PDB ID : 3W15 / pdb_00003w15
Title : Structure of peroxisomal targeting signal 2 (PTS2) of *Saccharomyces cerevisiae* 3-ketoacyl-CoA thiolase in complex with Pex7p and Pex21p
Authors : Pan, D.; Nakatsu, T.; Kato, H.
Deposited on : 2012-11-06
Resolution : 1.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	368	9% 82% 10% 9%
2	B	101	15% 69% 29%
3	C	389	% 97% .

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6824 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Peroxisomal targeting signal 2 receptor.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	336	2650	1688	455	492	15	0	7	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-1	GLY	-	expression tag	UNP P39108
A	0	SER	-	expression tag	UNP P39108
A	?	-	GLU	deletion	UNP P39108
A	?	-	SER	deletion	UNP P39108
A	?	-	ALA	deletion	UNP P39108
A	?	-	THR	deletion	UNP P39108
A	?	-	ILE	deletion	UNP P39108
A	?	-	LYS	deletion	UNP P39108
A	?	-	ARG	deletion	UNP P39108
A	?	-	THR	deletion	UNP P39108
A	?	-	VAL	deletion	UNP P39108

- Molecule 2 is a protein called Peroxisomal membrane protein PEX21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	72	548	338	93	114	3	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	188	GLY	-	expression tag	UNP P50091
B	189	SER	-	expression tag	UNP P50091


- Molecule 3 is a protein called 3-ketoacyl-CoA thiolase, peroxisomal, Maltose-binding periplasmic protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	389	Total	C	N	O	S	0	4	0
			3021	1939	491	584	7			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	-1	GLY	-	expression tag	UNP P27796
C	0	SER	-	expression tag	UNP P27796
C	16	ARG	-	linker	UNP P0AEX9
C	17	SER	-	linker	UNP P0AEX9

- Molecule 4 is NITRATE ION (CCD ID: NO3) (formula: NO₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	N	O	0	0
			4	1	3		
4	A	1	Total	N	O	0	0
			4	1	3		
4	B	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	C	1	Total	N	O	0	0
			4	1	3		
4	C	1	Total	N	O	0	0
			4	1	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	1	Total	Mg	0	0
			1	1		
5	C	5	Total	Mg	0	0
			5	5		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	136	Total	O	0	0
			136	136		
6	B	28	Total	O	0	0
			28	28		
6	C	399	Total	O	0	0
			399	399		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Peroxisomal targeting signal 2 receptor



- Molecule 2: Peroxisomal membrane protein PEX21



- Molecule 3: 3-ketoacyl-CoA thiolase, peroxisomal, Maltose-binding periplasmic protein



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	78.89Å 52.75Å 97.57Å 90.00° 106.59° 90.00°	Depositor
Resolution (Å)	32.20 – 1.80 32.20 – 1.80	Depositor EDS
% Data completeness (in resolution range)	99.7 (32.20-1.80) 99.7 (32.20-1.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	6.94 (at 1.80Å)	Xtriage
Refinement program	REFMAC refmac_5.5.0102	Depositor
R, R_{free}	0.189 , 0.224 0.188 , 0.225	Depositor DCC
R_{free} test set	3618 reflections (5.05%)	wwPDB-VP
Wilson B-factor (Å ²)	18.0	Xtriage
Anisotropy	0.667	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 38.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6824	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.09% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NO3, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/2738	0.73	1/3725 (0.0%)
2	B	0.44	0/550	0.79	0/734
3	C	0.50	1/3103 (0.0%)	0.73	0/4214
All	All	0.48	1/6391 (0.0%)	0.73	1/8673 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	149	ILE	CA-CB	5.45	1.57	1.54

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	218	HIS	N-CA-C	6.04	117.69	110.44

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2650	0	2507	22	0
2	B	548	0	536	1	0
3	C	3021	0	2975	5	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	8	0	0	0	0
4	B	4	0	0	0	0
4	C	24	0	0	0	0
5	A	1	0	0	0	0
5	C	5	0	0	0	0
6	A	136	0	0	0	0
6	B	28	0	0	0	0
6	C	399	0	0	0	0
All	All	6824	0	6018	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:249:ASP:OD2	1:A:251:ARG:HG2	1.83	0.79
1:A:125:TRP:CD1	1:A:176[A]:CYS:HG	2.08	0.72
1:A:116:ASN:HD22	1:A:187:GLN:HE21	1.46	0.64
1:A:149:GLU:O	1:A:153:MET:HG2	2.07	0.55
1:A:113:ASN:O	1:A:117:ARG:HA	2.09	0.52
1:A:125:TRP:CD1	1:A:176[A]:CYS:SG	3.05	0.50
2:B:267:ASN:HB3	2:B:269:GLU:HG2	1.94	0.50
1:A:43:LEU:HD12	1:A:368:LEU:HD22	1.94	0.49
1:A:185:HIS:CD2	1:A:233:PRO:HA	2.47	0.49
1:A:219:SER:HB3	3:C:67:VAL:HB	1.95	0.48
1:A:308[A]:CYS:SG	1:A:345:VAL:HG11	2.54	0.48
1:A:121:LEU:CD1	1:A:131:ILE:HG12	2.44	0.48
1:A:155:ASP:O	1:A:158:ASN:HB2	2.13	0.48
1:A:125:TRP:NE1	1:A:176[A]:CYS:HG	2.14	0.45
3:C:95:GLU:HG3	3:C:119:LYS:HB3	1.99	0.45
1:A:125:TRP:NE1	1:A:176[A]:CYS:SG	2.90	0.44
1:A:234:TYR:O	1:A:250:ILE:HG12	2.18	0.43
3:C:308[B]:GLU:OE2	3:C:312:LYS:HD3	2.19	0.43
3:C:246:PRO:HA	3:C:249:TRP:CE2	2.55	0.41
1:A:121:LEU:HD21	1:A:190:VAL:HG13	2.02	0.41
1:A:193[A]:CYS:SG	1:A:226:CYS:HB3	2.60	0.41
1:A:312:ARG:HB2	1:A:335:ILE:HD11	2.02	0.41
1:A:2:LEU:HB2	1:A:370:VAL:HB	2.03	0.41
1:A:24:LEU:C	1:A:24:LEU:HD12	2.45	0.40
1:A:127:GLY:HA2	1:A:175:ASN:O	2.22	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:C:188:TYR:OH	3:C:191:GLY:HA2	2.21	0.40
1:A:121:LEU:HD13	1:A:131:ILE:HG12	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	335/368 (91%)	327 (98%)	8 (2%)	0	100	100
2	B	66/101 (65%)	63 (96%)	3 (4%)	0	100	100
3	C	391/389 (100%)	386 (99%)	5 (1%)	0	100	100
All	All	792/858 (92%)	776 (98%)	16 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	287/320 (90%)	283 (99%)	4 (1%)	59	52
2	B	65/92 (71%)	65 (100%)	0	100	100
3	C	313/315 (99%)	313 (100%)	0	100	100
All	All	665/727 (92%)	661 (99%)	4 (1%)	81	77

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	LEU
1	A	66	LEU
1	A	181[A]	GLN
1	A	181[B]	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	187	GLN
1	A	316	ASN
3	C	299	ASN
3	C	366	ASN
3	C	372	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 15 ligands modelled in this entry, 6 are monoatomic - leaving 9 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	NO3	B	301	-	1,3,3	3.15	1 (100%)	0,3,3	-	-
4	NO3	A	402	-	1,3,3	3.24	1 (100%)	0,3,3	-	-
4	NO3	C	406	-	1,3,3	3.21	1 (100%)	0,3,3	-	-
4	NO3	A	401	-	1,3,3	3.29	1 (100%)	0,3,3	-	-
4	NO3	C	404	-	1,3,3	3.21	1 (100%)	0,3,3	-	-
4	NO3	C	401	-	1,3,3	3.04	1 (100%)	0,3,3	-	-
4	NO3	C	405	-	1,3,3	3.07	1 (100%)	0,3,3	-	-
4	NO3	C	403	-	1,3,3	3.30	1 (100%)	0,3,3	-	-
4	NO3	C	402	-	1,3,3	3.21	1 (100%)	0,3,3	-	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	C	403	NO3	O1-N	3.30	1.40	1.24
4	A	401	NO3	O1-N	3.29	1.40	1.24
4	A	402	NO3	O1-N	3.24	1.40	1.24
4	C	402	NO3	O1-N	3.21	1.40	1.24
4	C	406	NO3	O1-N	3.21	1.40	1.24
4	C	404	NO3	O1-N	3.21	1.40	1.24
4	B	301	NO3	O1-N	3.15	1.39	1.24
4	C	405	NO3	O1-N	3.07	1.39	1.24
4	C	401	NO3	O1-N	3.04	1.39	1.24

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues ⓘ

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2		OWAB(Å ²)	Q < 0.9
1	A	336/368 (91%)	0.41	33 (9%)	13 11	8, 20, 45, 49	7 (2%)
2	B	72/101 (71%)	1.27	15 (20%)	2 2	15, 36, 47, 49	0
3	C	389/389 (100%)	-0.21	3 (0%)	82 83	9, 16, 27, 33	4 (1%)
All	All	797/858 (92%)	0.19	51 (6%)	25 24	8, 18, 42, 49	11 (1%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	154	VAL	5.8
1	A	151	THR	5.5
1	A	158	ASN	4.5
1	A	152	LYS	4.3
1	A	1	MET	4.1
1	A	159	ALA	4.0
1	A	49	GLY	4.0
2	B	224	SER	3.9
1	A	150	ILE	3.6
2	B	265	SER	3.5
1	A	161	ILE	3.4
1	A	331	THR	3.4
1	A	148	LEU	3.3
2	B	263	PHE	3.3
1	A	160	ILE	3.2
1	A	147	PRO	3.2
2	B	246	GLY	3.2
1	A	155	ASP	3.1
2	B	210	CYS	3.1
2	B	250	LEU	3.0
1	A	47	ARG	3.0
2	B	197	SER	3.0
1	A	353	TRP	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	268	GLY	2.8
3	C	117	ASN	2.8
1	A	156	PRO	2.8
3	C	191	GLY	2.7
2	B	211	THR	2.7
1	A	356	PRO	2.7
1	A	354	GLY	2.7
1	A	330	ALA	2.6
1	A	162	LEU	2.6
2	B	249	THR	2.5
1	A	153	MET	2.5
1	A	252	MET	2.4
1	A	327	SER	2.4
3	C	190	ASN	2.4
1	A	48	SER	2.4
2	B	267	ASN	2.4
2	B	227	VAL	2.4
1	A	273	SER	2.4
1	A	355	LYS	2.4
1	A	157	LEU	2.3
1	A	275	ILE	2.3
1	A	328	THR	2.3
2	B	248	VAL	2.2
2	B	247	ASP	2.2
1	A	146	ARG	2.2
1	A	251	ARG	2.2
2	B	266	ASN	2.1
1	A	21	ASP	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	NO3	C	405	4/4	0.80	0.17	30,30,30,30	0
4	NO3	A	401	4/4	0.81	0.16	40,40,40,41	0
4	NO3	C	402	4/4	0.84	0.20	31,32,33,33	0
4	NO3	C	404	4/4	0.84	0.20	36,36,36,37	0
4	NO3	A	402	4/4	0.84	0.13	48,48,48,48	0
4	NO3	C	406	4/4	0.84	0.17	47,47,47,47	0
4	NO3	C	401	4/4	0.91	0.10	25,26,26,27	0
4	NO3	C	403	4/4	0.92	0.11	26,27,27,27	0
4	NO3	B	301	4/4	0.92	0.09	28,28,28,28	0
5	MG	C	408	1/1	0.95	0.05	31,31,31,31	0
5	MG	C	411	1/1	0.95	0.05	39,39,39,39	0
5	MG	C	409	1/1	0.97	0.12	24,24,24,24	0
5	MG	C	410	1/1	0.98	0.05	18,18,18,18	0
5	MG	A	403	1/1	0.99	0.07	22,22,22,22	0
5	MG	C	407	1/1	0.99	0.10	19,19,19,19	0

6.5 Other polymers ⓘ

There are no such residues in this entry.