



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 12:25 PM UTC

PDB ID : 3W1Q / pdb_00003w1q
Title : Structure of Trypanosoma cruzi dihydroorotate dehydrogenase in complex with NL-2
Authors : Inaoka, D.K.; Iida, M.; Tabuchi, T.; Lee, N.; Matsuoka, S.; Shiba, T.; Sakamoto, K.; Suzuki, S.; Balogun, E.O.; Nara, T.; Aoki, T.; Inoue, M.; Honma, T.; Tanaka, A.; Harada, S.; Kita, K.
Deposited on : 2012-11-20
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	313	4% 86% 12% .
1	B	313	2% 90% 9% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	YRO	A	403	-	-	X	-
5	GOL	A	406	-	-	X	-

2 Entry composition [i](#)


There are 6 unique types of molecules in this entry. The entry contains 5480 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Dihydroorotate dehydrogenase (fumarate).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	313	Total	C	N	O	S	0	0	0
			2388	1530	396	444	18			
1	B	313	Total	C	N	O	S	0	0	0
			2388	1530	396	444	18			

- Molecule 2 is FLAVIN MONONUCLEOTIDE (CCD ID: FMN) (formula: $C_{17}H_{21}N_4O_9P$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	P	0	0
			31	17	4	9	1		
2	B	1	Total	C	N	O	P	0	0
			31	17	4	9	1		

- Molecule 3 is COBALT HEXAMMINE(III) (CCD ID: NCO) (formula: $CoH_{18}N_6$).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	Co	N	0	0
			7	1	6		

- Molecule 4 is 5-(3,3-dimethylbutyl)-2,6-dioxo-1,2,3,6-tetrahydropyrimidine-4-carboxylic acid (CCD ID: YRO) (formula: C₁₁H₁₆N₂O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			17	11	2	4		
4	B	1	Total	C	N	O	0	0
			17	11	2	4		

- Molecule 5 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	A	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0
5	B	1	Total C O 6 3 3	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	287	Total O 287 287	0	0
6	B	272	Total O 272 272	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



- Molecule 1: Dihydroorotate dehydrogenase (fumarate)



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	68.11Å 71.89Å 129.55Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	34.63 – 1.85 34.63 – 1.85	Depositor EDS
% Data completeness (in resolution range)	96.1 (34.63-1.85) 96.1 (34.63-1.85)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	7.47 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.7.0029	Depositor
R, R_{free}	0.151 , 0.190 (Not available) , 0.181	Depositor DCC
R_{free} test set	2793 reflections (5.07%)	wwPDB-VP
Wilson B-factor (Å ²)	12.0	Xtriage
Anisotropy	0.073	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 39.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	5480	wwPDB-VP
Average B, all atoms (Å ²)	14.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 27.07 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.3507e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, YRO, NCO, FMN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	1.34	12/2440 (0.5%)	1.14	7/3305 (0.2%)
1	B	1.34	5/2440 (0.2%)	1.10	5/3305 (0.2%)
All	All	1.34	17/4880 (0.3%)	1.12	12/6610 (0.2%)

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	126	LEU	C-O	-10.46	1.11	1.24
1	B	268	VAL	C-O	-9.04	1.14	1.24
1	B	163	VAL	C-O	-8.06	1.16	1.24
1	A	23	CYS	C-O	-7.87	1.18	1.23
1	A	130	CYS	N-CA	7.64	1.53	1.45
1	A	4	LEU	CA-C	6.38	1.60	1.52
1	B	131	PRO	N-CD	5.93	1.56	1.47
1	A	191	THR	C-O	5.82	1.31	1.24
1	B	127	ASN	C-O	-5.72	1.16	1.23
1	A	14	PRO	N-CA	-5.61	1.40	1.47
1	A	127	ASN	C-O	-5.48	1.17	1.24
1	A	130	CYS	CA-C	5.40	1.58	1.52
1	A	4	LEU	C-O	-5.20	1.17	1.23
1	B	131	PRO	C-O	-5.19	1.18	1.23
1	A	218	GLY	N-CA	5.18	1.51	1.45
1	A	163	VAL	C-O	-5.18	1.18	1.24
1	A	251	VAL	CA-CB	5.09	1.59	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	4	LEU	N-CA-C	8.07	121.16	109.14
1	A	278	GLY	CA-C-N	6.44	126.07	119.56

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	278	GLY	C-N-CA	6.44	126.07	119.56
1	A	3	LYS	CA-C-O	-6.32	114.86	121.55
1	B	130	CYS	CA-C-N	-6.25	113.52	120.14
1	B	130	CYS	C-N-CA	-6.25	113.52	120.14
1	A	5	ASN	N-CA-C	5.92	120.08	107.70
1	B	197	GLY	N-CA-C	5.79	118.13	111.36
1	A	3	LYS	CA-C-N	5.73	131.53	122.36
1	A	3	LYS	C-N-CA	5.73	131.53	122.36
1	B	1	CYS	N-CA-CB	-5.21	102.28	110.77
1	B	225	ILE	N-CA-C	5.02	116.94	112.17

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2388	0	2391	26	0
1	B	2388	0	2391	16	0
2	A	31	0	19	1	0
2	B	31	0	19	1	0
3	A	7	0	0	0	0
4	A	17	0	15	13	0
4	B	17	0	15	3	0
5	A	18	0	24	4	0
5	B	24	0	32	0	0
6	A	287	0	0	3	0
6	B	272	0	0	5	0
All	All	5480	0	4906	52	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (52) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:138:GLN:HG2	6:A:775:HOH:O	1.73	0.88
1:A:132:ASN:HB2	6:A:605:HOH:O	1.84	0.75
1:B:59:MET:HE3	6:B:613:HOH:O	1.88	0.73
1:A:147:ARG:O	1:A:151:GLN:HG3	1.90	0.71
1:A:130:CYS:HA	4:A:403:YRO:H10	1.75	0.68
4:A:403:YRO:CAM	4:A:403:YRO:H12	2.24	0.68
1:A:130:CYS:C	4:A:403:YRO:H8	2.20	0.66
1:A:234:ASN:HD21	1:A:238:ARG:HE	1.44	0.66
1:A:130:CYS:CA	4:A:403:YRO:CAB	2.76	0.63
1:A:130:CYS:HA	4:A:403:YRO:CAB	2.29	0.63
1:A:82:TYR:OH	1:A:87:HIS:HD2	1.83	0.61
1:A:130:CYS:C	4:A:403:YRO:CAB	2.74	0.61
4:B:402:YRO:H15	4:B:402:YRO:CAM	2.32	0.59
4:A:403:YRO:H12	4:A:403:YRO:CAN	2.34	0.57
1:B:234:ASN:HD21	1:B:238:ARG:HE	1.49	0.57
4:A:403:YRO:H13	5:A:406:GOL:H11	1.86	0.56
1:B:146:MET:HE2	1:B:180:VAL:HG11	1.88	0.55
1:B:7:LEU:O	1:B:9:HIS:HD2	1.90	0.55
1:B:59:MET:CE	6:B:613:HOH:O	2.51	0.53
1:A:20:GLY:HA3	2:A:401:FMN:N5	2.23	0.53
1:B:304:PHE:HA	1:B:307:ARG:HG3	1.91	0.53
1:B:82:TYR:OH	1:B:87:HIS:HD2	1.93	0.52
1:B:20:GLY:HA3	2:B:401:FMN:N5	2.25	0.51
1:A:5:ASN:HD21	1:A:10:VAL:HG22	1.77	0.50
1:B:87:HIS:HE1	1:B:92:LYS:O	1.94	0.49
1:A:160:PRO:HB3	1:A:188:LYS:HG3	1.95	0.49
1:A:128:LEU:HG	1:A:163:VAL:HG13	1.97	0.46
1:A:98:ILE:O	1:A:98:ILE:HG13	2.16	0.46
4:A:403:YRO:H13	5:A:406:GOL:C1	2.46	0.46
1:A:71:LEU:N	1:A:72:PRO:CD	2.79	0.46
1:B:195:SER:HB3	6:B:758:HOH:O	2.15	0.46
4:A:403:YRO:CAM	4:A:403:YRO:CAC	2.92	0.45
1:A:87:HIS:HE1	1:A:92:LYS:O	2.00	0.45
1:B:97:SER:HA	1:B:125:GLU:O	2.17	0.44
1:A:234:ASN:ND2	1:A:238:ARG:HE	2.12	0.44
1:B:291:GLU:CD	6:B:608:HOH:O	2.61	0.43
1:B:205:GLU:CD	1:B:205:GLU:C	2.86	0.43
1:A:205:GLU:CD	1:A:205:GLU:C	2.86	0.43
4:A:403:YRO:CAC	5:A:406:GOL:C1	2.96	0.43
1:B:131:PRO:HD3	4:B:402:YRO:H13	1.99	0.43
1:B:160:PRO:HB3	1:B:188:LYS:HG3	2.00	0.43
1:A:128:LEU:CD1	1:A:163:VAL:HG11	2.49	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:2:LEU:HD13	1:A:13:ASN:HB2	2.00	0.42
1:A:97:SER:HA	1:A:125:GLU:O	2.20	0.42
5:A:406:GOL:H12	6:A:777:HOH:O	2.20	0.42
1:B:234:ASN:ND2	1:B:238:ARG:HE	2.17	0.42
1:A:136:LYS:HA	1:A:136:LYS:HE2	2.01	0.41
1:A:185:PRO:O	1:A:188:LYS:NZ	2.51	0.40
4:B:402:YRO:H14	6:B:771:HOH:O	2.21	0.40
1:A:226:LEU:HB3	1:A:227:PRO:HD3	2.02	0.40
4:A:403:YRO:CAL	4:A:403:YRO:H16	2.51	0.40
1:A:130:CYS:N	4:A:403:YRO:H9	2.36	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	311/313 (99%)	306 (98%)	5 (2%)	0	100	100
1	B	311/313 (99%)	300 (96%)	11 (4%)	0	100	100
All	All	622/626 (99%)	606 (97%)	16 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	258/258 (100%)	249 (96%)	9 (4%)	32	16
1	B	258/258 (100%)	254 (98%)	4 (2%)	55	44
All	All	516/516 (100%)	503 (98%)	13 (2%)	42	27

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	0	MET
1	A	1	CYS
1	A	2	LEU
1	A	98	ILE
1	A	133	VAL
1	A	138	GLN
1	A	262	LEU
1	A	295	ARG
1	A	312	GLU
1	B	4	LEU
1	B	133	VAL
1	B	307	ARG
1	B	312	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	ASN
1	A	9	HIS
1	A	87	HIS
1	A	138	GLN
1	A	234	ASN
1	A	275	GLN
1	B	9	HIS
1	B	87	HIS
1	B	138	GLN
1	B	152	GLN
1	B	234	ASN
1	B	275	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

12 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	FMN	B	401	-	33,33,33	1.43	7 (21%)	48,50,50	1.56	10 (20%)
2	FMN	A	401	-	33,33,33	1.45	4 (12%)	48,50,50	1.26	4 (8%)
5	GOL	A	406	-	5,5,5	0.48	0	5,5,5	0.87	0
5	GOL	A	405	-	5,5,5	0.58	0	5,5,5	0.82	0
4	YRO	B	402	-	17,17,17	1.59	3 (17%)	22,25,25	1.27	3 (13%)
5	GOL	B	403	-	5,5,5	0.46	0	5,5,5	0.65	0
4	YRO	A	403	-	17,17,17	1.87	5 (29%)	22,25,25	1.79	6 (27%)
5	GOL	B	406	-	5,5,5	0.88	0	5,5,5	0.71	0
5	GOL	B	405	-	5,5,5	0.69	0	5,5,5	0.61	0
3	NCO	A	402	-	6,6,6	2.21	4 (66%)	-		
5	GOL	B	404	-	5,5,5	0.60	0	5,5,5	0.92	0
5	GOL	A	404	-	5,5,5	0.75	0	5,5,5	1.76	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	B	401	-	-	1/18/18/18	0/3/3/3

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FMN	A	401	-	-	2/18/18/18	0/3/3/3
5	GOL	A	406	-	-	2/4/4/4	-
5	GOL	A	405	-	-	0/4/4/4	-
5	GOL	B	403	-	-	0/4/4/4	-
4	YRO	A	403	-	-	1/10/10/10	0/1/1/1
5	GOL	B	406	-	-	2/4/4/4	-
5	GOL	B	405	-	-	0/4/4/4	-
4	YRO	B	402	-	-	4/10/10/10	0/1/1/1
5	GOL	B	404	-	-	2/4/4/4	-
5	GOL	A	404	-	-	1/4/4/4	-

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	FMN	C9A-C5A	4.11	1.47	1.41
2	A	401	FMN	C4-N3	-4.01	1.31	1.38
4	A	403	YRO	CAN-NAJ	3.68	1.46	1.38
3	A	402	NCO	CO-N2	3.60	2.08	1.96
4	A	403	YRO	CAO-NAK	3.52	1.43	1.37
4	B	402	YRO	CAH-CAM	-3.17	1.43	1.51
4	B	402	YRO	CAN-CAL	-2.91	1.43	1.48
2	B	401	FMN	C9A-C5A	2.82	1.45	1.41
4	A	403	YRO	CAB-CAQ	-2.65	1.43	1.52
4	A	403	YRO	CAN-CAM	2.57	1.42	1.37
2	B	401	FMN	C4-N3	-2.53	1.34	1.38
2	B	401	FMN	C8-C7	2.52	1.47	1.40
2	B	401	FMN	C4A-N5	2.41	1.36	1.30
2	B	401	FMN	C6-C7	-2.34	1.36	1.39
3	A	402	NCO	CO-N1	2.33	2.04	1.96
2	B	401	FMN	C1'-C2'	-2.32	1.49	1.52
3	A	402	NCO	CO-N6	2.16	2.03	1.96
3	A	402	NCO	CO-N4	2.15	2.03	1.96
2	A	401	FMN	C9A-N10	-2.12	1.37	1.41
2	A	401	FMN	O4-C4	2.11	1.27	1.23
4	B	402	YRO	CAN-NAJ	2.10	1.42	1.38
4	A	403	YRO	CAN-CAL	-2.09	1.45	1.48
2	B	401	FMN	O4-C4	2.09	1.27	1.23

All (25) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	403	YRO	CAM-CAP-NAK	4.77	119.33	115.70
2	B	401	FMN	O2-C2-N1	-4.06	115.06	121.80
2	A	401	FMN	O4-C4-C4A	-3.46	117.41	126.53
2	B	401	FMN	O4-C4-C4A	-3.18	118.13	126.53
2	B	401	FMN	O5'-C5'-C4'	3.07	117.55	109.36
2	B	401	FMN	O4-C4-N3	3.05	125.85	120.11
2	B	401	FMN	C4-C4A-N5	3.03	122.40	118.21
2	B	401	FMN	C10-C4A-N5	-2.88	118.92	124.81
4	A	403	YRO	OAF-CAP-NAK	-2.84	114.77	120.11
2	A	401	FMN	O4-C4-N3	2.72	125.24	120.11
2	A	401	FMN	C9A-C5A-N5	-2.69	119.59	122.45
5	A	404	GOL	O2-C2-C1	-2.68	98.07	109.18
2	A	401	FMN	C6-C5A-N5	2.50	122.59	118.44
4	A	403	YRO	CAH-CAM-CAP	2.50	119.99	116.64
4	A	403	YRO	OAD-CAL-CAN	2.39	120.53	116.73
4	A	403	YRO	CAA-CAQ-CAI	2.39	119.76	109.83
4	B	402	YRO	OAG-CAL-CAN	2.23	124.08	120.20
4	B	402	YRO	CAH-CAM-CAP	2.22	119.61	116.64
4	B	402	YRO	OAD-CAL-OAG	-2.18	118.71	123.90
2	B	401	FMN	C5A-N5-C4A	2.17	121.59	118.09
2	B	401	FMN	C5'-C4'-C3'	-2.16	108.14	112.22
2	B	401	FMN	O2-C2-N3	2.08	122.58	118.58
4	A	403	YRO	CAP-NAK-CAO	-2.08	123.51	126.37
5	A	404	GOL	O3-C3-C2	2.07	119.70	110.38
2	B	401	FMN	C4A-C10-N10	2.03	119.38	116.48

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	403	YRO	OAD-CAL-CAN-NAJ
4	B	402	YRO	OAD-CAL-CAN-NAJ
5	B	404	GOL	O1-C1-C2-C3
5	A	406	GOL	O1-C1-C2-C3
5	A	404	GOL	O2-C2-C3-O3
5	B	404	GOL	O1-C1-C2-O2
5	A	406	GOL	O1-C1-C2-O2
5	B	406	GOL	O2-C2-C3-O3
2	A	401	FMN	C2'-C3'-C4'-O4'
4	B	402	YRO	OAD-CAL-CAN-CAM
2	A	401	FMN	C4'-C5'-O5'-P
4	B	402	YRO	OAG-CAL-CAN-NAJ
2	B	401	FMN	C4'-C5'-O5'-P

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
5	B	406	GOL	C1-C2-C3-O3
4	B	402	YRO	OAG-CAL-CAN-CAM


There are no ring outliers.

5 monomers are involved in 19 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	401	FMN	1	0
2	A	401	FMN	1	0
5	A	406	GOL	4	0
4	B	402	YRO	3	0
4	A	403	YRO	13	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	313/313 (100%)	-0.25	11 (3%) 47 51	6, 12, 30, 44	0
1	B	313/313 (100%)	-0.42	6 (1%) 66 70	6, 11, 26, 41	0
All	All	626/626 (100%)	-0.34	17 (2%) 56 59	6, 12, 27, 44	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	131	PRO	4.2
1	A	135	GLY	4.0
1	A	2	LEU	3.7
1	B	131	PRO	3.4
1	A	133	VAL	3.4
1	A	130	CYS	3.3
1	A	134	PRO	3.2
1	A	5	ASN	3.1
1	B	132	ASN	3.0
1	A	312	GLU	2.5
1	B	312	GLU	2.3
1	A	136	LYS	2.3
1	B	0	MET	2.2
1	B	144	GLU	2.2
1	B	130	CYS	2.2
1	A	0	MET	2.2
1	A	295	ARG	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	GOL	B	404	6/6	0.82	0.14	37,47,48,53	0
5	GOL	B	406	6/6	0.89	0.13	18,23,28,30	0
5	GOL	A	404	6/6	0.92	0.09	17,19,20,21	0
5	GOL	B	405	6/6	0.92	0.10	21,25,25,26	0
5	GOL	A	405	6/6	0.92	0.09	20,21,24,26	0
5	GOL	A	406	6/6	0.93	0.10	16,16,21,25	0
4	YRO	B	402	17/17	0.96	0.08	8,10,33,36	0
2	FMN	B	401	31/31	0.97	0.05	5,6,7,9	0
4	YRO	A	403	17/17	0.97	0.07	9,11,29,29	0
5	GOL	B	403	6/6	0.97	0.05	16,18,19,21	0
2	FMN	A	401	31/31	0.98	0.05	5,7,7,8	0
3	NCO	A	402	7/7	0.99	0.06	9,9,9,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around FMN B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around FMN A 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.