



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:28 AM UTC

PDB ID : 5W1Z / pdb_00005w1z
Title : Crystal Structure of inosine-substituted decamer duplex DNA (I4)
Authors : Pallan, P.S.; Egli, M.
Deposited on : 2017-06-05
Resolution : 1.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2145 (1.56-1.56)
Clashscore	190562	2189 (1.56-1.56)
RSRZ outliers	180081	2146 (1.56-1.56)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	10	 60% 40%
1	B	10	 70% 20% 10%
1	C	10	 70% 30%
1	D	10	 80% 10% 10%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 966 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (5'-D(*CP*CP*AP*IP*IP*CP*CP*(BRU)P*IP*I)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	10	Total	Br	C	N	O	P	0	0	0
			198	1	95	35	58	9			
1	B	10	Total	Br	C	N	O	P	0	0	0
			198	1	95	35	58	9			
1	C	10	Total	Br	C	N	O	P	0	0	0
			198	1	95	35	58	9			
1	D	10	Total	Br	C	N	O	P	0	1	0
			209	1	100	35	63	10			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	C	1	Total	Mg	0	0
			1	1		
2	D	1	Total	Mg	0	0
			1	1		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		
3	C	1	Total	Na	0	0
			1	1		
3	D	2	Total	Na	0	0
			2	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	37	Total 37	O 37	0	0
4	B	40	Total 40	O 40	0	0
4	C	37	Total 37	O 37	0	0
4	D	42	Total 42	O 42	0	0

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (5'-D(*CP*CP*AP*IP*IP*CP*CP*(BRU)P*IP*I)-3')

Chain A: 




- Molecule 1: DNA (5'-D(*CP*CP*AP*IP*IP*CP*CP*(BRU)P*IP*I)-3')

Chain B: 




- Molecule 1: DNA (5'-D(*CP*CP*AP*IP*IP*CP*CP*(BRU)P*IP*I)-3')

Chain C: 



- Molecule 1: DNA (5'-D(*CP*CP*AP*IP*IP*CP*CP*(BRU)P*IP*I)-3')

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	24.88Å 32.80Å 34.03Å 90.25° 107.32° 111.04°	Depositor
Resolution (Å)	32.24 – 1.55 32.24 – 1.55	Depositor EDS
% Data completeness (in resolution range)	94.8 (32.24-1.55) 97.1 (32.24-1.55)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.33 (at 1.53Å)	Xtriage
Refinement program	REFMAC 5.8.0158	Depositor
R, R_{free}	0.207 , 0.234 0.208 , 0.234	Depositor DCC
R_{free} test set	805 reflections (5.61%)	wwPDB-VP
Wilson B-factor (Å ²)	17.2	Xtriage
Anisotropy	0.506	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 31.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.51$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	966	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 21.84 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 6.4533e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BRU, MG, NA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.63	0/103	1.12	1/153 (0.7%)
1	B	0.64	0/103	0.99	1/153 (0.7%)
1	C	0.51	0/103	1.13	0/153
1	D	0.57	0/124	1.06	1/185 (0.5%)
All	All	0.59	0/433	1.07	3/644 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1
1	B	0	1
All	All	0	2

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	13	DA	C2'-C3'-O3'	-7.49	100.26	111.50
1	B	17	DC	C2'-C3'-O3'	5.69	120.04	111.50
1	A	7	DC	C2'-C3'-O3'	5.45	119.68	111.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	9	DI	Sidechain

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Group
1	B	19	DI	Sidechain

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	198	0	106	1	0
1	B	198	0	106	1	0
1	C	198	0	106	1	0
1	D	209	0	109	1	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0
3	C	1	0	0	0	0
3	D	2	0	0	0	0
4	A	37	0	0	0	0
4	B	40	0	0	0	0
4	C	37	0	0	0	0
4	D	42	0	0	1	0
All	All	966	0	427	3	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (3) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:18:BRU:H2'	4:D:234:HOH:O	2.01	0.59
1:C:2:DC:H2''	1:C:3:DA:N7	2.25	0.52
1:A:3:DA:C2	1:B:19:DI:C2	3.06	0.44

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

There are no protein molecules in this entry.

5.3.2 Protein sidechains [i](#)

There are no protein molecules in this entry.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	BRU	C	8	1	18,21,22	1.58	4 (22%)	25,30,33	1.89	8 (32%)
1	BRU	D	18	1,3	18,21,22	1.96	6 (33%)	25,30,33	2.32	10 (40%)
1	BRU	A	8	1	18,21,22	1.93	6 (33%)	25,30,33	2.91	10 (40%)
1	BRU	B	18	1	18,21,22	2.13	4 (22%)	25,30,33	2.89	11 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BRU	C	8	1	-	4/7/21/22	0/2/2/2
1	BRU	D	18	1,3	-	2/7/21/22	0/2/2/2
1	BRU	A	8	1	-	4/7/21/22	0/2/2/2

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	BRU	B	18	1	-	1/7/21/22	0/2/2/2

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	8	BRU	C4-N3	-5.17	1.29	1.38
1	B	18	BRU	C6-N1	-4.25	1.30	1.38
1	B	18	BRU	C4-N3	-4.12	1.31	1.38
1	D	18	BRU	C4-N3	-3.99	1.31	1.38
1	B	18	BRU	C2-N1	3.89	1.44	1.38
1	B	18	BRU	C2-N3	-3.48	1.31	1.38
1	D	18	BRU	C6-N1	-3.48	1.32	1.38
1	C	8	BRU	C4-N3	-3.40	1.32	1.38
1	C	8	BRU	C6-N1	-3.30	1.32	1.38
1	A	8	BRU	C6-N1	-3.03	1.32	1.38
1	D	18	BRU	C2-N1	2.99	1.43	1.38
1	C	8	BRU	C2-N3	-2.83	1.33	1.38
1	D	18	BRU	C2-N3	-2.73	1.33	1.38
1	A	8	BRU	C2-N1	2.73	1.42	1.38
1	A	8	BRU	C1'-N1	-2.67	1.41	1.48
1	A	8	BRU	C2-N3	-2.66	1.33	1.38
1	A	8	BRU	BR-C5	-2.33	1.83	1.88
1	D	18	BRU	BR-C5	-2.25	1.83	1.88
1	D	18	BRU	C1'-N1	-2.20	1.42	1.48
1	C	8	BRU	C6-C5	2.00	1.38	1.34

All (39) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	8	BRU	N3-C2-N1	7.36	124.47	114.89
1	A	8	BRU	O2-C2-N1	-6.64	114.15	122.80
1	B	18	BRU	N3-C2-N1	6.63	123.52	114.89
1	A	8	BRU	C6-N1-C2	-5.96	115.37	121.30
1	B	18	BRU	O4-C4-C5	-5.78	118.43	125.80
1	B	18	BRU	C6-N1-C2	-4.99	116.34	121.30
1	B	18	BRU	O2-C2-N1	-4.82	116.52	122.80
1	D	18	BRU	O2-C2-N1	-4.71	116.67	122.80
1	D	18	BRU	O4-C4-C5	-4.67	119.84	125.80
1	D	18	BRU	N3-C2-N1	4.47	120.71	114.89
1	A	8	BRU	O4-C4-C5	-4.42	120.16	125.80
1	B	18	BRU	C4-N3-C2	-4.07	122.00	127.34
1	A	8	BRU	C4-N3-C2	-4.04	122.04	127.34

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	18	BRU	BR-C5-C4	-3.55	113.93	118.02
1	D	18	BRU	C4-N3-C2	-3.42	122.85	127.34
1	C	8	BRU	C6-C5-C4	-3.38	117.24	120.67
1	C	8	BRU	O4-C4-C5	-3.36	121.52	125.80
1	C	8	BRU	C5-C4-N3	3.31	117.15	113.34
1	C	8	BRU	C6-N1-C2	-3.28	118.04	121.30
1	D	18	BRU	C6-N1-C2	-3.20	118.12	121.30
1	C	8	BRU	N3-C2-N1	3.19	119.04	114.89
1	B	18	BRU	O4-C4-N3	3.08	125.91	120.11
1	D	18	BRU	C5-C4-N3	2.94	116.72	113.34
1	D	18	BRU	C1'-N1-C6	2.88	125.57	120.74
1	B	18	BRU	BR-C5-C6	2.78	124.53	120.64
1	D	18	BRU	O4'-C1'-C2'	-2.74	101.13	106.25
1	A	8	BRU	C1'-N1-C6	2.68	125.24	120.74
1	C	8	BRU	O2-C2-N1	-2.64	119.36	122.80
1	A	8	BRU	O4-C4-N3	2.64	125.07	120.11
1	D	18	BRU	O4'-C1'-N1	2.54	112.37	107.86
1	B	18	BRU	O3'-C3'-C2'	-2.51	102.16	110.88
1	D	18	BRU	O3'-C3'-C2'	-2.46	102.33	110.88
1	A	8	BRU	O3'-C3'-C2'	-2.44	102.39	110.88
1	C	8	BRU	BR-C5-C6	2.39	123.99	120.64
1	A	8	BRU	O4'-C1'-C2'	-2.34	101.88	106.25
1	A	8	BRU	C6-C5-C4	2.34	123.05	120.67
1	B	18	BRU	O4'-C4'-C5'	2.13	116.17	109.33
1	C	8	BRU	C1'-N1-C6	2.09	124.26	120.74
1	B	18	BRU	C2'-C3'-C4'	2.04	106.94	102.80

There are no chirality outliers.

All (11) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	8	BRU	C2'-C1'-N1-C6
1	A	8	BRU	C2'-C1'-N1-C2
1	C	8	BRU	C2'-C1'-N1-C2
1	D	18	BRU	C2'-C1'-N1-C2
1	C	8	BRU	C2'-C1'-N1-C6
1	C	8	BRU	O4'-C1'-N1-C2
1	A	8	BRU	O4'-C1'-N1-C6
1	C	8	BRU	O4'-C1'-N1-C6
1	B	18	BRU	C2'-C1'-N1-C2
1	A	8	BRU	O4'-C1'-N1-C2
1	D	18	BRU	O4'-C1'-N1-C2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	D	18	BRU	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 7 ligands modelled in this entry, 7 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	5/10 (50%)	0.00	0 100 100	20, 21, 23, 24	0
1	B	5/10 (50%)	-0.14	0 100 100	19, 20, 23, 25	0
1	C	5/10 (50%)	-0.02	0 100 100	19, 20, 27, 28	0
1	D	5/10 (50%)	0.01	0 100 100	15, 19, 24, 24	1 (20%)
All	All	20/40 (50%)	-0.04	0 100 100	15, 21, 27, 28	1 (5%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	BRU	B	18	20/21	0.91	0.09	17,23,35,49	1
1	BRU	C	8	20/21	0.92	0.09	18,23,28,54	1
1	BRU	A	8	20/21	0.93	0.08	17,24,33,48	1
1	BRU	D	18	20/21	0.93	0.08	18,23,30,44	1

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	NA	C	102	1/1	0.88	0.16	32,32,32,32	1
2	MG	A	101	1/1	0.91	0.14	27,27,27,27	1
3	NA	D	102	1/1	0.93	0.07	33,33,33,33	0
2	MG	D	101	1/1	0.94	0.13	33,33,33,33	0
3	NA	D	103	1/1	0.94	0.08	31,31,31,31	0
2	MG	C	101	1/1	0.96	0.10	22,22,22,22	0
3	NA	A	102	1/1	0.97	0.10	30,30,30,30	0

6.5 Other polymers [i](#)

There are no such residues in this entry.