



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 02:53 PM UTC

PDB ID : 2W20 / pdb_00002w20
Title : Structure of the catalytic domain of the native NanA sialidase from *Streptococcus pneumoniae*
Authors : Xu, G.; Andrew, P.W.; Taylor, G.L.
Deposited on : 2008-10-21
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	4037 (1.50-1.50)
Clashscore	190562	4235 (1.50-1.50)
Ramachandran outliers	187476	4153 (1.50-1.50)
Sidechain outliers	187428	4150 (1.50-1.50)
RSRZ outliers	180081	4039 (1.50-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	471	4% 93% 7%
1	B	471	10% 93% 6%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	GOL	A	1794	-	-	X	-

2 Entry composition [i](#)


There are 5 unique types of molecules in this entry. The entry contains 8499 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SIALIDASE A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	470	Total	C	N	O	S	0	0	0
			3717	2329	656	720	12			
1	B	469	Total	C	N	O	S	0	0	0
			3712	2326	655	719	12			

- Molecule 2 is 2-(N-MORPHOLINO)-ETHANESULFONIC ACID (CCD ID: MES) (formula: $C_6H_{13}NO_4S$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
2	A	1	Total	C	N	O	S	0	0
			12	6	1	4	1		
2	B	1	Total	C	N	O	S	0	0
			12	6	1	4	1		

- Molecule 3 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	2	Total Cl 2 2	0	0
3	B	1	Total Cl 1 1	0	0

- Molecule 4 is GLYCEROL (CCD ID: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C O 6 3 3	0	0
4	A	1	Total C O 6 3 3	0	0
4	B	1	Total C O 6 3 3	0	0


- Molecule 5 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	560	Total O 560 560	0	0
5	B	465	Total O 465 465	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: SIALIDASE A



• Molecule 1: SIALIDASE A



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	47.20Å 96.60Å 218.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.23 – 1.49 40.23 – 1.49	Depositor EDS
% Data completeness (in resolution range)	97.7 (40.23-1.49) 97.6 (40.23-1.49)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.25 (at 1.49Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE)	Depositor
R, R_{free}	0.200 , 0.225 0.194 , 0.220	Depositor DCC
R_{free} test set	8003 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	14.9	Xtriage
Anisotropy	0.298	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 37.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	8499	wwPDB-VP
Average B, all atoms (Å ²)	18.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.71% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, CL, MES

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/3798	0.70	0/5131
1	B	0.36	0/3793	0.70	0/5124
All	All	0.37	0/7591	0.70	0/10255

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3717	0	3633	30	0
1	B	3712	0	3628	17	0
2	A	12	0	12	4	0
2	B	12	0	12	5	0
3	A	2	0	0	1	0
3	B	1	0	0	0	0
4	A	12	0	16	6	0
4	B	6	0	8	1	0
5	A	560	0	0	1	0
5	B	465	0	0	0	0
All	All	8499	0	7309	50	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:734:ASN:H	1:A:735:GLY:HA2	1.17	1.02
1:A:783:LYS:HE3	4:A:1794:GOL:O1	1.84	0.76
1:B:769:HIS:NE2	1:B:771:GLU:HG3	2.02	0.73
2:A:1791:MES:H31	1:B:632:LYS:HE3	1.70	0.73
1:A:734:ASN:N	1:A:735:GLY:HA2	1.89	0.69
1:A:465:TYR:CD1	2:A:1791:MES:H22	2.28	0.69
3:A:1793:CL:CL	5:A:2403:HOH:O	2.51	0.64
1:B:626:ARG:CZ	1:B:685:ILE:HD12	2.31	0.60
1:A:734:ASN:HB2	1:A:736:GLU:H	1.65	0.60
1:A:386:ASN:HD22	4:A:1794:GOL:C1	2.17	0.57
1:B:465:TYR:CD1	2:B:1791:MES:H51	2.39	0.57
1:A:729:ALA:HB2	1:A:739:TRP:CE3	2.40	0.57
1:A:783:LYS:HE3	4:A:1794:GOL:HO1	1.72	0.53
1:A:386:ASN:HD22	4:A:1794:GOL:H11	1.75	0.52
1:A:510:LYS:O	1:A:513:GLN:HG2	2.10	0.51
1:B:729:ALA:HB2	1:B:739:TRP:CE3	2.46	0.51
1:A:499:LYS:HD3	1:A:504:ASP:HB3	1.93	0.50
1:B:348:ILE:HG21	1:B:417:ASP:HA	1.92	0.49
1:A:330:PHE:HB3	1:A:346:TYR:CG	2.48	0.49
1:A:348:ILE:HG21	1:A:417:ASP:HA	1.95	0.48
1:A:380:ILE:HD12	1:A:380:ILE:C	2.38	0.48
1:A:386:ASN:HB3	4:A:1794:GOL:H11	1.96	0.48
1:B:369:HIS:CE1	1:B:371:SER:HB2	2.49	0.47
1:A:632:LYS:HE2	2:B:1791:MES:H61	1.96	0.47
1:B:469:GLU:O	2:B:1791:MES:H72	2.15	0.47
2:B:1791:MES:H71	4:B:1793:GOL:H12	1.97	0.47
1:B:330:PHE:HB3	1:B:346:TYR:CG	2.50	0.47
1:A:734:ASN:HB2	1:A:736:GLU:HG3	1.98	0.46
1:A:386:ASN:HD22	4:A:1794:GOL:H12	1.80	0.46
1:A:465:TYR:CE1	2:A:1791:MES:H22	2.50	0.46
1:A:565:PHE:CZ	1:A:567:GLY:HA3	2.50	0.45
1:B:386:ASN:OD1	1:B:758:LEU:HG	2.17	0.45
1:A:606:ILE:C	1:A:606:ILE:HD12	2.42	0.45
1:A:734:ASN:N	1:A:735:GLY:CA	2.70	0.45
1:A:734:ASN:HB2	1:A:736:GLU:N	2.31	0.45
1:B:776:ALA:HA	1:B:777:TYR:HA	1.70	0.45
1:A:378:MET:HA	1:A:378:MET:HE2	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:606:ILE:C	1:B:606:ILE:HD12	2.43	0.44
2:A:1791:MES:H52	2:A:1791:MES:H82	1.61	0.43
1:A:419:VAL:HA	1:A:571:GLY:O	2.17	0.43
1:A:326:LYS:HG3	1:A:780:SER:HB3	2.00	0.43
1:B:732:GLU:C	1:B:733:GLU:HG3	2.43	0.42
1:B:733:GLU:CG	1:B:734:ASN:H	2.33	0.42
1:A:718:GLY:HA2	1:A:720:LYS:O	2.19	0.42
1:B:662:MET:HE1	1:B:685:ILE:HD13	2.02	0.42
1:B:565:PHE:CZ	1:B:567:GLY:HA3	2.55	0.41
1:A:725:MET:HE3	1:A:725:MET:HB3	1.96	0.41
1:A:585:ILE:O	1:A:608:TYR:HA	2.21	0.40
1:B:522:THR:O	1:B:523:ASN:C	2.65	0.40
1:A:632:LYS:HD2	2:B:1791:MES:H22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	468/471 (99%)	447 (96%)	21 (4%)	0	100	100
1	B	467/471 (99%)	446 (96%)	20 (4%)	1 (0%)	43	22
All	All	935/942 (99%)	893 (96%)	41 (4%)	1 (0%)	48	24

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	733	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	401/402 (100%)	398 (99%)	3 (1%)	76	57
1	B	401/402 (100%)	395 (98%)	6 (2%)	57	31
All	All	802/804 (100%)	793 (99%)	9 (1%)	65	41

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	404	LYS
1	A	499	LYS
1	A	645	ASN
1	B	404	LYS
1	B	477	GLU
1	B	536	LEU
1	B	645	ASN
1	B	733	GLU
1	B	758	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	386	ASN
1	A	523	ASN
1	B	478	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 3 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
4	GOL	A	1794	-	5,5,5	0.44	0	5,5,5	0.48	0
4	GOL	B	1793	-	5,5,5	0.38	0	5,5,5	0.34	0
2	MES	B	1791	-	12,12,12	2.22	1 (8%)	15,16,16	5.60	7 (46%)
2	MES	A	1791	-	12,12,12	2.13	1 (8%)	15,16,16	5.91	8 (53%)
4	GOL	A	1795	-	5,5,5	0.38	0	5,5,5	0.47	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	A	1794	-	-	2/4/4/4	-
4	GOL	B	1793	-	-	1/4/4/4	-
2	MES	B	1791	-	-	3/6/14/14	0/1/1/1
2	MES	A	1791	-	-	5/6/14/14	0/1/1/1
4	GOL	A	1795	-	-	4/4/4/4	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1791	MES	C8-S	-7.33	1.67	1.77
2	A	1791	MES	C8-S	-7.05	1.67	1.77

All (15) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	1791	MES	O1S-S-C8	-13.81	85.87	106.73
2	B	1791	MES	O1S-S-C8	-12.92	87.20	106.73
2	A	1791	MES	O2S-S-C8	-12.51	87.83	106.73
2	B	1791	MES	O2S-S-C8	-11.32	89.63	106.73
2	A	1791	MES	O3S-S-C8	-9.23	87.94	106.00
2	B	1791	MES	O3S-S-C8	-9.09	88.22	106.00
2	B	1791	MES	C5-N4-C3	7.03	123.99	108.84
2	A	1791	MES	C5-N4-C3	6.59	123.03	108.84
2	A	1791	MES	O3S-S-O1S	3.65	120.54	111.40
2	B	1791	MES	O3S-S-O1S	3.50	120.17	111.40
2	A	1791	MES	C7-N4-C5	3.41	120.33	111.24
2	B	1791	MES	C7-N4-C3	3.32	120.08	111.24
2	B	1791	MES	O3S-S-O2S	3.31	119.69	111.40
2	A	1791	MES	O3S-S-O2S	3.30	119.65	111.40
2	A	1791	MES	C7-N4-C3	2.00	116.58	111.24

There are no chirality outliers.

All (15) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	1791	MES	C8-C7-N4-C5
2	A	1791	MES	N4-C7-C8-S
2	B	1791	MES	C8-C7-N4-C3
4	A	1794	GOL	C1-C2-C3-O3
4	A	1795	GOL	O1-C1-C2-C3
4	A	1795	GOL	C1-C2-C3-O3
4	A	1794	GOL	O2-C2-C3-O3
4	A	1795	GOL	O2-C2-C3-O3
4	A	1795	GOL	O1-C1-C2-O2
4	B	1793	GOL	O1-C1-C2-O2
2	A	1791	MES	C7-C8-S-O2S
2	B	1791	MES	N4-C7-C8-S
2	B	1791	MES	C8-C7-N4-C5
2	A	1791	MES	C7-C8-S-O3S
2	A	1791	MES	C7-C8-S-O1S

There are no ring outliers.

4 monomers are involved in 15 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	1794	GOL	6	0
4	B	1793	GOL	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1791	MES	5	0
2	A	1791	MES	4	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	470/471 (99%)	0.11	18 (3%) 44 48	7, 14, 28, 47	0
1	B	469/471 (99%)	0.57	45 (9%) 13 14	9, 18, 33, 45	0
All	All	939/942 (99%)	0.34	63 (6%) 24 26	7, 16, 31, 47	0

All (63) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	734	ASN	7.0
1	B	734	ASN	5.7
1	B	458	GLY	4.7
1	B	733	GLU	4.3
1	A	733	GLU	4.2
1	A	457	ASP	3.9
1	B	790	SER	3.6
1	A	321	ALA	3.6
1	B	322	ALA	3.5
1	A	736	GLU	3.4
1	B	786	TRP	3.4
1	B	628	VAL	3.3
1	B	336	GLY	3.3
1	A	513	GLN	3.3
1	B	334	ARG	3.2
1	B	327	THR	3.2
1	A	485	ASP	3.2
1	A	790	SER	3.2
1	A	735	GLY	3.1
1	B	457	ASP	3.1
1	A	487	LYS	3.0
1	B	477	GLU	3.0
1	B	735	GLY	2.9
1	B	512	ASN	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	338	PRO	2.8
1	B	392	ASP	2.7
1	B	459	LYS	2.6
1	B	787	ASP	2.6
1	A	334	ARG	2.6
1	B	324	THR	2.6
1	A	467	GLU	2.5
1	B	631	GLN	2.5
1	B	343	ILE	2.5
1	B	629	ASP	2.5
1	B	425	GLU	2.5
1	B	719	PRO	2.5
1	A	401	ASP	2.5
1	B	630	GLY	2.4
1	B	739	TRP	2.4
1	B	393	ARG	2.4
1	B	773	GLY	2.4
1	B	706	GLU	2.4
1	A	511	GLY	2.3
1	B	340	LYS	2.3
1	B	747	LYS	2.3
1	B	323	LEU	2.3
1	B	685	ILE	2.3
1	B	333	GLY	2.2
1	B	771	GLU	2.2
1	B	683	LYS	2.2
1	B	708	LYS	2.2
1	A	514	LEU	2.1
1	A	512	ASN	2.1
1	B	627	GLN	2.1
1	B	485	ASP	2.1
1	A	477	GLU	2.1
1	B	342	GLY	2.1
1	B	760	ASN	2.1
1	B	389	THR	2.0
1	B	642	ARG	2.0
1	A	723	ASN	2.0
1	B	339	ASN	2.0
1	B	355	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	GOL	B	1793	6/6	0.72	0.17	27,29,34,40	0
4	GOL	A	1794	6/6	0.74	0.18	15,26,34,41	0
4	GOL	A	1795	6/6	0.76	0.14	24,31,34,35	0
3	CL	A	1793	1/1	0.89	0.31	35,35,35,35	0
2	MES	A	1791	12/12	0.89	0.13	20,22,25,26	8
2	MES	B	1791	12/12	0.91	0.10	16,18,24,24	6
3	CL	A	1792	1/1	0.97	0.07	14,14,14,14	0
3	CL	B	1792	1/1	0.99	0.04	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.