



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 11:34 AM UTC

PDB ID : 8W2C / pdb_00008w2c
Title : Thioesterase domain structure from Sulfazecin biosynthetic nonribosomal peptide synthetase SulM
Authors : Patel, K.D.; Gulick, A.M.
Deposited on : 2024-02-20
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7789 (1.90-1.90)
Clashscore	190562	8410 (1.90-1.90)
Ramachandran outliers	187476	8333 (1.90-1.90)
Sidechain outliers	187428	8333 (1.90-1.90)
RSRZ outliers	180081	7790 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	282	3% 85% 10%
1	B	282	3% 84% 6% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	EDO	A	3003	-	-	-	X
2	EDO	A	3004	-	-	-	X

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7927 atoms, of which 3789 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Non-ribosomal peptide synthetase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	253	Total	C	H	N	O	S	0	1	0
			3756	1197	1857	329	358	15			
1	B	256	Total	C	H	N	O	S	0	1	0
			3817	1214	1890	334	364	15			

There are 40 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	2703	GLY	-	expression tag	UNP A0A1I9RH13
A	2704	SER	-	expression tag	UNP A0A1I9RH13
A	2705	SER	-	expression tag	UNP A0A1I9RH13
A	2706	HIS	-	expression tag	UNP A0A1I9RH13
A	2707	HIS	-	expression tag	UNP A0A1I9RH13
A	2708	HIS	-	expression tag	UNP A0A1I9RH13
A	2709	HIS	-	expression tag	UNP A0A1I9RH13
A	2710	HIS	-	expression tag	UNP A0A1I9RH13
A	2711	HIS	-	expression tag	UNP A0A1I9RH13
A	2712	SER	-	expression tag	UNP A0A1I9RH13
A	2713	SER	-	expression tag	UNP A0A1I9RH13
A	2714	GLY	-	expression tag	UNP A0A1I9RH13
A	2715	LEU	-	expression tag	UNP A0A1I9RH13
A	2716	VAL	-	expression tag	UNP A0A1I9RH13
A	2717	PRO	-	expression tag	UNP A0A1I9RH13
A	2718	ARG	-	expression tag	UNP A0A1I9RH13
A	2719	GLY	-	expression tag	UNP A0A1I9RH13
A	2720	SER	-	expression tag	UNP A0A1I9RH13
A	2721	HIS	-	expression tag	UNP A0A1I9RH13
A	2722	MET	-	expression tag	UNP A0A1I9RH13
B	2703	GLY	-	expression tag	UNP A0A1I9RH13
B	2704	SER	-	expression tag	UNP A0A1I9RH13
B	2705	SER	-	expression tag	UNP A0A1I9RH13
B	2706	HIS	-	expression tag	UNP A0A1I9RH13
B	2707	HIS	-	expression tag	UNP A0A1I9RH13

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	2708	HIS	-	expression tag	UNP A0A1I9RH13
B	2709	HIS	-	expression tag	UNP A0A1I9RH13
B	2710	HIS	-	expression tag	UNP A0A1I9RH13
B	2711	HIS	-	expression tag	UNP A0A1I9RH13
B	2712	SER	-	expression tag	UNP A0A1I9RH13
B	2713	SER	-	expression tag	UNP A0A1I9RH13
B	2714	GLY	-	expression tag	UNP A0A1I9RH13
B	2715	LEU	-	expression tag	UNP A0A1I9RH13
B	2716	VAL	-	expression tag	UNP A0A1I9RH13
B	2717	PRO	-	expression tag	UNP A0A1I9RH13
B	2718	ARG	-	expression tag	UNP A0A1I9RH13
B	2719	GLY	-	expression tag	UNP A0A1I9RH13
B	2720	SER	-	expression tag	UNP A0A1I9RH13
B	2721	HIS	-	expression tag	UNP A0A1I9RH13
B	2722	MET	-	expression tag	UNP A0A1I9RH13

- Molecule 2 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	A	1	Total	C	H	O	0	0
			10	2	6	2		
2	B	1	Total	C	H	O	0	0
			10	2	6	2		

- Molecule 3 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Na	0	0
			1	1		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	133	Total	O	0	0
			133	133		
4	B	150	Total	O	0	0
			150	150		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Non-ribosomal peptide synthetase

Chain A: 



- Molecule 1: Non-ribosomal peptide synthetase

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	47.10Å 83.19Å 69.38Å 90.00° 101.36° 90.00°	Depositor
Resolution (Å)	42.27 – 1.90 42.27 – 1.90	Depositor EDS
% Data completeness (in resolution range)	95.5 (42.27-1.90) 87.6 (42.27-1.90)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	0.52 (at 1.90Å)	Xtriage
Refinement program	PHENIX 1.13_2998	Depositor
R, R_{free}	0.175 , 0.215 0.177 , 0.215	Depositor DCC
R_{free} test set	1995 reflections (4.82%)	wwPDB-VP
Wilson B-factor (Å ²)	25.3	Xtriage
Anisotropy	0.493	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.42 , 47.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7927	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.58% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NA, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.46	0/1940	0.54	0/2647
1	B	0.50	0/1969	0.56	0/2685
All	All	0.48	0/3909	0.55	0/5332

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1899	1857	1856	7	0
1	B	1927	1890	1890	13	0
2	A	24	36	36	0	0
2	B	4	6	5	0	0
3	A	1	0	0	0	0
4	A	133	0	0	1	0
4	B	150	0	0	0	0
All	All	4138	3789	3787	20	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:2818[A]:CYS:SG	1:B:2819:ASN:N	2.64	0.70
1:A:2816:ALA:HB2	1:A:2824:ALA:HB2	1.85	0.57
1:B:2816:ALA:HB2	1:B:2824:ALA:HB2	1.89	0.53
1:A:2837:ASP:OD2	4:A:3101:HOH:O	2.19	0.52
1:B:2875:PRO:HG2	1:B:2881:GLN:HA	1.92	0.51
1:B:2749:ASN:CB	1:B:2818[B]:CYS:SG	3.00	0.50
1:B:2940:SER:HA	1:B:2943:GLY:O	2.14	0.48
1:B:2757:PHE:HB3	1:B:2773:MET:CE	2.46	0.46
1:B:2757:PHE:HB3	1:B:2773:MET:HE1	1.99	0.45
1:A:2940:SER:HA	1:A:2943:GLY:O	2.18	0.44
1:A:2818[A]:CYS:SG	1:A:2819:ASN:N	2.92	0.43
1:A:2749:ASN:CB	1:A:2818[B]:CYS:SG	3.07	0.42
1:B:2894:LEU:HD23	1:B:2894:LEU:C	2.45	0.42
1:B:2932:TYR:N	1:B:2933:PRO:CD	2.83	0.41
1:B:2872:SER:HB2	1:B:2957:TYR:OH	2.20	0.41
1:B:2746:CYS:HB3	1:B:2757:PHE:CZ	2.56	0.41
1:B:2749:ASN:HB2	1:B:2818[B]:CYS:SG	2.61	0.41
1:A:2928:LEU:HD12	1:A:2956:HIS:CD2	2.56	0.41
1:B:2978:SER:O	1:B:2979:CYS:HB2	2.21	0.40
1:A:2757:PHE:HB3	1:A:2773:MET:CE	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/282 (89%)	248 (98%)	4 (2%)	0	100	100
1	B	255/282 (90%)	251 (98%)	4 (2%)	0	100	100
All	All	507/564 (90%)	499 (98%)	8 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/226 (88%)	199 (100%)	0	100	100
1	B	203/226 (90%)	201 (99%)	2 (1%)	68	70
All	All	402/452 (89%)	400 (100%)	2 (0%)	86	84

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	2818[A]	CYS
1	B	2818[B]	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	2901	HIS
1	A	2937	GLN
1	B	2976	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 8 ligands modelled in this entry, 1 is monoatomic - leaving 7 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	3003	-	3,3,3	0.44	0	2,2,2	0.40	0
2	EDO	A	3005	-	3,3,3	0.39	0	2,2,2	0.45	0
2	EDO	A	3004	-	3,3,3	0.40	0	2,2,2	0.47	0
2	EDO	A	3006	-	3,3,3	0.38	0	2,2,2	0.54	0
2	EDO	B	3001	-	3,3,3	0.87	0	2,2,2	0.78	0
2	EDO	A	3001	-	3,3,3	0.39	0	2,2,2	0.33	0
2	EDO	A	3002	-	3,3,3	0.43	0	2,2,2	0.21	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	3003	-	-	0/1/1/1	-
2	EDO	A	3005	-	-	1/1/1/1	-
2	EDO	A	3004	-	-	0/1/1/1	-
2	EDO	A	3006	-	-	0/1/1/1	-
2	EDO	B	3001	-	-	1/1/1/1	-
2	EDO	A	3001	-	-	0/1/1/1	-
2	EDO	A	3002	-	-	0/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	3005	EDO	O1-C1-C2-O2
2	B	3001	EDO	O1-C1-C2-O2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2			OWAB(Å ²)	Q < 0.9
1	A	253/282 (89%)	0.02	9 (3%)	46	49	17, 39, 74, 102	1 (0%)
1	B	256/282 (90%)	-0.15	8 (3%)	51	55	17, 34, 73, 97	1 (0%)
All	All	509/564 (90%)	-0.06	17 (3%)	49	52	17, 36, 74, 102	2 (0%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	2980	LYS	3.7
1	A	2878	PRO	3.4
1	A	2979	CYS	3.1
1	A	2876	ALA	2.8
1	B	2726	GLU	2.8
1	B	2877	GLN	2.7
1	B	2883	PHE	2.7
1	A	2877	GLN	2.6
1	A	2879	ASP	2.5
1	B	2981	GLY	2.4
1	A	2738	ALA	2.4
1	B	2794	ARG	2.4
1	B	2727	SER	2.4
1	A	2883	PHE	2.1
1	B	2784	ASP	2.1
1	A	2737	PRO	2.0
1	A	2784	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	EDO	A	3003	4/4	0.58	0.41	144,172,184,188	0
2	EDO	A	3004	4/4	0.71	0.43	132,158,167,167	0
2	EDO	A	3001	4/4	0.84	0.15	58,70,80,81	0
2	EDO	A	3002	4/4	0.89	0.11	31,37,37,42	0
2	EDO	A	3005	4/4	0.91	0.21	68,82,91,91	0
2	EDO	A	3006	4/4	0.91	0.16	49,59,70,70	0
2	EDO	B	3001	4/4	0.92	0.17	14,29,35,38	0
3	NA	A	3007	1/1	0.99	0.17	18,18,18,18	0

6.5 Other polymers [i](#)

There are no such residues in this entry.