



wwPDB EM Validation Summary Report ⓘ

Mar 9, 2026 – 01:05 PM UTC

PDB ID : 8W2R / pdb_00008w2r
EMDB ID : EMD-43756
Title : HIV-1 P5-IN intasome core
Authors : Li, M.; Craigie, R.
Deposited on : 2024-02-21
Resolution : 3.23 Å (reported)
Based on initial model : 5U1C

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev132
Mogul : 2022.3.0, CSD as543be (2022)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 3.23 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
Q-score	-	25397	14612 (2.73 - 3.73)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	362	
1	B	362	
1	C	362	
1	D	362	

Continued on next page...

Continued from previous page...


Mol	Chain	Length	Quality of chain
1	I	362	58%13%29%
1	K	362	57%10%33%
1	L	362	14%82%
1	M	362	10%87%
2	E	27	48%22%30%
2	N	27	52%19%30%
3	F	25	64%32%
3	O	25	64%32%

ENTRY-COMPOSITION INFOmissingINFO


2 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Integrase





• Molecule 1: Integrase



• Molecule 1: Integrase














● Molecule 1: Integrase

Chain K: 57% 10% 33%




● Molecule 1: Integrase

Chain L: 14% . 82%



● Molecule 1: Integrase

Chain M: 10% . 87%



GLY GLU ALA MET HIS GLY GLN VAL ASP CYS SER PRO GLY ILE TRP GLN LEU ASP CYS THR HIS LEU GLU GLY LYS VAL ILE VAL HIS VAL HIS LEU GLU PHE TRP THR VAL ARG

ARG TRP PRO VAL HIS THR VAL HIS THR ASP ASN GLY SER ASN PHE THR SER THR VAL LYS ALA GLY ALA TRP LYS TRP ALA GLY ILE VAL ALA VAL HIS VAL PHE GLY ILE SER PRO GLY TYR ASN PRO GLU ALA GLN GLY ILE VAL PRO ILE ALA THR THR GLY THR SER MET GLN ASN LYS GLU LEU LYS THR PHE ILE LEU ILE GLY VAL ARG

ASP GLN ALA HIS LEU LYS THR ALA VAL GLN MET ALA VAL PHE ILE THR HIS ASN PHE LYS ARG ASP TYR GLY LYS ILE GLY TYR SER ALA ASP GLY GLU ARG ILE VAL ASP ILE ILE THR ALA ILE ASP ILE THR LYS GLU LEU LYS GLN ILE THR LYS ILE ILE GLN N222 V225 D229

S230 R231 A239 L242 Q252 N254 S255 D256 R263 K266 I268 ARG ASP TYR GLY LYS MET GLY TYR GLY ASP ASP CYS VAL ALA SER ARG GLN ASP GLU ASP

- Molecule 2: DNA (5'-D(*AP*CP*TP*GP*CP*TP*AP*GP*AP*GP*AP*TP*TP*TP*TP*CP*CP*CP*G)-3')

Chain E:  48% 22% 30%

A15 C19 T20 A21 G22 A23 T29 C30 G33 DC DC DC DA DA DG DC DT

- Molecule 2: DNA (5'-D(*AP*CP*TP*GP*CP*TP*AP*GP*AP*GP*AP*TP*TP*TP*TP*CP*CP*CP*G)-3')

Chain N:  52% 19% 30%

A15 C19 T20 A23 T29 C30 G33 DC DC DC DA DA DG DC DT

- Molecule 3: DNA (5'-D(P*CP*GP*GP*GP*AP*AP*AP*AP*TP*CP*TP*CP*TP*AP*GP*CP*P*A)-3')

Chain F:  64% 1% 32%

DA DG DC DG DT DG DG DG C9 T17 A21

- Molecule 3: DNA (5'-D(P*CP*GP*GP*GP*AP*AP*AP*AP*TP*CP*TP*CP*TP*AP*GP*CP*P*A)-3')

Chain O:  64% 1% 32%

DA DG DC DG DT DG DG DG C5 T17 A21

3 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	154000	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	NONE	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	54	Depositor
Minimum defocus (nm)	1400	Depositor
Maximum defocus (nm)	2600	Depositor
Magnification	105000	Depositor
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.052	Depositor
Minimum map value	-0.012	Depositor
Average map value	0.000	Depositor
Map value standard deviation	0.001	Depositor
Recommended contour level	0.01	Depositor
Map size ($\text{\AA}$)	381.59998, 381.59998, 381.59998	wwPDB
Map dimensions	360, 360, 360	wwPDB
Map angles ($^\circ$)	90.0, 90.0, 90.0	wwPDB
Pixel spacing ($\text{\AA}$)	1.06, 1.06, 1.06	Depositor

4 Model quality [i](#)

4.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: DLU, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.17	0/2071	0.31	0/2796
1	B	0.13	0/1951	0.28	0/2631
1	C	0.15	0/544	0.27	0/729
1	D	0.18	0/397	0.30	0/535
1	I	0.16	0/2062	0.28	0/2785
1	K	0.15	0/1951	0.31	0/2631
1	L	0.15	0/544	0.28	0/729
1	M	0.18	0/397	0.30	0/535
2	E	0.24	0/429	0.42	0/660
2	N	0.24	0/429	0.42	0/660
3	F	0.22	0/393	0.38	0/604
3	O	0.22	0/393	0.38	0/604
All	All	0.17	0/11561	0.31	0/15899

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

4.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2030	0	2047	36	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	1914	0	1936	27	0
1	C	534	0	561	12	0
1	D	388	0	406	7	0
1	I	2021	0	2034	36	0
1	K	1914	0	1938	25	0
1	L	534	0	561	10	0
1	M	388	0	406	7	0
2	E	384	0	217	5	0
2	N	384	0	217	3	0
3	F	350	0	191	1	0
3	O	350	0	191	1	0
4	A	2	0	0	0	0
4	I	2	0	0	0	0
5	A	1	0	0	0	0
5	I	1	0	0	0	0
6	A	30	0	18	0	0
6	I	30	0	18	0	0
All	All	11257	0	10741	146	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

The worst 5 of 146 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:K:74:LEU:HD12	1:K:89:ILE:HD12	1.69	0.74
1:K:12:HIS:HE2	1:K:43:CYS:HG	1.35	0.73
1:D:266:LYS:NZ	2:N:23:DA:OP1	2.21	0.71
2:E:23:DA:OP1	1:M:266:LYS:NZ	2.24	0.69
1:B:242:LEU:HD11	1:B:252:GLN:HG2	1.75	0.68

There are no symmetry-related clashes.

4.3 Torsion angles [i](#)

4.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/362 (70%)	249 (98%)	6 (2%)	0	100	100
1	B	234/362 (65%)	230 (98%)	4 (2%)	0	100	100
1	C	62/362 (17%)	59 (95%)	3 (5%)	0	100	100
1	D	45/362 (12%)	42 (93%)	3 (7%)	0	100	100
1	I	254/362 (70%)	248 (98%)	6 (2%)	0	100	100
1	K	234/362 (65%)	225 (96%)	8 (3%)	1 (0%)	30	60
1	L	62/362 (17%)	60 (97%)	2 (3%)	0	100	100
1	M	45/362 (12%)	42 (93%)	3 (7%)	0	100	100
All	All	1191/2896 (41%)	1155 (97%)	35 (3%)	1 (0%)	49	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	K	3	ASP

4.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	215/293 (73%)	215 (100%)	0	100	100
1	B	202/293 (69%)	202 (100%)	0	100	100
1	C	57/293 (20%)	57 (100%)	0	100	100
1	D	41/293 (14%)	41 (100%)	0	100	100
1	I	214/293 (73%)	214 (100%)	0	100	100
1	K	202/293 (69%)	202 (100%)	0	100	100
1	L	57/293 (20%)	57 (100%)	0	100	100
1	M	41/293 (14%)	41 (100%)	0	100	100
All	All	1029/2344 (44%)	1029 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 7 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	221	GLN
1	C	274	GLN
1	L	216	GLN
1	K	164	GLN
1	A	252	GLN

4.3.3 RNA [i](#)

There are no RNA molecules in this entry.

4.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

4.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

4.6 Ligand geometry [i](#)

Of 8 ligands modelled in this entry, 6 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
6	DLU	I	304	4	30,33,33	2.00	7 (23%)	34,49,49	1.50	6 (17%)
6	DLU	A	304	4	30,33,33	1.99	7 (23%)	34,49,49	1.50	6 (17%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the

Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	DLU	I	304	4	-	0/9/35/35	0/4/4/4
6	DLU	A	304	4	-	0/9/35/35	0/4/4/4

The worst 5 of 14 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	I	304	DLU	CAW-CAZ	-4.67	1.38	1.50
6	A	304	DLU	CAW-CAZ	-4.66	1.38	1.50
6	I	304	DLU	FAF-CAT	-4.50	1.25	1.36
6	A	304	DLU	FAF-CAT	-4.47	1.25	1.36
6	I	304	DLU	FAG-CAU	-4.43	1.23	1.35

The worst 5 of 12 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	I	304	DLU	OAC-CAS-CAY	-3.71	118.78	126.38
6	A	304	DLU	OAC-CAS-CAY	-3.68	118.84	126.38
6	A	304	DLU	CAM-CBA-NBC	-3.49	106.08	109.42
6	I	304	DLU	CAM-CBA-NBC	-3.47	106.10	109.42
6	I	304	DLU	CAA-CBA-CAM	-2.48	108.13	112.74


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



4.7 Other polymers [i](#)

There are no such residues in this entry.

4.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


5 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-43756. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

5.1 Orthogonal projections [i](#)


5.1.1 Primary map



X




Y




Z


5.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

5.2 Central slices [i](#)


5.2.1 Primary map



X Index: 180




Y Index: 180




Z Index: 180


5.2.2 Raw map



X Index: 180



Y Index: 180




Z Index: 180


The images above show central slices of the map in three orthogonal directions.

5.3 Largest variance slices [i](#)


5.3.1 Primary map



X Index: 167



Y Index: 205




Z Index: 164


5.3.2 Raw map



X Index: 167



Y Index: 155



Z Index: 164


The images above show the largest variance slices of the map in three orthogonal directions.

5.4 Orthogonal standard-deviation projections (False-color) [i](#)


5.4.1 Primary map



X



Y




Z


5.4.2 Raw map



X



Y




Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


5.5 Orthogonal surface views [i](#)

5.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.01. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

5.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


5.6 Mask visualisation [i](#)

This section was not generated. No masks/segmentation were deposited.

6 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

6.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


6.2 Volume estimate [i](#)



The volume at the recommended contour level is 164 nm³; this corresponds to an approximate mass of 148 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

6.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.310 Å⁻¹

7 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

7.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of $0.310 \text{ \AA}^{-1}$

7.2 Resolution estimates [i](#)

Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	3.23	-	-
Author-provided FSC curve	3.25	3.59	3.27
Unmasked-calculated*	3.84	4.42	3.89

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.84 differs from the reported value 3.23 by more than 10 %


8 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-43756 and PDB model 8W2R. Per-residue inclusion information can be found in section 2 on page 4.

8.1 Map-model overlay [i](#)

This section was not generated.

8.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

8.3 Atom inclusion mapped to coordinate model [i](#)

This section was not generated.

8.4 Atom inclusion [i](#)



At the recommended contour level, 96% of all backbone atoms, 86% of all non-hydrogen atoms, are inside the map.

8.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.01) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8640	0.4320
A	0.8710	0.4500
B	0.8210	0.3920
C	0.8670	0.4620
D	0.8660	0.4920
E	0.9690	0.4430
F	0.8910	0.3800
I	0.8760	0.4610
K	0.8330	0.3930
L	0.8610	0.4670
M	0.8660	0.4940
N	0.9610	0.4410
O	0.8940	0.3840

