



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:56 PM UTC

PDB ID : 5W3K / pdb_00005w3k
Title : Crystal structure of Staphylococcus aureus ketol-acid reductoisomerase in complex NADPH, Mg²⁺ and CPD
Authors : Patel, K.M.; Teran, D.; Zheng, S.; Kandale, A.; Schembri, M.; McGeary, R.P.; Schenk, G.; Guddat, L.W.
Deposited on : 2017-06-08
Resolution : 1.59 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.59 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	340	% 92% 5% .
1	B	340	91% 5% .

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 6070 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Ketol-acid reductoisomerase (NADP(+)).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	326	Total	C	N	O	S	0	10	0
			2609	1658	444	499	8			
1	B	326	Total	C	N	O	S	0	13	0
			2622	1670	441	503	8			

There are 12 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	335	HIS	-	expression tag	UNP A0A145BYP4
A	336	HIS	-	expression tag	UNP A0A145BYP4
A	337	HIS	-	expression tag	UNP A0A145BYP4
A	338	HIS	-	expression tag	UNP A0A145BYP4
A	339	HIS	-	expression tag	UNP A0A145BYP4
A	340	HIS	-	expression tag	UNP A0A145BYP4
B	335	HIS	-	expression tag	UNP A0A145BYP4
B	336	HIS	-	expression tag	UNP A0A145BYP4
B	337	HIS	-	expression tag	UNP A0A145BYP4
B	338	HIS	-	expression tag	UNP A0A145BYP4
B	339	HIS	-	expression tag	UNP A0A145BYP4
B	340	HIS	-	expression tag	UNP A0A145BYP4

- Molecule 2 is cyclopropane-1,1-dicarboxylic acid (CCD ID: 9TY) (formula: C₅H₆O₄) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			9	5	4		
2	B	1	Total	C	O	0	0
			9	5	4		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	Mg	0	0
			4	4		
3	B	2	Total	Mg	0	0
			2	2		

- Molecule 4 is NADPH DIHYDRO-NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (CCD ID: NDP) (formula: C₂₁H₃₀N₇O₁₇P₃).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	A	1	Total	C	N	O	P	0	1
			96	42	14	34	6		
4	B	1	Total	C	N	O	P	0	0
			48	21	7	17	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	344	Total	O	0	0
			344	344		
5	B	327	Total	O	0	0
			327	327		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Ketol-acid reductoisomerase (NADP(+))

Chain A:  92%



- Molecule 1: Ketol-acid reductoisomerase (NADP(+))

Chain B:  91% 5%



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	63.94Å 80.77Å 66.75Å 90.00° 92.49° 90.00°	Depositor
Resolution (Å)	34.55 – 1.59 34.55 – 1.59	Depositor EDS
% Data completeness (in resolution range)	98.7 (34.55-1.59) 98.6 (34.55-1.59)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.60 (at 1.59Å)	Xtriage
Refinement program	PHENIX (1.11.1_2575)	Depositor
R, R_{free}	0.160 , 0.187 (Not available) , 0.179	Depositor DCC
R_{free} test set	1987 reflections (2.18%)	wwPDB-VP
Wilson B-factor (Å ²)	16.7	Xtriage
Anisotropy	0.248	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for l,k,-h 0.022 for h,-k,-l 0.017 for l,-k,h	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	6070	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.30% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: MG, NDP, 9TY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.35	0/2682	0.55	0/3620
1	B	0.34	0/2704	0.55	0/3650
All	All	0.34	0/5386	0.55	0/7270

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2609	0	2572	13	0
1	B	2622	0	2594	15	0
2	A	9	0	0	0	0
2	B	9	0	0	1	0
3	A	4	0	0	0	0
3	B	2	0	0	0	0
4	A	96	0	52	4	0
4	B	48	0	26	0	0
5	A	344	0	0	2	1
5	B	327	0	0	0	1
All	All	6070	0	5244	27	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:81:PRO:HD3	4:A:406[A]:NDP:H51A	1.74	0.68
1:A:5:TYR:OH	1:A:161[A]:ARG:NH2	2.35	0.59
1:B:51[A]:ARG:HE	1:B:55[A]:LYS:HE2	1.71	0.55
1:B:289:ARG:NH2	1:B:300:GLU:OE2	2.26	0.54
1:A:132:PRO:HA	4:A:406[B]:NDP:O7N	2.08	0.54
1:B:51[A]:ARG:HE	1:B:55[A]:LYS:HG3	1.73	0.53
1:A:314:GLU:OE2	1:B:243:MET:HB2	2.09	0.52
1:A:92[B]:GLU:OE1	5:A:501:HOH:O	2.20	0.50
1:B:51[A]:ARG:NE	1:B:55[A]:LYS:HG3	2.28	0.48
1:A:37:LYS:NZ	5:A:511:HOH:O	2.48	0.47
1:A:247:ARG:HG2	1:A:255:GLU:HG3	1.97	0.46
1:A:315[B]:LYS:HG2	1:A:318:ARG:NH2	2.30	0.46
1:A:321:ARG:HA	1:A:324:MET:HE3	1.99	0.45
1:A:315[B]:LYS:HG2	1:A:318:ARG:HH21	1.81	0.45
1:A:99:LYS:HB2	1:A:99:LYS:HE2	1.81	0.44
1:B:326:PHE:CD2	1:B:327:ILE:HG12	2.53	0.44
1:B:327:ILE:HD13	1:B:327:ILE:HA	1.85	0.43
1:B:71:LYS:HE2	1:B:96:ASN:OD1	2.18	0.43
1:B:51[A]:ARG:HA	1:B:51[A]:ARG:HD2	1.79	0.42
1:B:53:PHE:CE1	1:B:57:LYS:HD2	2.53	0.42
1:B:315:LYS:O	1:B:319:GLU:HG3	2.21	0.41
1:B:222:LEU:C	1:B:222:LEU:HD23	2.45	0.41
1:B:228:LEU:HD23	1:B:228:LEU:C	2.46	0.41
4:A:406[B]:NDP:H41N	2:B:401:9TY:C02	2.51	0.41
1:B:47:ILE:HG22	1:B:63:VAL:HG13	2.02	0.41
1:A:314:GLU:OE1	1:B:244[A]:GLU:OE2	2.38	0.41
1:A:132:PRO:HA	4:A:406[A]:NDP:O7N	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:681:HOH:O	5:B:665:HOH:O[2_545]	2.18	0.02

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	334/340 (98%)	328 (98%)	6 (2%)	0	100	100
1	B	337/340 (99%)	329 (98%)	8 (2%)	0	100	100
All	All	671/680 (99%)	657 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/283 (99%)	279 (100%)	0	100	100
1	B	282/283 (100%)	282 (100%)	0	100	100
All	All	561/566 (99%)	561 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	85	GLN
1	A	219	GLN
1	A	252	ASN
1	A	309	HIS
1	B	34	GLN
1	B	85	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	110	ASN
1	B	219	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	NDP	A	406[B]	-	51,52,52	2.38	10 (19%)	71,80,80	1.39	13 (18%)
2	9TY	B	401	3	9,9,9	0.96	0	13,14,14	1.73	4 (30%)
4	NDP	A	406[A]	-	51,52,52	2.50	11 (21%)	71,80,80	1.28	8 (11%)
2	9TY	A	401	3	9,9,9	1.18	0	13,14,14	1.14	1 (7%)
4	NDP	B	404	-	51,52,52	1.97	8 (15%)	71,80,80	1.25	7 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	NDP	A	406[B]	-	-	9/34/77/77	0/5/5/5
2	9TY	B	401	3	-	0/12/16/16	0/1/1/1
4	NDP	A	406[A]	-	-	8/34/77/77	0/5/5/5
2	9TY	A	401	3	-	3/12/16/16	0/1/1/1
4	NDP	B	404	-	-	4/34/77/77	0/5/5/5

All (29) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	406[B]	NDP	P2B-O2B	12.94	1.82	1.59
4	A	406[A]	NDP	P2B-O2B	12.00	1.80	1.59
4	B	404	NDP	P2B-O2B	8.84	1.75	1.59
4	A	406[A]	NDP	PA-O3	8.05	1.68	1.59
4	B	404	NDP	PA-O3	6.71	1.66	1.59
4	A	406[B]	NDP	PA-O3	5.88	1.65	1.59
4	A	406[A]	NDP	PN-O3	3.80	1.63	1.59
4	A	406[B]	NDP	PN-O5D	3.70	1.73	1.59
4	A	406[A]	NDP	C2A-N1A	3.27	1.39	1.33
4	A	406[A]	NDP	PN-O5D	3.08	1.71	1.59
4	B	404	NDP	PN-O3	2.74	1.62	1.59
4	A	406[A]	NDP	O2B-C2B	-2.56	1.35	1.44
4	A	406[B]	NDP	O2B-C2B	-2.53	1.35	1.44
4	B	404	NDP	C6A-N6A	2.48	1.40	1.34
4	A	406[B]	NDP	O4B-C4B	-2.43	1.39	1.45
4	B	404	NDP	C8A-N7A	2.42	1.36	1.31
4	A	406[A]	NDP	C3D-C4D	2.41	1.59	1.53
4	A	406[B]	NDP	C5A-C4A	2.38	1.43	1.39
4	A	406[A]	NDP	C3B-C4B	2.31	1.58	1.53
4	A	406[B]	NDP	C3D-C4D	2.28	1.58	1.53
4	A	406[A]	NDP	O5B-C5B	-2.26	1.36	1.44
4	A	406[A]	NDP	C6N-N1N	2.23	1.42	1.37
4	A	406[B]	NDP	C2A-N1A	2.22	1.37	1.33
4	A	406[A]	NDP	C5A-C4A	2.20	1.43	1.39
4	A	406[B]	NDP	C4A-N3A	2.12	1.38	1.34
4	B	404	NDP	C5A-C4A	2.07	1.42	1.39
4	B	404	NDP	C5B-C4B	2.06	1.57	1.51
4	A	406[B]	NDP	C7N-N7N	2.03	1.39	1.33
4	B	404	NDP	C4A-N3A	2.02	1.38	1.34

All (33) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	406[B]	NDP	P2B-O2B-C2B	-3.34	114.50	123.43
2	B	401	9TY	O06-C04-C03	-3.19	114.63	122.14
4	A	406[A]	NDP	O2B-P2B-O1X	-3.19	97.97	109.33
4	B	404	NDP	C3N-C2N-N1N	-2.95	118.88	123.20
4	A	406[B]	NDP	C2B-C1B-N9A	-2.92	108.94	113.75
4	A	406[A]	NDP	C3N-C2N-N1N	-2.82	119.06	123.20
2	B	401	9TY	C08-C03-C07	2.65	59.74	57.45
4	B	404	NDP	C3N-C7N-N7N	2.64	122.35	117.67
4	B	404	NDP	P2B-O2B-C2B	-2.63	116.40	123.43
2	B	401	9TY	O05-C04-C03	2.63	121.98	114.87
4	B	404	NDP	C3B-C2B-C1B	-2.62	97.79	102.81
4	A	406[B]	NDP	PA-O5B-C5B	-2.62	106.36	121.35
2	A	401	9TY	C08-C03-C07	2.61	59.71	57.45
4	A	406[B]	NDP	N3A-C4A-N9A	2.60	131.59	127.17
4	A	406[B]	NDP	O2N-PN-O1N	2.50	124.08	112.44
4	A	406[A]	NDP	O2N-PN-O3	2.49	114.01	107.27
4	A	406[B]	NDP	O3X-P2B-O2X	2.48	117.09	107.80
4	A	406[A]	NDP	PA-O5B-C5B	-2.40	107.59	121.35
4	A	406[A]	NDP	O3X-P2B-O2X	2.40	116.79	107.80
4	A	406[A]	NDP	C3B-C2B-C1B	-2.35	98.31	102.81
4	A	406[B]	NDP	O2B-P2B-O1X	-2.34	100.98	109.33
4	A	406[B]	NDP	C3N-C2N-N1N	-2.32	119.80	123.20
4	A	406[A]	NDP	O7N-C7N-N7N	-2.29	117.75	122.89
4	A	406[B]	NDP	O3X-P2B-O2B	-2.29	96.93	105.85
4	A	406[B]	NDP	O4B-C4B-C3B	2.27	109.66	105.15
2	B	401	9TY	O09-C02-C03	2.26	120.98	114.87
4	B	404	NDP	O3X-P2B-O2X	2.20	116.05	107.80
4	A	406[B]	NDP	O2N-PN-O3	2.20	113.21	107.27
4	B	404	NDP	C2B-C1B-N9A	-2.11	110.28	113.75
4	A	406[B]	NDP	O7N-C7N-N7N	-2.08	118.23	122.89
4	A	406[B]	NDP	C2A-N1A-C6A	-2.08	115.32	118.73
4	A	406[A]	NDP	C1D-N1N-C6N	-2.06	116.41	120.77
4	B	404	NDP	O2N-PN-O3	2.06	112.84	107.27

There are no chirality outliers.

All (24) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	406[A]	NDP	C5B-O5B-PA-O1A
4	A	406[A]	NDP	C5B-O5B-PA-O2A
4	A	406[B]	NDP	C5B-O5B-PA-O2A
4	A	406[B]	NDP	C5B-O5B-PA-O3
4	A	406[A]	NDP	O4B-C4B-C5B-O5B

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	406[B]	NDP	O4B-C4B-C5B-O5B
4	A	406[A]	NDP	C3B-C4B-C5B-O5B
4	A	406[B]	NDP	C3B-C4B-C5B-O5B
4	B	404	NDP	C2D-C1D-N1N-C6N
2	A	401	9TY	O09-C02-C03-C07
4	A	406[A]	NDP	C2D-C1D-N1N-C6N
4	A	406[A]	NDP	C5B-O5B-PA-O3
4	A	406[B]	NDP	C5B-O5B-PA-O1A
4	A	406[A]	NDP	O4D-C1D-N1N-C6N
4	B	404	NDP	O4D-C1D-N1N-C6N
4	A	406[B]	NDP	O4D-C1D-N1N-C6N
4	A	406[A]	NDP	C2B-O2B-P2B-O1X
4	B	404	NDP	O4B-C4B-C5B-O5B
4	A	406[B]	NDP	C2D-C1D-N1N-C6N
4	A	406[B]	NDP	PN-O3-PA-O1A
4	A	406[B]	NDP	PN-O3-PA-O2A
4	B	404	NDP	PN-O3-PA-O2A
2	A	401	9TY	O01-C02-C03-C07
2	A	401	9TY	O09-C02-C03-C08


There are no ring outliers.

3 monomers are involved in 4 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	406[B]	NDP	2	0
2	B	401	9TY	1	0
4	A	406[A]	NDP	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.


Ligand NDP A 406 (B)




Ligand 9TY B 401




Bond lengths



Bond angles




Torsions





Rings

Ligand NDP A 406 (A)



Ligand 9TY A 401





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	326/340 (95%)	-0.15	2 (0%) 85 89	7, 18, 29, 37	10 (3%)
1	B	326/340 (95%)	-0.09	1 (0%) 90 91	7, 19, 32, 46	13 (3%)
All	All	652/680 (95%)	-0.12	3 (0%) 87 90	7, 18, 31, 46	23 (3%)

All (3) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	327	ILE	4.0
1	A	327	ILE	2.6
1	A	309	HIS	2.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	MG	A	405	1/1	0.92	0.10	49,49,49,49	0
3	MG	A	404	1/1	0.95	0.12	25,25,25,25	0

Continued on next page...

Continued from previous page...




Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	NDP	A	406[A]	48/48	0.97	0.06	11,13,19,21	48
4	NDP	A	406[B]	48/48	0.97	0.06	10,14,19,21	48
2	9TY	A	401	9/9	0.98	0.04	10,11,12,12	0
2	9TY	B	401	9/9	0.99	0.03	10,12,13,15	0
3	MG	A	403	1/1	0.99	0.02	12,12,12,12	0
4	NDP	B	404	48/48	0.99	0.04	10,15,20,23	0
3	MG	A	402	1/1	1.00	0.01	12,12,12,12	0
3	MG	B	402	1/1	1.00	0.03	13,13,13,13	0
3	MG	B	403	1/1	1.00	0.01	12,12,12,12	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.






Electron density around NDP A 406 (B):

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 9TY A 401:


$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 9TY B 401:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.