



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 04:08 PM UTC

PDB ID : 7W4X / pdb_00007w4x
Title : Crystal structure of PDE4D catalytic domain complexed with 17
Authors : Huang, Y.-Y.; Luo, H.-B.
Deposited on : 2021-11-29
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6164 (2.20-2.20)
Clashscore	190562	6851 (2.20-2.20)
Ramachandran outliers	187476	6768 (2.20-2.20)
Sidechain outliers	187428	6769 (2.20-2.20)
RSRZ outliers	180081	6166 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	373	% 82% 5% 13%
1	B	373	19% 73% 14% • 13%

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 5558 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called cAMP-specific 3',5'-cyclic phosphodiesterase 4D.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	324	Total	C	N	O	S	0	0	0
			2622	1659	448	501	14			
1	B	324	Total	C	N	O	S	0	0	0
			2622	1659	448	501	14			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	450	ALA	-	expression tag	UNP Q08499
B	450	ALA	-	expression tag	UNP Q08499


- Molecule 2 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Zn	0	0
			1	1		
2	B	1	Total	Zn	0	0
			1	1		

- Molecule 3 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Mg	0	0
			1	1		
3	B	1	Total	Mg	0	0
			1	1		

- Molecule 4 is (2R,4S)-2-(2-hydroxyethyl)-2,6-dimethyl-4-(2-methylprop-1-enyl)-3,4-dihydropyrano[3,2-c][1,8]naphthyridin-5-one (CCD ID: 8G7) (formula: C₁₉H₂₄N₂O₃) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	A	1	Total	C	N	O	0	0
			24	19	2	3		
4	B	1	Total	C	N	O	0	0
			24	19	2	3		

- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	157	Total	O	0	0
			157	157		
5	B	105	Total	O	0	0
			105	105		

- Molecule 1: cAMP-specific 3',5'-cyclic phosphodiesterase 4D



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	57.81Å 80.30Å 163.29Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	22.63 – 2.20 22.63 – 2.20	Depositor EDS
% Data completeness (in resolution range)	99.6 (22.63-2.20) 99.5 (22.63-2.20)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.01 (at 2.19Å)	Xtriage
Refinement program	PHENIX 1.10_2155	Depositor
R, R_{free}	0.229 , 0.253 0.230 , 0.254	Depositor DCC
R_{free} test set	1881 reflections (4.77%)	wwPDB-VP
Wilson B-factor (Å ²)	25.9	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	5558	wwPDB-VP
Average B, all atoms (Å ²)	32.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.32% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: MG, ZN, 8G7

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	0/2676	0.30	0/3636
1	B	0.22	1/2676 (0.0%)	0.40	0/3636
All	All	0.20	1/5352 (0.0%)	0.35	0/7272

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	255	LYS	CD-CE	-5.14	1.37	1.52

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2622	0	2578	19	0
1	B	2622	0	2578	49	1
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
4	A	24	0	0	0	0
4	B	24	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	157	0	0	15	2
5	B	105	0	0	16	1
All	All	5558	0	5156	67	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:186:THR:O	1:B:190:ILE:HD13	1.67	0.95
1:B:101:LYS:O	5:B:601:HOH:O	1.88	0.90
1:B:397:ASP:OD1	5:B:602:HOH:O	1.91	0.88
1:B:352:MET:SD	5:B:636:HOH:O	2.35	0.85
1:A:130:ASP:O	5:A:602:HOH:O	1.98	0.81
1:A:127:GLN:NE2	5:A:606:HOH:O	2.13	0.81
1:B:186:THR:O	1:B:190:ILE:CD1	2.29	0.81
1:A:288:GLU:OE1	5:A:601:HOH:O	1.97	0.80
1:B:176:LEU:HD13	1:B:190:ILE:HG23	1.63	0.80
1:A:393:GLN:OE1	5:A:604:HOH:O	2.04	0.74
1:B:313:MET:SD	5:B:610:HOH:O	2.45	0.73
1:B:389:HIS:HB3	5:B:605:HOH:O	1.87	0.73
1:B:92:LEU:HD23	1:B:93:ALA:N	2.08	0.69
1:B:89:GLU:C	1:B:92:LEU:HD22	2.19	0.68
1:B:290:LYS:NZ	5:B:605:HOH:O	2.27	0.68
1:B:391:ASP:O	5:B:603:HOH:O	2.12	0.68
1:A:128:GLU:OE2	5:A:605:HOH:O	2.12	0.67
1:B:183:ALA:O	1:B:185:PHE:N	2.29	0.66
1:B:104:LEU:HB2	5:B:601:HOH:O	1.95	0.66
1:B:94:LYS:HA	1:B:97:GLU:HG3	1.79	0.65
1:B:282:ASP:HB3	1:B:308:GLN:HE21	1.62	0.65
1:A:178:THR:O	5:A:607:HOH:O	2.15	0.64
1:B:292:VAL:O	1:B:293:THR:OG1	2.16	0.63
1:A:133:LYS:NZ	5:A:609:HOH:O	2.16	0.62
1:A:133:LYS:N	5:A:602:HOH:O	2.32	0.62
1:B:182:GLU:O	1:B:184:VAL:N	2.26	0.61
1:A:218:GLU:OE1	1:B:239:LYS:NZ	2.28	0.60
1:A:88:GLN:N	5:A:612:HOH:O	2.35	0.59
1:A:151:ASP:O	5:A:608:HOH:O	2.16	0.58
1:B:350:ARG:HD3	5:B:636:HOH:O	2.02	0.58
1:B:92:LEU:O	1:B:96:LEU:HG	2.03	0.58

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:411:PRO:O	5:B:604:HOH:O	2.17	0.57
1:B:257:ARG:NH1	5:B:607:HOH:O	2.38	0.57
1:A:95:GLU:HG2	5:A:646:HOH:O	2.04	0.57
1:B:93:ALA:HA	1:B:96:LEU:HD12	1.87	0.56
1:B:176:LEU:HG	5:B:610:HOH:O	2.06	0.55
1:B:92:LEU:HD23	1:B:93:ALA:H	1.71	0.55
1:B:179:PRO:HA	1:B:182:GLU:OE1	2.07	0.54
1:B:190:ILE:HD12	1:B:190:ILE:N	2.23	0.53
1:B:179:PRO:O	1:B:182:GLU:HG2	2.07	0.53
1:A:91:VAL:HG23	5:A:665:HOH:O	2.08	0.53
1:B:168:VAL:O	1:B:172:THR:HG23	2.09	0.53
1:A:255:LYS:HD3	1:A:255:LYS:C	2.34	0.52
1:B:292:VAL:HG23	1:B:298:LEU:HA	1.92	0.52
1:B:179:PRO:HA	1:B:182:GLU:CD	2.34	0.52
1:B:141:THR:HG22	1:B:248:ILE:HG22	1.91	0.52
1:B:176:LEU:CD1	1:B:190:ILE:HG23	2.36	0.51
1:B:302:ASN:ND2	5:B:615:HOH:O	2.44	0.50
1:B:293:THR:O	1:B:295:SER:N	2.47	0.48
1:A:104:LEU:HD21	1:A:109:ILE:HD12	1.96	0.47
1:B:132:LEU:HG	1:B:137:ILE:O	2.14	0.47
1:A:407:GLN:NE2	5:A:603:HOH:O	1.98	0.47
1:B:171:SER:O	1:B:175:LEU:HG	2.14	0.47
1:A:407:GLN:HG2	5:A:603:HOH:O	2.12	0.47
1:B:132:LEU:HD11	1:B:139:VAL:HB	1.98	0.46
1:B:350:ARG:NH2	5:B:618:HOH:O	2.49	0.46
1:B:182:GLU:HA	5:B:614:HOH:O	2.15	0.45
1:B:323:THR:HB	1:B:395:ILE:HG23	1.98	0.44
1:B:182:GLU:C	1:B:184:VAL:H	2.22	0.44
1:B:293:THR:C	1:B:296:GLY:H	2.26	0.44
1:B:183:ALA:N	5:B:614:HOH:O	2.44	0.43
1:A:290:LYS:NZ	5:A:614:HOH:O	2.43	0.43
1:B:104:LEU:HD21	1:B:109:ILE:HD12	2.01	0.43
1:A:98:ASP:HB2	1:A:104:LEU:HD12	2.01	0.42
1:B:262:LYS:HE3	1:B:266:ASP:OD2	2.19	0.42
1:B:389:HIS:CE1	1:B:390:PRO:HB3	2.55	0.41
1:B:389:HIS:HA	1:B:390:PRO:HA	1.70	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:744:HOH:O	5:B:615:HOH:O[3_644]	2.08	0.12
1:B:301:ASP:OD1	5:A:608:HOH:O[3_654]	2.16	0.04

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	322/373 (86%)	317 (98%)	5 (2%)	0	100	100
1	B	322/373 (86%)	304 (94%)	15 (5%)	3 (1%)	14	14
All	All	644/746 (86%)	621 (96%)	20 (3%)	3 (0%)	24	27

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	184	VAL
1	B	293	THR
1	B	294	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	296/339 (87%)	296 (100%)	0	100	100
1	B	296/339 (87%)	295 (100%)	1 (0%)	86	93
All	All	592/678 (87%)	591 (100%)	1 (0%)	87	94

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	148	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	250	GLN
1	A	369	GLN
1	B	210	GLN
1	B	245	ASN
1	B	302	ASN
1	B	308	GLN
1	B	321	ASN
1	B	369	GLN
1	B	378	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

Of 6 ligands modelled in this entry, 4 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	8G7	B	503	2	26,26,26	0.30	0	32,39,39	0.91	1 (3%)
4	8G7	A	503	-	26,26,26	0.34	0	32,39,39	0.91	1 (3%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	8G7	B	503	2	-	3/8/22/22	0/3/3/3
4	8G7	A	503	-	-	1/8/22/22	0/3/3/3

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	503	8G7	O1-C3-C12	4.10	110.43	104.10
4	B	503	8G7	O1-C3-C12	4.04	110.33	104.10

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	A	503	8G7	C3-C4-C5-O2
4	B	503	8G7	C7-C8-C9-C11
4	B	503	8G7	C3-C4-C5-O2
4	B	503	8G7	C7-C8-C9-C10


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.

The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	324/373 (86%)	-0.03	2 (0%) 85 83	12, 25, 41, 54	0
1	B	324/373 (86%)	1.12	71 (21%) 2 1	16, 39, 63, 87	0
All	All	648/746 (86%)	0.54	73 (11%) 10 8	12, 30, 58, 87	0

All (73) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	294	SER	5.4
1	B	297	VAL	5.3
1	B	105	HIS	4.7
1	B	295	SER	4.6
1	B	184	VAL	4.5
1	B	132	LEU	4.3
1	B	91	VAL	4.2
1	B	245	ASN	4.1
1	B	387	LEU	4.0
1	B	296	GLY	4.0
1	B	92	LEU	3.8
1	B	179	PRO	3.8
1	B	178	THR	3.7
1	B	190	ILE	3.6
1	B	182	GLU	3.6
1	B	293	THR	3.6
1	B	393	GLN	3.4
1	B	289	THR	3.3
1	B	298	LEU	3.2
1	B	176	LEU	3.2
1	B	183	ALA	3.2
1	B	384	TRP	3.1
1	B	135	PHE	3.1
1	B	181	LEU	3.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	99	VAL	3.0
1	B	185	PHE	3.0
1	B	392	ALA	3.0
1	B	385	ALA	2.9
1	B	292	VAL	2.9
1	B	88	GLN	2.9
1	B	351	GLY	2.9
1	B	103	GLY	2.9
1	B	130	ASP	2.9
1	B	93	ALA	2.9
1	B	177	SER	2.8
1	B	131	LEU	2.8
1	B	102	TRP	2.7
1	B	112	LEU	2.7
1	B	386	ASP	2.6
1	B	306	ARG	2.6
1	B	127	GLN	2.6
1	B	191	LEU	2.6
1	B	180	ALA	2.5
1	B	126	PHE	2.5
1	B	125	ILE	2.5
1	B	362	ASN	2.5
1	B	133	LYS	2.4
1	B	141	THR	2.4
1	B	391	ASP	2.4
1	B	398	THR	2.4
1	B	101	LYS	2.4
1	B	139	VAL	2.3
1	B	98	ASP	2.3
1	B	147	MET	2.3
1	A	409	THR	2.2
1	B	299	LEU	2.2
1	B	174	VAL	2.2
1	A	364	SER	2.2
1	B	143	ILE	2.2
1	B	390	PRO	2.2
1	B	95	GLU	2.2
1	B	395	ILE	2.1
1	B	254	LYS	2.1
1	B	350	ARG	2.1
1	B	109	ILE	2.1
1	B	287	VAL	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	186	THR	2.0
1	B	253	THR	2.0
1	B	246	CYS	2.0
1	B	389	HIS	2.0
1	B	137	ILE	2.0
1	B	286	MET	2.0
1	B	96	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	8G7	B	503	24/24	0.86	0.12	17,29,37,39	0
4	8G7	A	503	24/24	0.88	0.11	19,28,34,45	0
3	MG	B	502	1/1	0.95	0.05	17,17,17,17	0
3	MG	A	502	1/1	0.96	0.03	15,15,15,15	0
2	ZN	A	501	1/1	0.99	0.03	20,20,20,20	0
2	ZN	B	501	1/1	0.99	0.03	28,28,28,28	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around 8G7 B 503:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around 8G7 A 503:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.