



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 01:26 AM UTC

PDB ID : 8W4N / pdb_00008w4n
Title : Crystal structure of EndoSz mutant D234M, in space group P21, in complex with oligosaccharide G2S1
Authors : Guan, H.H.; Lin, C.C.; Hsieh, Y.C.; Chen, C.J.
Deposited on : 2023-08-24
Resolution : 3.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1456 (3.10-3.10)
Clashscore	190562	1539 (3.10-3.10)
Ramachandran outliers	187476	1467 (3.10-3.10)
Sidechain outliers	187428	1467 (3.10-3.10)
RSRZ outliers	180081	1456 (3.10-3.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	992	
2	B	9	

2 Entry composition [i](#)


There are 3 unique types of molecules in this entry. The entry contains 7021 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Glycoside hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	876	Total	C	N	O	S	0	0	0
			6902	4375	1152	1360	15			

- Molecule 2 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-6)-beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-3)-[beta-D-galactopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-2)-alpha-D-mannopyranose-(1-6)]beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	B	9	Total	C	N	O	0	0	0
			118	65	4	49			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	1	Total	Ca	0	0
			1	1		

- Molecule 1: Glycoside hydrolase



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.59Å 101.69Å 129.80Å 90.00° 90.72° 90.00°	Depositor
Resolution (Å)	28.33 – 3.10 28.33 – 3.10	Depositor EDS
% Data completeness (in resolution range)	98.4 (28.33-3.10) 98.2 (28.33-3.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.46 (at 3.11Å)	Xtriage
Refinement program	PHENIX (1.18.2_3874: ???)	Depositor
R, R_{free}	0.227 , 0.273 0.230 , 0.278	Depositor DCC
R_{free} test set	1990 reflections (8.12%)	wwPDB-VP
Wilson B-factor (Å ²)	71.0	Xtriage
Anisotropy	0.457	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.32 , 56.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.30$	Xtriage
Estimated twinning fraction	0.017 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7021	wwPDB-VP
Average B, all atoms (Å ²)	79.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GAL, NAG, BMA, SIA, CA, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.19	1/7035 (0.0%)	0.44	6/9501 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	604	ARG	CA-C	5.06	1.59	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	616	LYS	N-CA-C	-8.93	100.39	111.11
1	A	605	GLN	N-CA-C	7.17	120.51	111.69
1	A	603	ASN	CA-C-N	6.19	133.36	121.54
1	A	603	ASN	C-N-CA	6.19	133.36	121.54
1	A	615	THR	N-CA-C	-5.63	105.51	112.72
1	A	623	LYS	N-CA-C	-5.11	107.43	113.97

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6902	0	6887	79	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	118	0	100	1	0
3	A	1	0	0	0	0
All	All	7021	0	6987	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (79) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:797:THR:HG22	1:A:798:SER:H	1.44	0.82
1:A:742:ASN:HD21	1:A:745:LYS:HE2	1.50	0.74
1:A:943:LYS:HE3	1:A:967:ASP:HB3	1.70	0.73
1:A:597:LEU:HD22	1:A:607:LEU:HD12	1.72	0.72
1:A:151:PHE:HE1	1:A:234:MET:HE1	1.56	0.70
1:A:156:LYS:HB2	1:A:159:SER:HB2	1.76	0.67
1:A:496:LEU:HD13	1:A:499:LEU:HD22	1.79	0.63
1:A:228:LEU:O	1:A:275:ARG:NH1	2.30	0.63
1:A:151:PHE:CE1	1:A:234:MET:HE1	2.36	0.61
1:A:579:ALA:HB2	1:A:606:ILE:HG23	1.82	0.60
1:A:795:LEU:H	1:A:795:LEU:HD23	1.68	0.59
1:A:280:ASP:OD2	1:A:304:GLN:NE2	2.35	0.59
1:A:583:THR:O	1:A:613:THR:CG2	2.51	0.58
1:A:282:THR:HG22	1:A:304:GLN:HB2	1.85	0.58
1:A:614:VAL:HG22	1:A:614:VAL:O	2.05	0.56
1:A:182:ARG:HB2	1:A:228:LEU:HD12	1.87	0.56
1:A:678:TYR:OH	1:A:696:TYR:O	2.20	0.55
1:A:301:LEU:HD13	1:A:337:ILE:HD13	1.87	0.55
1:A:717:LEU:HB2	1:A:905:ILE:HD13	1.90	0.54
1:A:302:LEU:HB3	1:A:399:PHE:CE1	2.43	0.54
1:A:958:ALA:O	1:A:962:ASN:ND2	2.36	0.52
1:A:603:ASN:O	1:A:603:ASN:OD1	2.26	0.52
1:A:839:VAL:C	1:A:841:HIS:H	2.17	0.52
1:A:446:GLU:OE2	1:A:470:ARG:NH1	2.42	0.52
1:A:789:ASN:ND2	1:A:822:SER:OG	2.43	0.52
1:A:797:THR:HG22	1:A:798:SER:N	2.20	0.51
1:A:951:LEU:HD23	1:A:961:ILE:HD13	1.92	0.51
1:A:798:SER:O	1:A:895:GLY:N	2.44	0.51
1:A:614:VAL:HG22	1:A:619:GLY:O	2.12	0.50
1:A:839:VAL:O	1:A:841:HIS:N	2.42	0.50
1:A:833:ALA:HB3	1:A:864:TYR:HB3	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:118:TYR:HA	1:A:149:PHE:HB2	1.94	0.50
1:A:785:ASP:OD1	1:A:786:GLY:N	2.45	0.49
1:A:779:ASN:ND2	1:A:794:THR:OG1	2.45	0.49
1:A:456:LYS:HG2	1:A:459:ARG:HH12	1.77	0.49
1:A:583:THR:HA	1:A:613:THR:HG21	1.95	0.48
1:A:123:HIS:HE1	1:A:424:ILE:HD12	1.77	0.48
1:A:574:LEU:HD12	1:A:603:ASN:HB2	1.96	0.47
1:A:742:ASN:ND2	1:A:745:LYS:HB2	2.29	0.47
1:A:138:MET:HE1	1:A:150:VAL:HG22	1.97	0.47
1:A:574:LEU:HB2	1:A:603:ASN:HB3	1.96	0.47
1:A:488:LYS:HG2	1:A:509:GLN:OE1	2.14	0.47
1:A:607:LEU:HD21	1:A:625:PHE:HE2	1.80	0.47
1:A:604:ARG:HH12	1:A:686:ILE:HG23	1.80	0.46
1:A:791:ASP:OD1	1:A:793:THR:HG23	2.16	0.46
1:A:799:ASN:HB2	1:A:893:GLN:O	2.15	0.46
1:A:553:LEU:HD11	1:A:555:ILE:HD11	1.98	0.46
1:A:456:LYS:HG2	1:A:459:ARG:HH22	1.80	0.46
1:A:604:ARG:NH1	1:A:608:ASP:OD1	2.48	0.46
1:A:712:VAL:HB	1:A:720:THR:HG23	1.97	0.46
1:A:302:LEU:HD22	1:A:343:MET:HB2	1.97	0.46
1:A:821:ASP:O	1:A:823:LYS:N	2.48	0.46
1:A:304:GLN:NE2	2:B:1:NAG:H83	2.31	0.46
1:A:186:TRP:HB3	1:A:239:SER:O	2.16	0.45
1:A:587:LYS:HG3	1:A:624:THR:HA	1.98	0.45
1:A:630:GLN:HG3	1:A:691:PHE:HD2	1.81	0.45
1:A:579:ALA:HB1	1:A:610:MET:HG3	1.98	0.45
1:A:940:LEU:HD13	1:A:971:TYR:HB2	1.99	0.45
1:A:849:LYS:HG3	1:A:850:ASP:N	2.32	0.45
1:A:604:ARG:NH1	1:A:686:ILE:HG23	2.33	0.44
1:A:574:LEU:HG	1:A:595:LEU:HD22	1.99	0.44
1:A:376:ILE:HG22	1:A:431:VAL:HG13	2.00	0.43
1:A:515:SER:HB3	1:A:569:PHE:HZ	1.83	0.43
1:A:280:ASP:HA	1:A:302:LEU:O	2.17	0.43
1:A:614:VAL:HA	1:A:617:HIS:HB3	2.01	0.43
1:A:822:SER:C	1:A:824:ILE:H	2.27	0.43
1:A:329:ARG:O	1:A:332:SER:OG	2.33	0.43
1:A:564:LEU:HD23	1:A:610:MET:HE1	2.01	0.42
1:A:672:ILE:HG23	1:A:677:ASP:HB3	2.00	0.42
1:A:570:ASP:HB3	1:A:594:LYS:HE2	2.01	0.42
1:A:165:LEU:HA	1:A:169:HIS:HB2	2.01	0.42
1:A:345:GLY:HA3	1:A:399:PHE:CZ	2.55	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:481:ARG:NH1	1:A:503:GLU:OE1	2.46	0.42
1:A:152:HIS:CE1	1:A:155:THR:HB	2.55	0.41
1:A:817:ARG:HG3	1:A:872:SER:HB3	2.01	0.41
1:A:878:ILE:HG22	1:A:880:ALA:H	1.84	0.41
1:A:822:SER:C	1:A:824:ILE:N	2.78	0.41
1:A:787:LEU:HD22	1:A:791:ASP:H	1.85	0.41
1:A:268:PRO:HD2	1:A:297:TYR:O	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	874/992 (88%)	834 (95%)	37 (4%)	3 (0%)	36 67

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	604	ARG
1	A	791	ASP
1	A	841	HIS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	753/847 (89%)	750 (100%)	3 (0%)	84 86

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	616	LYS
1	A	842	LEU
1	A	845	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (10) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	152	HIS
1	A	169	HIS
1	A	174	ASN
1	A	414	ASN
1	A	673	ASN
1	A	742	ASN
1	A	779	ASN
1	A	789	ASN
1	A	877	ASN
1	A	957	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	NAG	B	1	2	15,15,15	0.60	0	21,21,21	0.97	1 (4%)
2	BMA	B	2	2	11,11,12	0.92	0	15,15,17	1.23	1 (6%)
2	MAN	B	3	2	11,11,12	0.75	0	15,15,17	1.10	2 (13%)
2	NAG	B	4	2	14,14,15	0.23	0	17,19,21	0.43	0
2	GAL	B	5	2	11,11,12	0.79	1 (9%)	15,15,17	0.98	0
2	SIA	B	6	2	20,20,21	1.62	2 (10%)	21,28,31	1.51	2 (9%)
2	MAN	B	7	2	11,11,12	0.64	0	15,15,17	1.10	2 (13%)
2	NAG	B	8	2	14,14,15	0.39	0	17,19,21	0.55	0
2	GAL	B	9	2	11,11,12	0.52	0	15,15,17	1.05	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	B	1	2	-	4/6/26/26	0/1/1/1
2	BMA	B	2	2	-	0/2/19/22	0/1/1/1
2	MAN	B	3	2	-	0/2/19/22	0/1/1/1
2	NAG	B	4	2	-	2/6/23/26	0/1/1/1
2	GAL	B	5	2	-	2/2/19/22	0/1/1/1
2	SIA	B	6	2	-	14/18/34/38	0/1/1/1
2	MAN	B	7	2	-	2/2/19/22	0/1/1/1
2	NAG	B	8	2	-	0/6/23/26	0/1/1/1
2	GAL	B	9	2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	6	SIA	C2-C1	5.58	1.59	1.52
2	B	6	SIA	O6-C2	2.23	1.47	1.43
2	B	5	GAL	C1-C2	2.07	1.57	1.52

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	6	SIA	O1A-C1-C2	-4.86	112.34	122.85
2	B	7	MAN	C1-O5-C5	3.23	116.51	112.19
2	B	2	BMA	C1-C2-C3	3.05	114.09	109.64
2	B	3	MAN	C1-O5-C5	2.71	115.82	112.19
2	B	9	GAL	C1-O5-C5	2.69	115.79	112.19
2	B	6	SIA	O1B-C1-O1A	2.63	130.04	124.08
2	B	1	NAG	C1-O5-C5	2.51	118.51	113.65
2	B	7	MAN	O2-C2-C3	-2.43	105.12	110.15
2	B	3	MAN	O2-C2-C3	-2.39	105.20	110.15

There are no chirality outliers.

All (24) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	6	SIA	C5-C6-C7-C8
2	B	6	SIA	C5-C6-C7-O7
2	B	6	SIA	O6-C6-C7-C8
2	B	6	SIA	O6-C6-C7-O7
2	B	6	SIA	O8-C8-C9-O9
2	B	6	SIA	C7-C8-C9-O9
2	B	7	MAN	O5-C5-C6-O6
2	B	1	NAG	O5-C5-C6-O6
2	B	7	MAN	C4-C5-C6-O6
2	B	6	SIA	C11-C10-N5-C5
2	B	6	SIA	O10-C10-N5-C5
2	B	1	NAG	C4-C5-C6-O6
2	B	6	SIA	C6-C7-C8-O8
2	B	5	GAL	O5-C5-C6-O6
2	B	5	GAL	C4-C5-C6-O6
2	B	6	SIA	C6-C7-C8-C9
2	B	6	SIA	O7-C7-C8-O8
2	B	1	NAG	C3-C2-N2-C7
2	B	4	NAG	C4-C5-C6-O6
2	B	6	SIA	O7-C7-C8-C9
2	B	6	SIA	O1A-C1-C2-O6
2	B	6	SIA	O1A-C1-C2-C3
2	B	1	NAG	C1-C2-N2-C7
2	B	4	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	1	NAG	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	876/992 (88%)	0.31	24 (2%) 56 35	30, 75, 132, 184	0

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	939	GLU	4.3
1	A	937	LEU	4.2
1	A	940	LEU	4.1
1	A	788	LEU	3.7
1	A	853	GLU	3.2
1	A	622	GLU	3.1
1	A	947	LEU	3.0
1	A	793	THR	3.0
1	A	797	THR	2.8
1	A	790	ASN	2.8
1	A	604	ARG	2.6
1	A	598	ALA	2.5
1	A	942	VAL	2.5
1	A	842	LEU	2.5
1	A	787	LEU	2.4
1	A	789	ASN	2.3
1	A	599	ALA	2.3
1	A	792	THR	2.3
1	A	799	ASN	2.2
1	A	684	GLN	2.2
1	A	971	TYR	2.2
1	A	798	SER	2.2
1	A	538	ASN	2.2
1	A	794	THR	2.1

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	GAL	B	9	11/12	0.20	0.22	173,195,201,208	0
2	SIA	B	6	20/21	0.52	0.16	154,176,184,184	0
2	NAG	B	8	14/15	0.56	0.24	101,146,184,192	0
2	GAL	B	5	11/12	0.60	0.12	103,124,150,160	0
2	MAN	B	7	11/12	0.82	0.19	71,99,112,130	0
2	NAG	B	4	14/15	0.83	0.14	68,83,104,111	0
2	NAG	B	1	15/15	0.87	0.16	45,61,85,96	0
2	MAN	B	3	11/12	0.91	0.10	51,56,71,80	0
2	BMA	B	2	11/12	0.95	0.11	52,61,80,95	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	A	1101	1/1	0.72	0.11	142,142,142,142	0

6.5 Other polymers [i](#)

There are no such residues in this entry.