



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 07:11 AM UTC

PDB ID : 5W5C / pdb_00005w5c
Title : Crystal structure of the primed SNARE-Complexin-Synaptotagmin-1 C2AB complex
Authors : Zhou, Q.; Brunger, A.T.
Deposited on : 2017-06-14
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	40	
2	B	67	
3	C	77	
4	D	65	
5	E	83	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
6	F	282	25% 75% 18% 7%

2 Entry composition

There are 9 unique types of molecules in this entry. The entry contains 4044 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vesicle-associated membrane protein 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	38	Total	C	N	O	S	0	1	0
			319	191	62	65	1			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	27	GLY	-	expression tag	UNP P63045

- Molecule 2 is a protein called Syntaxin-1A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	54	Total	C	N	O	S	0	0	0
			433	268	73	87	5			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	190	MET	-	initiating methionine	UNP P32851

- Molecule 3 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	C	65	Total	C	N	O	S	0	1	0
			501	297	91	108	5			

- Molecule 4 is a protein called Synaptosomal-associated protein 25.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	D	55	Total	C	N	O	S	1	0	0
			407	241	77	85	4			

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
D	140	MET	-	initiating methionine	UNP P60881

- Molecule 5 is a protein called Complexin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	E	25	Total	C	N	O	S	0	1	0
			193	118	40	33	2			


- Molecule 6 is a protein called Synaptotagmin-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	F	261	Total	C	N	O	S	2	1	0
			1881	1201	320	355	5			

- Molecule 7 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	C	1	Total	Mg	0	0
			1	1		

- Molecule 8 is GLYCEROL (CCD ID: GOL) (formula: C₃H₈O₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	D	1	Total	C	O	0	0
			6	3	3		
8	F	1	Total	C	O	0	0
			6	3	3		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
8	F	1	Total	C	O	0	0
			6	3	3		
8	F	1	Total	C	O	0	0
			6	3	3		
8	F	1	Total	C	O	0	0
			6	3	3		

- Molecule 9 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	A	23	Total	O	0	0
			23	23		
9	B	21	Total	O	0	0
			21	21		
9	C	43	Total	O	0	0
			43	43		
9	D	35	Total	O	0	0
			35	35		
9	E	12	Total	O	0	0
			12	12		
9	F	145	Total	O	0	0
			145	145		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vesicle-associated membrane protein 2

Chain A: 



- Molecule 2: Syntaxin-1A

Chain B: 




- Molecule 3: Synaptosomal-associated protein 25

Chain C: 




- Molecule 4: Synaptosomal-associated protein 25


Chain D: 




- Molecule 5: Complexin-1

Chain E: 





● Molecule 6: Synaptotagmin-1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	85.66Å 89.68Å 91.73Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	64.13 – 1.85 64.13 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.6 (64.13-1.85) 99.7 (64.13-1.85)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.05 (at 1.86Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.195 , 0.232 0.202 , 0.236	Depositor DCC
R_{free} test set	2007 reflections (3.29%)	wwPDB-VP
Wilson B-factor (Å ²)	41.5	Xtriage
Anisotropy	0.546	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 71.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.010 for -h,l,k	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4044	wwPDB-VP
Average B, all atoms (Å ²)	75.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.14% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.75	0/321	0.73	0/430
2	B	0.75	0/437	0.73	0/586
3	C	0.67	0/504	0.69	0/675
4	D	0.78	0/407	0.76	0/547
5	E	0.59	0/196	0.73	0/258
6	F	0.73	0/1920	0.72	0/2616
All	All	0.73	0/3785	0.72	0/5112

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	319	0	321	6	0
2	B	433	0	416	5	0
3	C	501	0	463	9	0
4	D	407	0	371	4	0
5	E	193	0	187	5	0
6	F	1881	0	1676	45	0
7	C	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
8	D	6	0	8	1	0
8	F	24	0	32	6	0
9	A	23	0	0	3	0
9	B	21	0	0	0	0
9	C	43	0	0	1	1
9	D	35	0	0	0	2
9	E	12	0	0	1	0
9	F	145	0	0	11	3
All	All	4044	0	3474	64	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:366:LYS:NZ	9:F:601:HOH:O	2.01	0.92
3:C:41[B]:ASP:OD2	3:C:45:ARG:NH2	2.03	0.92
6:F:324:LYS:HE2	6:F:326:LYS:HE2	1.67	0.77
6:F:381:ASN:O	6:F:381:ASN:ND2	2.18	0.76
3:C:52:GLU:OE1	9:C:201:HOH:O	2.08	0.72
6:F:327:LYS:NZ	9:F:603:HOH:O	2.24	0.70
6:F:281:ARG:NH2	9:F:604:HOH:O	2.26	0.68
1:A:44[A]:ASP:OD1	9:A:101:HOH:O	2.11	0.67
6:F:319:ASN:H	8:F:503:GOL:H2	1.63	0.63
6:F:354:LYS:HG2	8:F:504:GOL:H2	1.81	0.61
3:C:71:MET:HB3	4:D:191:ARG:HG2	1.85	0.59
4:D:155:GLY:HA2	8:D:301:GOL:H11	1.85	0.59
1:A:44[B]:ASP:OD1	9:A:102:HOH:O	2.17	0.59
1:A:51:ASP:OD1	5:E:67[A]:ARG:NH1	2.35	0.59
6:F:246:PRO:O	6:F:249:THR:HG22	2.03	0.59
6:F:348:PRO:HG2	6:F:351:GLN:HG2	1.85	0.57
6:F:326:LYS:NZ	9:F:605:HOH:O	2.30	0.54
6:F:380:TYR:CE2	8:F:503:GOL:H32	2.42	0.54
2:B:199:HIS:HE1	3:C:24:GLU:HB3	1.73	0.54
6:F:184:PHE:CB	6:F:225:VAL:O	2.56	0.53
6:F:281:ARG:NH2	6:F:398:ARG:O	2.42	0.53
3:C:20:GLN:NE2	6:F:350:GLU:HG2	2.23	0.53
6:F:324:LYS:HG3	9:F:719:HOH:O	2.09	0.53
3:C:17:ARG:NH1	6:F:349:PHE:HB3	2.26	0.51
1:A:47:ARG:N	2:B:215:MET:HE1	2.26	0.51

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:F:182:LYS:HA	6:F:193:PHE:O	2.11	0.51
6:F:275:ASP:OD2	9:F:602:HOH:O	2.20	0.50
5:E:69:LYS:HE3	5:E:70:TYR:CZ	2.47	0.50
5:E:68:ASP:OD1	9:E:101:HOH:O	2.18	0.50
3:C:17:ARG:HH12	6:F:349:PHE:HB3	1.76	0.50
6:F:272:LYS:NZ	9:F:614:HOH:O	2.45	0.49
6:F:307:LEU:O	9:F:601:HOH:O	2.20	0.49
1:A:31:ARG:HD2	9:A:111:HOH:O	2.12	0.48
2:B:199:HIS:CE1	3:C:24:GLU:HB3	2.49	0.48
6:F:220:GLY:HA2	6:F:247:MET:CB	2.44	0.47
6:F:220:GLY:HA3	6:F:248:ASN:HB3	1.95	0.47
6:F:388[A]:ARG:NH2	9:F:616:HOH:O	2.46	0.47
6:F:354:LYS:HG2	8:F:504:GOL:C2	2.44	0.46
4:D:190:THR:O	4:D:194:GLU:HG3	2.16	0.46
6:F:327:LYS:HG3	8:F:502:GOL:H32	1.97	0.46
8:F:502:GOL:H11	9:F:664:HOH:O	2.15	0.46
3:C:20:GLN:HE22	6:F:350:GLU:HG2	1.81	0.45
1:A:47:ARG:CA	2:B:215:MET:HE1	2.46	0.45
6:F:223:THR:HG23	6:F:245:VAL:O	2.16	0.45
6:F:348:PRO:HB2	6:F:350:GLU:OE1	2.15	0.45
6:F:307:LEU:HD22	6:F:333:ASN:OD1	2.17	0.45
5:E:62:MET:HE3	5:E:62:MET:HB3	1.83	0.45
6:F:178:ASP:O	6:F:231:PHE:N	2.42	0.45
6:F:186:LEU:CB	6:F:223:THR:H	2.29	0.44
6:F:301:LYS:HE3	6:F:301:LYS:HB3	1.69	0.44
6:F:379:GLY:HA2	6:F:390:TRP:CE2	2.53	0.43
6:F:311:TYR:HB3	6:F:330:ILE:HD13	2.00	0.43
6:F:168:LEU:HB2	6:F:202:LEU:O	2.19	0.43
6:F:284:PRO:HB3	6:F:349:PHE:CD2	2.53	0.43
6:F:366:LYS:NZ	9:F:607:HOH:O	2.52	0.42
6:F:310:PRO:HA	6:F:362:LEU:O	2.19	0.42
6:F:163:ILE:HA	6:F:207:ASN:H	1.84	0.41
4:D:153:VAL:O	4:D:157:ILE:HG12	2.21	0.41
2:B:198:ARG:HD3	2:B:198:ARG:HA	1.93	0.41
6:F:302:MET:HB2	6:F:366:LYS:HE2	2.02	0.41
5:E:68:ASP:HB3	6:F:408:GLN:HG2	2.03	0.40
6:F:147:TYR:O	6:F:257:GLU:HA	2.20	0.40
6:F:223:THR:OG1	6:F:246:PRO:HA	2.21	0.40
6:F:285:THR:H	6:F:285:THR:HG23	1.64	0.40

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
9:D:432:HOH:O	9:F:713:HOH:O[3_444]	2.04	0.16
9:C:233:HOH:O	9:F:666:HOH:O[3_444]	2.10	0.10
9:D:435:HOH:O	9:F:704:HOH:O[3_444]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	37/40 (92%)	36 (97%)	1 (3%)	0	100	100
2	B	52/67 (78%)	51 (98%)	1 (2%)	0	100	100
3	C	64/77 (83%)	63 (98%)	1 (2%)	0	100	100
4	D	53/65 (82%)	53 (100%)	0	0	100	100
5	E	24/83 (29%)	24 (100%)	0	0	100	100
6	F	254/282 (90%)	237 (93%)	17 (7%)	0	100	100
All	All	484/614 (79%)	464 (96%)	20 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	37/38 (97%)	37 (100%)	0	100	100
2	B	48/61 (79%)	48 (100%)	0	100	100
3	C	52/69 (75%)	52 (100%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	D	40/56 (71%)	40 (100%)	0	100	100
5	E	16/65 (25%)	16 (100%)	0	100	100
6	F	170/252 (68%)	170 (100%)	0	100	100
All	All	363/541 (67%)	363 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
6	F	319	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
8	GOL	F	503	-	5,5,5	0.40	0	5,5,5	0.62	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
8	GOL	F	504	-	5,5,5	0.45	0	5,5,5	0.52	0
8	GOL	F	501	-	5,5,5	0.31	0	5,5,5	0.66	0
8	GOL	F	502	-	5,5,5	0.36	0	5,5,5	1.26	1 (20%)
8	GOL	D	301	-	5,5,5	0.65	0	5,5,5	1.45	2 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
8	GOL	F	503	-	-	2/4/4/4	-
8	GOL	F	504	-	-	4/4/4/4	-
8	GOL	F	501	-	-	2/4/4/4	-
8	GOL	F	502	-	-	2/4/4/4	-
8	GOL	D	301	-	-	2/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
8	F	502	GOL	C3-C2-C1	-2.23	103.63	111.80
8	D	301	GOL	O3-C3-C2	2.11	119.90	110.38
8	D	301	GOL	O2-C2-C3	2.01	117.51	109.18

There are no chirality outliers.

All (12) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
8	D	301	GOL	O1-C1-C2-C3
8	F	501	GOL	O1-C1-C2-C3
8	F	502	GOL	O1-C1-C2-C3
8	F	503	GOL	O1-C1-C2-C3
8	F	504	GOL	O1-C1-C2-C3
8	F	504	GOL	C1-C2-C3-O3
8	F	501	GOL	O1-C1-C2-O2
8	F	502	GOL	O1-C1-C2-O2
8	F	503	GOL	O1-C1-C2-O2
8	D	301	GOL	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
8	F	504	GOL	O1-C1-C2-O2
8	F	504	GOL	O2-C2-C3-O3

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
8	F	503	GOL	2	0
8	F	504	GOL	2	0
8	F	502	GOL	2	0
8	D	301	GOL	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	38/40 (95%)	0.64	0 100 100	29, 52, 94, 122	1 (2%)
2	B	54/67 (80%)	0.89	11 (20%) 3 2	35, 62, 113, 128	1 (1%)
3	C	65/77 (84%)	0.86	7 (10%) 11 10	26, 64, 134, 154	3 (4%)
4	D	55/65 (84%)	0.67	4 (7%) 21 23	33, 55, 140, 142	1 (1%)
5	E	25/83 (30%)	1.11	2 (8%) 18 19	33, 66, 129, 135	1 (4%)
6	F	261/282 (92%)	1.26	71 (27%) 1 1	28, 73, 166, 181	5 (1%)
All	All	498/614 (81%)	1.04	95 (19%) 3 3	26, 66, 159, 181	12 (2%)

All (95) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
6	F	243	PHE	5.1
6	F	216	TYR	4.6
6	F	147	TYR	4.5
6	F	219	LEU	4.1
6	F	245	VAL	4.0
6	F	239	ILE	3.9
6	F	206	PHE	3.9
6	F	349	PHE	3.8
6	F	186	LEU	3.8
6	F	231	PHE	3.7
6	F	145	LEU	3.7
5	E	52	TYR	3.6
3	C	11	LEU	3.6
6	F	352	ILE	3.5
6	F	169	PRO	3.4
6	F	252	PHE	3.4
6	F	181	VAL	3.4
6	F	227	ALA	3.4
6	F	224	LEU	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
6	F	184	PHE	3.3
6	F	214	VAL	3.3
4	D	192	ILE	3.2
6	F	241	GLY	3.2
6	F	230	ASP	3.2
3	C	21	LEU	3.1
6	F	247	MET	3.1
6	F	240	ILE	3.1
6	F	143	GLY	3.1
6	F	149	LEU	3.0
6	F	228	VAL	3.0
6	F	185	LEU	3.0
6	F	168	LEU	2.9
6	F	250	VAL	2.9
6	F	262	LEU	2.9
6	F	307	LEU	2.9
2	B	244	VAL	2.9
6	F	259	TRP	2.9
2	B	195	ILE	2.9
4	D	142	ARG	2.7
6	F	261	ASP	2.7
6	F	416	MET	2.7
6	F	151	TYR	2.7
6	F	254	HIS	2.7
6	F	229	TYR	2.7
3	C	71	MET	2.7
6	F	223	THR	2.7
6	F	202	LEU	2.6
2	B	239	HIS	2.6
6	F	153	PHE	2.6
6	F	193	PHE	2.6
6	F	187	PRO	2.6
6	F	246	PRO	2.6
6	F	225	VAL	2.6
6	F	304	VAL	2.6
6	F	418	ALA	2.6
6	F	195	THR	2.6
6	F	260	ARG	2.5
6	F	162	ILE	2.5
6	F	205	VAL	2.5
6	F	144	LYS	2.5
6	F	177	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	B	191	ALA	2.5
6	F	166	ALA	2.5
3	C	68	ASN	2.5
6	F	315	HIS	2.5
6	F	263	GLN	2.5
6	F	146	GLN	2.4
4	D	186	ASP	2.4
2	B	203	ILE	2.4
6	F	204	PRO	2.4
2	B	192	LEU	2.3
6	F	286	ALA	2.3
6	F	226	MET	2.3
3	C	74	ALA	2.3
3	C	14	MET	2.3
2	B	242	ASP	2.3
2	B	243	TYR	2.3
3	C	18	ALA	2.3
6	F	221	GLY	2.3
6	F	179	PRO	2.2
2	B	237	VAL	2.2
6	F	353	GLN	2.2
6	F	196	LYS	2.2
6	F	167	GLU	2.2
6	F	323	LEU	2.2
6	F	165	ALA	2.1
2	B	235	TYR	2.1
6	F	256	THR	2.1
4	D	144	ASN	2.1
6	F	163	ILE	2.0
2	B	240	ALA	2.0
6	F	268	GLU	2.0
5	E	75	LYS	2.0
6	F	191	LYS	2.0
6	F	211	THR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
8	GOL	F	504	6/6	0.65	0.25	83,93,96,99	0
8	GOL	F	503	6/6	0.80	0.18	72,75,79,82	0
8	GOL	D	301	6/6	0.85	0.18	59,70,73,79	0
8	GOL	F	502	6/6	0.89	0.16	41,65,71,79	0
8	GOL	F	501	6/6	0.92	0.12	53,72,88,89	0
7	MG	C	101	1/1	0.99	0.05	68,68,68,68	0

6.5 Other polymers [i](#)

There are no such residues in this entry.