



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 25, 2026 – 11:26 AM EDT

PDB ID : 2W62 / pdb_00002w62
Title : Saccharomyces cerevisiae Gas2p in complex with laminaripentaose
Authors : Schuettelkopf, A.W.; Hurtado-Guerrero, R.; van Aalten, D.M.F.
Deposited on : 2008-12-16
Resolution : 1.85 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.85 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3428 (1.86-1.86)
Clashscore	190562	3579 (1.86-1.86)
Ramachandran outliers	187476	3553 (1.86-1.86)
Sidechain outliers	187428	3553 (1.86-1.86)
RSRZ outliers	180081	3429 (1.86-1.86)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	555	11% 71% 7% 21%
2	B	5	100%
2	C	5	60% 40%

2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 3930 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called GLYCOLIPID-ANCHORED SURFACE PROTEIN 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	436	Total	C	N	O	S	0	6	0
			3499	2234	558	684	23			

- Molecule 2 is an oligosaccharide called beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	B	5	Total	C	O	0	0	0
			56	30	26			
2	C	5	Total	C	O	0	0	0
			56	30	26			

- Molecule 3 is 1,4-BUTANEDIOL (CCD ID: BU1) (formula: C₄H₁₀O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			6	4	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	313	Total	O	0	0
			313	313		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: GLYCOLIPID-ANCHORED SURFACE PROTEIN 2



• Molecule 2: beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose



• Molecule 2: beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose-(1-3)-beta-D-glucopyranose



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.04Å 70.84Å 149.15Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	23.00 – 1.85 23.00 – 1.85	Depositor EDS
% Data completeness (in resolution range)	99.5 (23.00-1.85) 99.4 (23.00-1.85)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.11 (at 1.85Å)	Xtriage
Refinement program	REFMAC 5.4.0069	Depositor
R, R_{free}	0.183 , 0.210 0.188 , 0.213	Depositor DCC
R_{free} test set	555 reflections (1.21%)	wwPDB-VP
Wilson B-factor (Å ²)	28.2	Xtriage
Anisotropy	0.228	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	3930	wwPDB-VP
Average B, all atoms (Å ²)	42.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.27% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: BGC, BU1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.67	0/3601	0.77	0/4869

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3499	0	3338	25	0
2	B	56	0	48	1	0
2	C	56	0	48	0	0
3	A	6	0	10	0	0
4	A	313	0	0	2	0
All	All	3930	0	3444	25	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (25) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:308:MET:HE2	1:A:310:PHE:H	1.23	1.00
1:A:29:SER:OG	1:A:31[B]:GLU:HG2	1.78	0.82
1:A:60:GLN:H	1:A:308:MET:HE1	1.49	0.77
1:A:220:MET:HG2	4:A:2168:HOH:O	1.91	0.71
1:A:42:LYS:HE3	1:A:348:ILE:HD11	1.72	0.70
1:A:392:CYS:O	1:A:396:ILE:HG12	1.93	0.69
1:A:433:THR:OG1	1:A:435:GLU:HG3	1.94	0.67
1:A:235:LYS:HE3	4:A:2176:HOH:O	2.01	0.60
1:A:191:LYS:HD2	1:A:228:TYR:CE2	2.39	0.57
1:A:149:VAL:O	1:A:153:GLU:HG2	2.07	0.55
1:A:282:ARG:HA	1:A:283:PRO:C	2.34	0.53
1:A:29:SER:HG	1:A:31[B]:GLU:HG2	1.73	0.51
1:A:395:GLU:O	1:A:478:GLU:HG2	2.12	0.49
1:A:220:MET:N	1:A:220:MET:HE2	2.28	0.48
1:A:139[B]:SER:OG	2:B:3:BGC:H2	2.14	0.47
1:A:74:ALA:C	1:A:75:PHE:HD1	2.22	0.47
1:A:400:GLU:HB2	1:A:474:TYR:CE1	2.51	0.46
1:A:478:GLU:HA	1:A:481:GLN:HG2	1.97	0.45
1:A:415:PHE:CZ	1:A:448:LEU:HG	2.51	0.45
1:A:152:PHE:CE2	1:A:156:LYS:HE2	2.52	0.44
1:A:246:TRP:CG	1:A:288:GLU:HB3	2.52	0.44
1:A:392:CYS:HA	1:A:395:GLU:HG3	2.00	0.43
1:A:191:LYS:CD	1:A:228:TYR:CE2	3.05	0.40
1:A:31[B]:GLU:HG2	1:A:31[B]:GLU:H	1.70	0.40
1:A:389:LYS:O	1:A:392:CYS:HB2	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	433/555 (78%)	429 (99%)	3 (1%)	1 (0%)	43 31

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	177	VAL

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	388/485 (80%)	380 (98%)	8 (2%)	47 33

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	61	LEU
1	A	168	LEU
1	A	312	GLU
1	A	363	GLU
1	A	365	VAL
1	A	395	GLU
1	A	471	LYS
1	A	489	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	62	GLN
1	A	203	HIS
1	A	205	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

10 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
2	BGC	B	1	2	12,12,12	1.01	0	17,17,17	0.92	2 (11%)
2	BGC	B	2	2	11,11,12	0.98	0	15,15,17	1.06	2 (13%)
2	BGC	B	3	2	11,11,12	0.82	0	15,15,17	0.86	0
2	BGC	B	4	2	11,11,12	1.01	1 (9%)	15,15,17	1.58	1 (6%)
2	BGC	B	5	2	11,11,12	0.47	0	15,15,17	1.06	2 (13%)
2	BGC	C	1	2	12,12,12	0.86	0	17,17,17	0.53	0
2	BGC	C	2	2	11,11,12	0.68	0	15,15,17	1.14	1 (6%)
2	BGC	C	3	2	11,11,12	0.99	0	15,15,17	1.05	0
2	BGC	C	4	2	11,11,12	0.81	0	15,15,17	0.64	0
2	BGC	C	5	2	11,11,12	1.35	2 (18%)	15,15,17	1.68	3 (20%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	B	1	2	-	0/2/22/22	0/1/1/1
2	BGC	B	2	2	-	0/2/19/22	0/1/1/1
2	BGC	B	3	2	-	0/2/19/22	0/1/1/1
2	BGC	B	4	2	-	0/2/19/22	0/1/1/1
2	BGC	B	5	2	-	0/2/19/22	0/1/1/1
2	BGC	C	1	2	-	0/2/22/22	0/1/1/1
2	BGC	C	2	2	-	0/2/19/22	0/1/1/1
2	BGC	C	3	2	-	0/2/19/22	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	C	4	2	-	0/2/19/22	0/1/1/1
2	BGC	C	5	2	-	0/2/19/22	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	C	5	BGC	C2-C3	3.39	1.57	1.52
2	B	4	BGC	O5-C5	-2.17	1.39	1.43
2	C	5	BGC	O2-C2	2.02	1.47	1.43

All (11) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	4	BGC	O3-C3-C2	-5.55	98.73	110.05
2	C	5	BGC	C1-O5-C5	3.09	116.33	112.19
2	C	2	BGC	O3-C3-C2	-3.03	103.86	110.05
2	C	5	BGC	O2-C2-C3	-2.95	104.04	110.15
2	C	5	BGC	O2-C2-C1	2.83	115.69	109.22
2	B	2	BGC	C1-C2-C3	-2.23	106.40	109.64
2	B	2	BGC	O3-C3-C4	-2.22	105.15	110.38
2	B	5	BGC	C1-O5-C5	2.14	115.05	112.19
2	B	1	BGC	O5-C5-C4	-2.13	105.86	109.70
2	B	1	BGC	O3-C3-C4	-2.07	105.50	110.38
2	B	5	BGC	C3-C4-C5	2.06	113.97	110.23

There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	BGC	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	BU1	A	1500	-	5,5,5	0.67	0	4,4,4	0.37	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	BU1	A	1500	-	-	1/3/3/3	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1500	BU1	O5-C1-C2-C3

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	436/555 (78%)	1.18	62 (14%) 6 6	24, 40, 53, 73	11 (2%)

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	414	TYR	9.7
1	A	346	LYS	8.1
1	A	412	GLU	5.9
1	A	459	ILE	5.2
1	A	458	LYS	5.2
1	A	482	PRO	5.1
1	A	74	ALA	4.9
1	A	404	PHE	4.9
1	A	489	CYS	4.4
1	A	413	GLU	4.3
1	A	73	GLY	4.0
1	A	401	ILE	3.8
1	A	75	PHE	3.8
1	A	474	TYR	3.7
1	A	468	LEU	3.6
1	A	295	ASN	3.5
1	A	452	LEU	3.6
1	A	481	GLN	3.4
1	A	480	LEU	3.4
1	A	415	PHE	3.1
1	A	461	ALA	3.1
1	A	355	THR	3.1
1	A	457	CYS	3.0
1	A	467	PRO	2.9
1	A	348	ILE	2.9
1	A	466	CYS	2.9
1	A	396	ILE	2.8

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	393	LEU	2.8
1	A	475	PHE	2.7
1	A	411	TYR	2.7
1	A	477	LEU	2.7
1	A	473	VAL	2.7
1	A	252	TYR	2.6
1	A	370	ILE	2.6
1	A	434	GLY	2.6
1	A	460	GLY	2.6
1	A	397	LEU	2.6
1	A	435	GLU	2.5
1	A	439	PHE	2.5
1	A	402	VAL	2.5
1	A	409	GLY	2.4
1	A	313	GLU	2.4
1	A	455	LEU	2.3
1	A	373	GLY	2.3
1	A	417	TYR	2.3
1	A	291	ALA	2.3
1	A	354	LEU	2.3
1	A	450	LEU	2.3
1	A	428	LEU	2.2
1	A	410	LYS	2.2
1	A	436	TYR	2.1
1	A	372	VAL	2.1
1	A	62	GLN	2.1
1	A	395	GLU	2.1
1	A	465	HIS	2.1
1	A	398	PRO	2.1
1	A	424	CYS	2.1
1	A	429	ALA	2.1
1	A	369	HIS	2.0
1	A	437	GLY	2.0
1	A	448	LEU	2.0
1	A	403	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ


There are no non-standard protein/DNA/RNA residues in this entry.


6.3 Carbohydrates [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	BGC	B	5	11/12	0.74	0.17	54,59,62,65	0
2	BGC	C	1	12/12	0.81	0.14	41,46,49,50	0
2	BGC	C	5	11/12	0.85	0.13	43,48,52,56	0
2	BGC	C	4	11/12	0.92	0.10	38,43,44,45	0
2	BGC	C	2	11/12	0.92	0.10	40,40,41,41	0
2	BGC	C	3	11/12	0.93	0.10	39,41,42,42	0
2	BGC	B	4	11/12	0.94	0.08	39,42,43,46	0
2	BGC	B	1	12/12	0.95	0.08	41,42,47,47	0
2	BGC	B	3	11/12	0.96	0.07	36,38,39,39	0
2	BGC	B	2	11/12	0.96	0.07	39,41,43,44	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	BU1	A	1500	6/6	0.77	0.16	48,52,53,58	0

6.5 Other polymers [i](#)

There are no such residues in this entry.