



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 6, 2026 – 04:08 PM UTC

PDB ID : 2W6G / pdb_00002w6g
Title : Low resolution structures of bovine mitochondrial F1-ATPase during controlled dehydration: Hydration State 3.
Authors : Sanchez-Weatherby, J.; Felisaz, F.; Gobbo, A.; Huet, J.; Ravelli, R.B.G.; Bowler, M.W.; Cipriani, F.
Deposited on : 2008-12-18
Resolution : 6.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 6.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	190562	1210 (8.00-4.00)
Ramachandran outliers	187476	1034 (8.00-4.00)
Sidechain outliers	187428	1000 (8.00-4.00)
RSRZ outliers	180081	1136 (8.00-4.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	553	3% 41% 36% 10% 12%
1	B	553	2% 41% 36% 8% 13%
1	C	553	3% 44% 35% 8% 11%
2	D	528	3% 51% 29% 7% 12%
2	E	528	2% 38% 38% 11% 12%
2	F	528	4% 49% 32% 6% 12%
3	G	298	2% 33% 13% 53%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 22795 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	487	Total	C	N	O	S	0	0	0
			3715	2341	656	706	12			
1	B	479	Total	C	N	O	S	0	0	0
			3656	2303	647	694	12			
1	C	492	Total	C	N	O	S	0	0	0
			3748	2360	661	715	12			

- Molecule 2 is a protein called ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	D	467	Total	C	N	O	S	0	0	0
			3539	2243	601	684	11			
2	E	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			
2	F	466	Total	C	N	O	S	0	0	0
			3530	2238	600	681	11			


- Molecule 3 is a protein called ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	G	140	Total	C	N	O	S	0	0	0
			1077	671	195	205	6			


3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL




• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL






• Molecule 1: ATP SYNTHASE SUBUNIT ALPHA HEART ISOFORM, MITOCHONDRIAL

Chain C: 3% 44% 35% 8% 11%







• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL




• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL








• Molecule 2: ATP SYNTHASE SUBUNIT BETA, MITOCHONDRIAL



• Molecule 3: ATP SYNTHASE SUBUNIT GAMMA, MITOCHONDRIAL





ASP

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	109.35Å 132.93Å 275.42Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	101.53 – 6.00 101.53 – 6.00	Depositor EDS
% Data completeness (in resolution range)	100.0 (101.53-6.00) 65.8 (101.53-6.00)	Depositor EDS
R_{merge}	0.21	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.88 (at 6.18Å)	Xtriage
Refinement program	REFMAC 5.5.0038	Depositor
R, R_{free}	0.330 , (Not available) 0.329 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	156.4	Xtriage
Anisotropy	0.372	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 50.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.36$, $\langle L^2 \rangle = 0.20$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.77	EDS
Total number of atoms	22795	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.56% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.76	0/3766	1.74	68/5080 (1.3%)
1	B	0.77	0/3704	1.76	70/4995 (1.4%)
1	C	0.80	1/3799 (0.0%)	1.80	73/5126 (1.4%)
2	D	0.81	1/3596 (0.0%)	1.78	69/4879 (1.4%)
2	E	0.78	0/3587	1.78	86/4867 (1.8%)
2	F	0.80	1/3587 (0.0%)	1.79	93/4867 (1.9%)
3	G	0.41	0/1083	0.93	2/1448 (0.1%)
All	All	0.77	3/23122 (0.0%)	1.74	461/31262 (1.5%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	96	ASN	CA-CB	5.51	1.62	1.53
2	D	96	ASN	CA-CB	5.26	1.61	1.53
1	C	290	GLY	N-CA	5.04	1.49	1.44

The worst 5 of 461 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	408	ARG	CD-NE-CZ	22.73	156.22	124.40
2	F	282	GLN	CB-CG-CD	15.17	138.39	112.60
2	E	408	ARG	CD-NE-CZ	13.40	143.16	124.40
1	B	351	PHE	CA-CB-CG	11.08	124.88	113.80
1	A	270	ASP	CA-CB-CG	10.88	123.48	112.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3715	0	3813	197	0
1	B	3656	0	3764	165	58
1	C	3748	0	3843	180	56
2	D	3539	0	3593	167	0
2	E	3530	0	3587	234	0
2	F	3530	0	3587	131	2
3	G	1077	0	1139	70	0
All	All	22795	0	23326	1034	58

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 22.

The worst 5 of 1034 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:390:ILE:CD1	3:G:242:MET:HE1	1.54	1.37
2:F:390:ILE:HD11	3:G:242:MET:CE	1.54	1.36
2:F:390:ILE:CD1	3:G:242:MET:CE	2.07	1.29
2:E:390:ILE:HG13	3:G:25:MET:SD	1.75	1.25
1:C:127:ARG:HH12	1:C:255:GLU:HB2	1.00	1.13

The worst 5 of 58 symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:314:ASP:O	1:C:22:SER:C[4_455]	0.42	1.78
1:B:314:ASP:CA	1:C:22:SER:N[4_455]	0.82	1.38
1:B:318:GLY:C	1:C:20:ASP:OD2[4_455]	0.87	1.33
1:B:314:ASP:O	1:C:23:VAL:N[4_455]	0.93	1.27
1:B:314:ASP:OD1	1:C:20:ASP:O[4_455]	0.94	1.26

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	485/553 (88%)	443 (91%)	35 (7%)	7 (1%)	9	40
1	B	475/553 (86%)	427 (90%)	41 (9%)	7 (2%)	8	39
1	C	490/553 (89%)	444 (91%)	38 (8%)	8 (2%)	7	37
2	D	465/528 (88%)	419 (90%)	43 (9%)	3 (1%)	21	59
2	E	464/528 (88%)	407 (88%)	48 (10%)	9 (2%)	6	32
2	F	464/528 (88%)	432 (93%)	30 (6%)	2 (0%)	30	67
3	G	134/298 (45%)	130 (97%)	3 (2%)	1 (1%)	18	56
All	All	2977/3541 (84%)	2702 (91%)	238 (8%)	37 (1%)	10	44

5 of 37 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	407	GLY
2	E	393	MET
1	A	57	SER
1	A	405	GLN
1	A	409	ASP

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	393/444 (88%)	343 (87%)	50 (13%)	4	16
1	B	388/444 (87%)	334 (86%)	54 (14%)	3	14
1	C	397/444 (89%)	359 (90%)	38 (10%)	8	25
2	D	377/417 (90%)	344 (91%)	33 (9%)	9	28
2	E	376/417 (90%)	335 (89%)	41 (11%)	6	21
2	F	376/417 (90%)	343 (91%)	33 (9%)	9	28
3	G	116/251 (46%)	109 (94%)	7 (6%)	17	39

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	2423/2834 (86%)	2167 (89%)	256 (11%)	6 21

5 of 256 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	F	210	ASP
2	F	387	ILE
1	B	474	SER
1	B	444	VAL
2	F	423	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 43 such sidechains are listed below:

Mol	Chain	Res	Type
2	E	442	GLN
2	F	282	GLN
2	F	51	GLN
2	F	198	HIS
2	F	308	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	487/553 (88%)	0.09	16 (3%) 49 44	20, 20, 20, 20	0
1	B	479/553 (86%)	0.09	10 (2%) 63 54	20, 20, 20, 20	0
1	C	492/553 (88%)	0.14	14 (2%) 55 47	20, 20, 20, 20	0
2	D	467/528 (88%)	0.13	18 (3%) 43 41	20, 20, 20, 20	0
2	E	466/528 (88%)	0.13	10 (2%) 63 54	20, 20, 20, 20	0
2	F	466/528 (88%)	0.20	22 (4%) 36 36	20, 20, 20, 20	0
3	G	140/298 (46%)	0.25	6 (4%) 40 39	36, 59, 81, 89	0
All	All	2997/3541 (84%)	0.13	96 (3%) 50 45	20, 20, 20, 89	0

The worst 5 of 96 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	220	GLY	10.5
1	A	234	ALA	7.5
2	F	335	LEU	5.4
1	B	274	GLN	4.9
1	B	489	ILE	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.