



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 03:19 PM UTC

PDB ID : 4W6R / pdb_00004w6r
Title : Crystal Structure of Full-Length Split GFP Mutant D102C Disulfide Dimer,
P 1 Space Group
Authors : Leibly, D.J.; Waldo, G.S.; Yeates, T.O.
Deposited on : 2014-08-20
Resolution : 3.47 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.47 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1083 (3.52-3.44)
Clashscore	190562	1139 (3.52-3.44)
Ramachandran outliers	187476	1111 (3.52-3.44)
Sidechain outliers	187428	1112 (3.52-3.44)
RSRZ outliers	180081	1082 (3.52-3.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	234	9% 55% 31% 6% 8%
1	B	234	7% 56% 29% .. 12%
1	C	234	12% 56% 30% • 12%
1	D	234	7% 53% 24% 5% 17%
1	E	234	15% 65% 24% • 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	234	
1	G	234	
1	H	234	
1	I	234	
1	J	234	
1	K	234	
1	L	234	
1	M	234	
1	N	234	
1	O	234	
1	P	234	

2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 25002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called fluorescent protein D102C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	215	Total	C	N	O	S	0	0	0
			1721	1098	291	327	5			
1	B	205	Total	C	N	O	S	0	0	0
			1649	1052	280	313	4			
1	C	207	Total	C	N	O	S	0	0	0
			1643	1052	277	309	5			
1	D	194	Total	C	N	O	S	0	0	0
			1552	993	261	293	5			
1	E	208	Total	C	N	O	S	0	0	0
			1661	1059	280	317	5			
1	F	201	Total	C	N	O	S	0	0	0
			1600	1020	270	307	3			
1	G	188	Total	C	N	O	S	0	0	0
			1497	949	262	282	4			
1	H	191	Total	C	N	O	S	0	0	0
			1537	980	264	289	4			
1	I	209	Total	C	N	O	S	0	0	0
			1671	1065	285	316	5			
1	J	184	Total	C	N	O	S	0	0	0
			1459	931	247	278	3			
1	K	175	Total	C	N	O	S	0	0	0
			1391	896	229	263	3			
1	L	197	Total	C	N	O	S	0	0	0
			1573	1002	269	298	4			
1	M	168	Total	C	N	O	S	0	0	0
			1352	864	233	251	4			
1	N	174	Total	C	N	O	S	0	0	0
			1387	876	235	273	3			
1	O	213	Total	C	N	O	S	0	0	0
			1677	1065	286	322	4			
1	P	204	Total	C	N	O	S	0	0	0
			1632	1038	281	308	5			


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: fluorescent protein D102C




• Molecule 1: fluorescent protein D102C




• Molecule 1: fluorescent protein D102C







• Molecule 1: fluorescent protein D102C



• Molecule 1: fluorescent protein D102C



• Molecule 1: fluorescent protein D102C



Chain G:

Metabolite list for Chain G (top to bottom): MET, ARG, LYS, GLY, GLU, GLU, L7, F8, V11, V12, P13, I14, L15, G20, G31, E32, G33, D36, I39, G40, K41, L42, S43, L44, K45, F46, L47, A48, K52, L53, P54, V55, P56, G57, I60, V61, THR, THR, LEU, LEU, CRO, V68, Q69, C70, F71, S72, R73, TYR, PRO, D76, H81, D82, F83.

Metabolite list for Chain G (bottom to top): R168, V171, D173, L178, A179, D180, N185, I188, GLY, ASP, PRO, PRO, VAL, ASP, LEU, P196, D197, D198, H199, T200, L201, S202, T203, Q204, T205, T206, L207, S208, D216, E217, M218, V219, L220, L221, E222, Y223, A227, G228, L229, THR, HIS, HIS, HIS, HIS, HIS.

Chain H:

Sequence logo for Chain H. The logo shows the relative frequency of amino acids at each position. The sequence is divided into four segments: 10% (red), 53% (green), 26% (yellow), and 18% (grey). The amino acids are listed below the logo, with red dots indicating specific residues at each position.

Position	Amino Acid
1	Met
2	Arg
3	Lys
4	Gly
5	Glu
6	Glu
7	Leu
8	Phe
9	Thr
10	Gly
11	Val
12	Val
13	Phe
14	Leu
15	Leu
16	Leu
17	Leu
18	Leu
19	Leu
20	Leu
21	Leu
22	Leu
23	Leu
24	Leu
25	Leu
26	Leu
27	Leu
28	Leu
29	Leu
30	Leu
31	Leu
32	Leu
33	Leu
34	Leu
35	Leu
36	Leu
37	Leu
38	Leu
39	Leu
40	Leu
41	Leu
42	Leu
43	Leu
44	Leu
45	Leu
46	Leu
47	Leu
48	Leu
49	Leu
50	Leu
51	Leu
52	Leu
53	Leu
54	Leu
55	Leu
56	Leu
57	Leu
58	Leu
59	Leu
60	Leu
61	Leu
62	Leu
63	Leu
64	Leu
65	Leu
66	Leu
67	Leu
68	Leu
69	Leu
70	Leu
71	Leu
72	Leu
73	Leu
74	Leu
75	Leu
76	Leu
77	Leu
78	Leu
79	Leu
80	Leu
81	Leu
82	Leu
83	Leu
84	Leu
85	Leu
86	Leu
87	Leu
88	Leu
89	Leu
90	Leu
91	Leu
92	Leu
93	Leu
94	Leu
95	Leu
96	Leu
97	Leu
98	Leu
99	Leu
100	Leu
101	Leu
102	Leu
103	Leu
104	Leu
105	Leu
106	Leu
107	Leu
108	Leu
109	Leu
110	Leu
111	Leu
112	Leu
113	Leu
114	Leu
115	Leu
116	Leu
117	Leu
118	Leu
119	Leu
120	Leu
121	Leu
122	Leu
123	Leu
124	Leu
125	Leu
126	Leu
127	Leu
128	Leu
129	Leu
130	Leu
131	Leu
132	Leu
133	Leu
134	Leu
135	Leu
136	Leu
137	Leu
138	Leu
139	Leu
140	Leu
141	Leu
142	Leu
143	Leu
144	Leu
145	Leu
146	Leu
147	Leu
148	Leu
149	Leu
150	Leu
151	Leu
152	Leu
153	Leu
154	Leu
155	Leu
156	Leu
157	Leu
158	Leu
159	Leu
160	Leu
161	Leu
162	Leu
163	Leu
164	Leu
165	Leu
166	Leu
167	Leu
168	Leu
169	Leu
170	Leu
171	Leu
172	Leu
173	Leu
174	Leu
175	Leu
176	Leu
177	Leu
178	Leu
179	Leu
180	Leu
181	Leu
182	Leu
183	Leu
184	Leu
185	Leu
186	Leu
187	Leu
188	Leu
189	Leu
190	Leu
191	Leu
192	Leu
193	Leu
194	Leu
195	Leu
196	Leu
197	Leu
198	Leu
199	Leu
200	Leu
201	Leu
202	Leu
203	Leu
204	Leu
205	Leu
206	Leu
207	Leu
208	Leu
209	Leu
210	Leu
211	Leu
212	Leu


Chain I:

Sequence logo for Chain I, showing amino acid conservation across 228 positions. The y-axis represents information content in bits (0.00 to 0.25). The x-axis shows positions 1 to 228. A color scale at the top indicates conservation levels: 7% (red), 55% (green), 29% (yellow), 5% (orange), and 11% (grey). Red dots above the bars indicate positions with high conservation (bits > 0.15).


Position	Amino Acid	Information Content (bits)
1	GLY	0.01
2	PRO	0.01
3	VAL	0.01
4	ASP	0.01
5	L195	0.01
6	P196	0.01
7	D197	0.01
8	L198	0.01
9	H199	0.01
10	Y200	0.01
11	L201	0.01
12	T205	0.01
13	L206	0.01
14	L207	0.01
15	S208	0.01
16	K209	0.01
17	D210	0.01
18	L211	0.01
19	D216	0.01
20	H217	0.01
21	L220	0.01
22	L221	0.01
23	E222	0.01
24	Y223	0.01
25	V224	0.01
26	T225	0.01
27	A226	0.01
28	ALA	0.01
29	GLY	0.01
30	ILE	0.01
31	THR	0.01
32	HIS	0.01
33	HIS	0.01
34	HIS	0.01
35	HIS	0.01
36	HIS	0.01
37	HIS	0.01
38	HIS	0.01
39	M88	0.01
40	R96	0.01
41	T97	0.01
42	F100	0.01
43	K101	0.01
44	C102	0.01
45	D103	0.01
46	G104	0.01
47	T105	0.01
48	Y106	0.01
49	K107	0.01
50	A110	0.01
51	E111	0.01
52	V112	0.01
53	L113	0.01
54	PHE	0.01
55	GLU	0.01
56	G116	0.01
57	L119	0.01
58	V120	0.01
59	I123	0.01
60	I128	0.01
61	D129	0.01
62	F130	0.01
63	K131	0.01
64	GLU	0.01
65	ASP	0.01
66	GLY	0.01
67	N135	0.01
68	I136	0.01
69	E142	0.01
70	Y143	0.01
71	N144	0.01
72	F145	0.01
73	K149	0.01
74	K156	0.01
75	N159	0.01
76	G160	0.01
77	I161	0.01
78	K162	0.01
79	N170	0.01
80	S175	0.01
81	V176	0.01
82	Q177	0.01
83	L178	0.01
84	G189	0.01
85	ASP	0.01
86	MET	0.01
87	ARG	0.01
88	LYS	0.01
89	G4	0.01
90	V11	0.01
91	V12	0.01
92	P13	0.01
93	L18	0.01
94	D19	0.01
95	G20	0.01
96	D21	0.01
97	V22	0.01
98	N23	0.01
99	F28	0.01
100	V29	0.01
101	R30	0.01
102	G31	0.01
103	E32	0.01
104	G33	0.01
105	D36	0.01
106	T39	0.01
107	G40	0.01
108	K41	0.01
109	L42	0.01
110	S43	0.01
111	L44	0.01
112	T49	0.01
113	K52	0.01
114	L53	0.01
115	P56	0.01
116	W57	0.01
117	P58	0.01
118	V61	0.01
119	T62	0.01
120	T63	0.01
121	L64	0.01
122	GRD	0.01
123	V68	0.01
124	F71	0.01
125	S72	0.01
126	R73	0.01
127	D76	0.01
128	K79	0.01
129	D82	0.01
130	F83	0.01
131	F84	0.01

Chain J:







• Molecule 1: fluorescent protein D102C




• Molecule 1: fluorescent protein D102C



• Molecule 1: fluorescent protein D102C





4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	92.42Å 92.56Å 124.53Å 94.94° 96.17° 102.25°	Depositor
Resolution (Å)	89.88 – 3.47 89.88 – 3.47	Depositor EDS
% Data completeness (in resolution range)	89.4 (89.88-3.47) 89.4 (89.88-3.47)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.09	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.70 (at 3.49Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: DEV_1555)	Depositor
R, R_{free}	0.307 , 0.357 0.324 , 0.342	Depositor DCC
R_{free} test set	4609 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	101.2	Xtriage
Anisotropy	0.575	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.28 , 64.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.46$, $\langle L^2 \rangle = 0.28$	Xtriage
Estimated twinning fraction	0.025 for -k,-h,-l	Xtriage
F_o, F_c correlation	0.85	EDS
Total number of atoms	25002	wwPDB-VP
Average B, all atoms (Å ²)	113.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.95% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	2/1758 (0.1%)	1.24	21/2375 (0.9%)
1	B	0.44	0/1681	1.17	13/2265 (0.6%)
1	C	0.43	0/1675	1.11	10/2258 (0.4%)
1	D	0.50	0/1579	1.25	16/2121 (0.8%)
1	E	0.45	1/1694 (0.1%)	1.09	5/2287 (0.2%)
1	F	0.46	0/1631	1.20	16/2199 (0.7%)
1	G	0.52	1/1524 (0.1%)	1.15	8/2053 (0.4%)
1	H	0.48	1/1567 (0.1%)	1.24	12/2112 (0.6%)
1	I	0.47	0/1705	1.22	19/2302 (0.8%)
1	J	0.50	0/1483	1.22	13/1998 (0.7%)
1	K	0.48	0/1417	1.23	11/1912 (0.6%)
1	L	0.41	0/1605	1.08	4/2164 (0.2%)
1	M	0.41	0/1373	1.27	13/1847 (0.7%)
1	N	0.43	0/1410	1.16	11/1899 (0.6%)
1	O	0.55	0/1712	1.18	10/2314 (0.4%)
1	P	0.44	0/1663	1.11	7/2239 (0.3%)
All	All	0.47	5/25477 (0.0%)	1.18	189/34345 (0.6%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	D	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	203	THR	CA-C	7.16	1.60	1.53
1	G	227	ALA	CA-C	-5.73	1.48	1.53
1	A	203	THR	CA-C	5.47	1.60	1.53
1	A	69	GLN	C-N	-5.34	1.26	1.33

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	E	26	LYS	CA-C	5.30	1.58	1.53

The worst 5 of 189 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	203	THR	N-CA-C	15.90	129.52	110.91
1	E	26	LYS	N-CA-C	15.00	128.46	110.91
1	A	203	THR	N-CA-C	13.75	130.21	111.24
1	H	204	GLN	N-CA-C	13.62	125.85	111.14
1	I	195	LEU	CA-C-N	12.37	135.31	119.84

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	D	111	GLU	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1721	0	1680	105	0
1	B	1649	0	1615	85	0
1	C	1643	0	1611	62	1
1	D	1552	0	1511	60	1
1	E	1661	0	1623	40	0
1	F	1600	0	1557	50	1
1	G	1497	0	1460	86	0
1	H	1537	0	1487	59	0
1	I	1671	0	1633	99	0
1	J	1459	0	1431	59	1
1	K	1391	0	1359	52	0
1	L	1573	0	1532	67	0
1	M	1352	0	1339	48	0
1	N	1387	0	1351	42	0
1	O	1677	0	1616	103	0
1	P	1632	0	1586	46	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	25002	0	24391	1023	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 21.

The worst 5 of 1023 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:I:28:PHE:CE2	1:I:30:ARG:HG3	1.44	1.51
1:A:152:ILE:O	1:A:153:THR:HG22	1.19	1.34
1:D:15:LEU:O	1:D:15:LEU:HD12	1.31	1.30
1:B:106:TYR:CE1	1:B:130:PHE:CE1	2.24	1.24
1:G:44:LEU:O	1:G:219:VAL:HG13	1.36	1.24

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:157:GLN:OE1	1:F:184:GLN:NE2[1_445]	2.14	0.06
1:C:17:GLU:OE2	1:J:50:THR:OG1[1_455]	2.17	0.03

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	207/234 (88%)	195 (94%)	11 (5%)	1 (0%)	24 58
1	B	193/234 (82%)	184 (95%)	8 (4%)	1 (0%)	24 58
1	C	195/234 (83%)	188 (96%)	6 (3%)	1 (0%)	24 58
1	D	174/234 (74%)	161 (92%)	12 (7%)	1 (1%)	21 53
1	E	198/234 (85%)	191 (96%)	6 (3%)	1 (0%)	24 58

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	187/234 (80%)	180 (96%)	6 (3%)	1 (0%)	24	58
1	G	176/234 (75%)	167 (95%)	8 (4%)	1 (1%)	21	53
1	H	175/234 (75%)	169 (97%)	6 (3%)	0	100	100
1	I	199/234 (85%)	192 (96%)	6 (3%)	1 (0%)	24	58
1	J	168/234 (72%)	163 (97%)	5 (3%)	0	100	100
1	K	159/234 (68%)	154 (97%)	4 (2%)	1 (1%)	21	53
1	L	187/234 (80%)	181 (97%)	5 (3%)	1 (0%)	24	58
1	M	152/234 (65%)	149 (98%)	2 (1%)	1 (1%)	18	51
1	N	162/234 (69%)	159 (98%)	3 (2%)	0	100	100
1	O	207/234 (88%)	198 (96%)	7 (3%)	2 (1%)	12	44
1	P	192/234 (82%)	186 (97%)	6 (3%)	0	100	100
All	All	2931/3744 (78%)	2817 (96%)	101 (3%)	13 (0%)	30	62

5 of 13 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	VAL
1	F	61	VAL
1	G	206	ILE
1	D	196	PRO
1	C	204	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	189/204 (93%)	186 (98%)	3 (2%)	55	70
1	B	181/204 (89%)	177 (98%)	4 (2%)	45	65
1	C	178/204 (87%)	175 (98%)	3 (2%)	53	69
1	D	170/204 (83%)	167 (98%)	3 (2%)	51	69
1	E	182/204 (89%)	180 (99%)	2 (1%)	65	74

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	F	175/204 (86%)	169 (97%)	6 (3%)	32	58
1	G	163/204 (80%)	157 (96%)	6 (4%)	30	56
1	H	168/204 (82%)	163 (97%)	5 (3%)	36	60
1	I	183/204 (90%)	178 (97%)	5 (3%)	39	62
1	J	160/204 (78%)	156 (98%)	4 (2%)	42	63
1	K	152/204 (74%)	150 (99%)	2 (1%)	61	72
1	L	171/204 (84%)	165 (96%)	6 (4%)	32	57
1	M	149/204 (73%)	146 (98%)	3 (2%)	48	66
1	N	153/204 (75%)	148 (97%)	5 (3%)	33	58
1	O	179/204 (88%)	178 (99%)	1 (1%)	78	79
1	P	177/204 (87%)	176 (99%)	1 (1%)	78	79
All	All	2730/3264 (84%)	2671 (98%)	59 (2%)	45	65

5 of 59 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	111	GLU
1	N	188	ILE
1	I	159	ASN
1	N	128	ILE
1	M	55	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 41 such sidechains are listed below:

Mol	Chain	Res	Type
1	L	184	GLN
1	O	139	HIS
1	M	158	ASN
1	N	170	ASN
1	P	94	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	215/234 (91%)	0.87	22 (10%) 12 9	50, 80, 148, 324	0
1	B	205/234 (87%)	0.70	16 (7%) 19 13	51, 87, 175, 274	0
1	C	207/234 (88%)	0.91	28 (13%) 7 6	55, 91, 164, 338	0
1	D	194/234 (82%)	0.86	17 (8%) 15 11	57, 92, 169, 237	0
1	E	208/234 (88%)	0.99	34 (16%) 4 4	64, 96, 157, 204	0
1	F	201/234 (85%)	0.69	18 (8%) 15 10	69, 106, 174, 357	0
1	G	188/234 (80%)	0.72	19 (10%) 12 9	52, 103, 181, 284	0
1	H	191/234 (81%)	0.99	24 (12%) 8 6	69, 102, 172, 382	0
1	I	209/234 (89%)	0.73	16 (7%) 19 13	64, 98, 165, 249	0
1	J	184/234 (78%)	0.90	25 (13%) 7 6	51, 95, 166, 270	0
1	K	175/234 (74%)	0.89	16 (9%) 15 10	73, 102, 181, 323	0
1	L	197/234 (84%)	0.77	19 (9%) 13 9	67, 114, 206, 293	0
1	M	168/234 (71%)	0.74	18 (10%) 11 8	75, 119, 203, 375	0
1	N	174/234 (74%)	0.83	19 (10%) 10 8	72, 130, 219, 388	0
1	O	213/234 (91%)	1.03	26 (12%) 8 7	54, 126, 205, 302	0
1	P	204/234 (87%)	0.84	17 (8%) 17 12	73, 134, 217, 464	0
All	All	3133/3744 (83%)	0.84	334 (10%) 11 8	50, 104, 188, 464	0

The worst 5 of 334 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	J	102	CYS	7.6
1	D	196	PRO	6.9
1	M	164	ASN	6.6
1	E	133	ASP	6.6
1	O	224	VAL	6.3

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.