



Full wwPDB X-ray Structure Validation Report ⓘ

Apr 15, 2026 – 01:06 AM UTC

PDB ID : 5W6K / pdb_00005w6k
Title : Structure of mutant Taq Polymerase incorporating unnatural base pairs Z:P
Authors : Singh, I.; Georgiadis, M.M.
Deposited on : 2017-06-16
Resolution : 2.34 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.34 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3031 (2.36-2.32)
Clashscore	190562	3127 (2.36-2.32)
Ramachandran outliers	187476	3095 (2.36-2.32)
Sidechain outliers	187428	3095 (2.36-2.32)
RSRZ outliers	180081	3033 (2.36-2.32)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	540	9% 91% 5% .
2	B	12	75% 25%
3	C	13	85% 8% 8%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9340 atoms, of which 4499 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA polymerase I, thermostable.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	522	Total	C	H	N	O	S	0	1	0
			8335	2624	4204	749	746	12			

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	444	VAL	MET	engineered mutation	UNP P19821
A	527	ALA	PRO	engineered mutation	UNP P19821
A	551	GLU	ASP	engineered mutation	UNP P19821
A	832	VAL	GLU	engineered mutation	UNP P19821


- Molecule 2 is a DNA chain called DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(DOC))-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	12	Total	C	H	N	O	P	0	0	0
			374	114	134	48	67	11			

- Molecule 3 is a DNA chain called DNA (5'-D(P*(1WA)P*GP*GP*CP*GP*CP*CP*GP*T P*GP*GP*TP*C)-3').

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	13	Total	C	H	N	O	P	0	0	0
			417	126	147	51	80	13			

- Molecule 4 is (1R)-1-[6-amino-5-(dihydroxyamino)-2-hydroxypyridin-3-yl]-1,4-anhydro-2-deoxy-5-O-[(S)-hydroxy{[(S)-hydroxy(phosphonooxy)phosphoryl]oxy}phosphoryl]-D-erythro-pentitol (CCD ID: A5J) (formula: C₁₀H₁₆N₃O₁₅P₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	P	0	0
			45	10	14	3	15	3		

- Molecule 5 is MAGNESIUM ION (CCD ID: MG) (formula: Mg) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	2	Total Mg 2 2	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	142	Total O 142 142	0	0
6	B	7	Total O 7 7	0	0
6	C	18	Total O 18 18	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: DNA polymerase I, thermostable



- Molecule 2: DNA (5'-D(*GP*AP*CP*CP*AP*CP*GP*GP*CP*GP*CP*(DOC))-3')



- Molecule 3: DNA (5'-D(P*(1WA)P*GP*GP*CP*GP*CP*CP*GP*TP*GP*GP*TP*C)-3')



4 Data and refinement statistics

Property	Value	Source
Space group	P 31 2 1	Depositor
Cell constants a, b, c, α , β , γ	109.31 Å 109.31 Å 90.87 Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	41.98 – 2.34 41.98 – 2.34	Depositor EDS
% Data completeness (in resolution range)	96.2 (41.98-2.34) 96.3 (41.98-2.34)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.61 (at 2.34 Å)	Xtriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, R_{free}	0.224 , 0.275 0.225 , 0.277	Depositor DCC
R_{free} test set	1229 reflections (4.57%)	wwPDB-VP
Wilson B-factor (Å ²)	29.2	Xtriage
Anisotropy	0.071	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.38 , 40.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	0.035 for -h,-k,l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	9340	wwPDB-VP
Average B, all atoms (Å ²)	44.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.69% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: A5J, 1WA, MG, DOC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.12	0/4218	0.29	0/5713
2	B	0.19	0/249	0.45	0/382
3	C	0.18	0/277	0.41	0/426
All	All	0.13	0/4744	0.31	0/6521

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4131	4204	4204	12	0
2	B	240	134	134	2	0
3	C	270	147	147	2	0
4	A	31	14	0	0	0
5	A	2	0	0	0	0
6	A	142	0	0	0	0
6	B	7	0	0	1	0
6	C	18	0	0	0	0
All	All	4841	4499	4485	13	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 1.

All (13) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:473:GLU:OE1	1:A:476:ARG:NH1	2.32	0.62
2:B:105:DA:N7	6:B:201:HOH:O	2.31	0.58
1:A:466:GLU:OE2	1:A:469:ARG:NH1	2.42	0.53
1:A:587:ARG:NH1	2:B:111:DC:OP2	2.42	0.52
1:A:631:VAL:HG11	1:A:638:ILE:HD11	1.92	0.51
1:A:523:ARG:NH2	1:A:530:GLU:OE2	2.46	0.46
1:A:677:ARG:NH2	3:C:204:1WA:OP1	2.49	0.45
1:A:576:SER:O	3:C:208:DG:H4'	2.17	0.44
1:A:299:TRP:CG	1:A:300:PRO:HA	2.53	0.44
1:A:643:ALA:HB2	1:A:659:ARG:HA	2.01	0.42
1:A:647:PHE:CE1	1:A:658:MET:HB2	2.55	0.42
1:A:721:GLU:HA	1:A:726:ARG:O	2.18	0.42
1:A:673:MET:CE	1:A:681:GLU:OE2	2.68	0.42

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	519/540 (96%)	498 (96%)	20 (4%)	1 (0%)	43	50

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	586	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	425/440 (97%)	419 (99%)	6 (1%)	59 71

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	332	VAL
1	A	411	ARG
1	A	524	GLU
1	A	644	SER
1	A	655	ASP
1	A	657	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	415	ASN
1	A	565	ASN
1	A	582	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the

expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	1WA	C	204	3	16,24,25	1.10	1 (6%)	16,35,38	2.32	3 (18%)
2	DOC	B	112	2,3	16,19,20	1.32	2 (12%)	20,26,29	1.25	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	1WA	C	204	3	-	1/5/21/22	0/2/3/3
2	DOC	B	112	2,3	-	0/7/18/19	0/2/2/2

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	204	1WA	C4-N3	-2.82	1.30	1.35
2	B	112	DOC	C6-N1	-2.65	1.31	1.38
2	B	112	DOC	C5-C4	-2.41	1.37	1.42

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	C	204	1WA	O6-C6-N1	6.14	128.03	115.34
3	C	204	1WA	C1'-N9-C8	-5.49	117.25	124.37
2	B	112	DOC	O4'-C4'-C5'	3.05	114.83	109.34
3	C	204	1WA	N3-C2-N1	2.52	127.94	123.32

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	C	204	1WA	O4'-C1'-N9-C8

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	C	204	1WA	1	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	A5J	A	901	5	30,32,32	1.80	11 (36%)	36,50,50	2.18	9 (25%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	A5J	A	901	5	-	3/23/38/38	0/2/2/2

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	A5J	O4'-C1'	-4.29	1.39	1.44
4	A	901	A5J	PB-O3A	-4.00	1.55	1.59
4	A	901	A5J	C6-C1	-3.65	1.33	1.39
4	A	901	A5J	ON2-N	-2.30	1.19	1.35
4	A	901	A5J	O4'-C4'	-2.27	1.39	1.45
4	A	901	A5J	PA-O3A	-2.13	1.57	1.59
4	A	901	A5J	O2-C2	-2.13	1.19	1.29
4	A	901	A5J	PB-O3B	-2.11	1.57	1.59
4	A	901	A5J	C5-N	-2.10	1.41	1.45

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	901	A5J	C6-C5	2.05	1.43	1.39
4	A	901	A5J	C5-C4	-2.02	1.39	1.42

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	901	A5J	C4-N3-C2	8.53	126.85	116.98
4	A	901	A5J	O2G-PG-O3B	3.47	116.26	104.64
4	A	901	A5J	O3G-PG-O3B	3.40	116.03	104.64
4	A	901	A5J	C2'-C1'-C1	-3.31	109.98	114.72
4	A	901	A5J	C6-C1-C1'	2.96	129.43	120.83
4	A	901	A5J	O3B-PB-O2B	2.54	118.36	110.70
4	A	901	A5J	C3'-C2'-C1'	-2.53	100.36	102.73
4	A	901	A5J	O1B-PB-O2B	-2.51	100.76	112.44
4	A	901	A5J	C5-C6-C1	2.05	122.50	119.51

There are no chirality outliers.


All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	901	A5J	PB-O3B-PG-O2G
4	A	901	A5J	PB-O3B-PG-O3G
4	A	901	A5J	PB-O3A-PA-O2A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	522/540 (96%)	0.67	51 (9%) 13 16	14, 42, 88, 134	1 (0%)
2	B	11/12 (91%)	-0.33	0 100 100	20, 30, 55, 57	0
3	C	12/13 (92%)	-0.51	0 100 100	18, 25, 40, 44	0
All	All	545/565 (96%)	0.62	51 (9%) 14 17	14, 41, 87, 134	1 (0%)

All (51) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	682	LEU	6.3
1	A	395	GLY	6.3
1	A	645	TRP	5.3
1	A	683	ALA	5.1
1	A	656	PRO	5.1
1	A	658	MET	5.1
1	A	685	PRO	5.1
1	A	701	PRO	4.3
1	A	525	ALA	4.3
1	A	526	HIS	4.1
1	A	329	GLY	4.1
1	A	653	ALA	3.9
1	A	411	ARG	3.7
1	A	679	SER	3.7
1	A	647	PHE	3.5
1	A	794	GLU	3.3
1	A	655	ASP	3.3
1	A	657	LEU	3.3
1	A	684	ILE	3.2
1	A	660	ARG	3.2
1	A	481	PRO	3.1
1	A	654	VAL	3.1
1	A	702	LYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	650	PRO	3.0
1	A	677	ARG	2.9
1	A	713	GLU	2.9
1	A	524	GLU	2.8
1	A	680	GLN	2.8
1	A	421	GLU	2.8
1	A	661	ALA	2.8
1	A	487	ARG	2.8
1	A	521	ALA	2.7
1	A	652	GLU	2.7
1	A	649	VAL	2.6
1	A	703	VAL	2.6
1	A	630	ARG	2.6
1	A	503	ILE	2.5
1	A	587	ARG	2.5
1	A	705	ALA	2.4
1	A	681	GLU	2.4
1	A	674	SER	2.4
1	A	651	ARG	2.3
1	A	700	PHE	2.3
1	A	676	HIS	2.2
1	A	296	GLU	2.2
1	A	678	LEU	2.1
1	A	640	THR	2.1
1	A	801	ARG	2.1
1	A	670	LEU	2.1
1	A	331	ARG	2.0
1	A	753	VAL	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	1WA	C	204	22/23	0.92	0.11	30,37,49,52	0
2	DOC	B	112	18/19	0.96	0.07	26,30,35,36	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
4	A5J	A	901	31/31	0.96	0.07	28,32,37,38	0
5	MG	A	903	1/1	0.97	0.11	25,25,25,25	0
5	MG	A	902	1/1	0.98	0.04	32,32,32,32	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around A5J A 901:

2mF_o-DF_c (at 0.7 rmsd) in gray
mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





Electron density around MG A 903:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around MG A 902:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers ⓘ

There are no such residues in this entry.