



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2026 – 08:35 AM UTC

PDB ID : 4W78 / pdb_00004w78
Title : Crystal structure of the ChsH1-ChsH2 complex from Mycobacterium tuberculosis
Authors : Guja, K.E.; Yang, M.; Sampson, N.; Garcia-Diaz, M.
Deposited on : 2014-08-21
Resolution : 1.54 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.54 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R _{free}	180053	1003 (1.54-1.54)
Clashscore	190562	1025 (1.54-1.54)
Ramachandran outliers	187476	1007 (1.54-1.54)
Sidechain outliers	187428	1007 (1.54-1.54)
RSRZ outliers	180081	1002 (1.54-1.54)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	178	6% 89% 6% 5%
1	C	178	13% 96% .
1	E	178	7% 90% . 7%
1	G	178	22% 83% 9% 8%
2	B	127	5% 96% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	D	127	4% 96% • •
2	F	127	9% 93% 6% •
2	H	127	3% 94% • • •

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 10103 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hydratase ChsH1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	169	Total	C	N	O	S	0	1	0
			1287	821	223	237	6			
1	C	178	Total	C	N	O	S	0	1	0
			1322	843	225	247	7			
1	E	165	Total	C	N	O	S	0	1	0
			1259	804	219	230	6			
1	G	163	Total	C	N	O	S	0	2	0
			1204	769	210	218	7			

- Molecule 2 is a protein called Hydratase ChsH2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	126	Total	C	N	O	S	0	3	0
			974	625	164	184	1			
2	D	126	Total	C	N	O	S	0	1	0
			957	612	159	185	1			
2	F	127	Total	C	N	O	S	0	0	0
			961	613	164	182	2			
2	H	126	Total	C	N	O	S	0	1	0
			956	613	159	182	2			

- Molecule 3 is CADMIUM ION (CCD ID: CD) (formula: Cd).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Cd	0	0
			1	1		
3	C	2	Total	Cd	0	0
			2	2		
3	D	3	Total	Cd	0	0
			3	3		
3	E	4	Total	Cd	0	0
			4	4		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	F	1	Total 1	Cd 1	0	0
3	G	1	Total 1	Cd 1	0	0
3	H	1	Total 1	Cd 1	0	0

- Molecule 4 is CHLORIDE ION (CCD ID: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	B	1	Total 1	Cl 1	0	0
4	D	1	Total 1	Cl 1	0	0
4	F	1	Total 1	Cl 1	0	0
4	H	1	Total 1	Cl 1	0	0

- Molecule 5 is SODIUM ION (CCD ID: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	D	1	Total 1	Na 1	0	0

- Molecule 6 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	E	1	Total 1	Ca 1	0	0
6	G	1	Total 1	Ca 1	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	190	Total 190	O 190	0	0
7	B	143	Total 143	O 143	0	0
7	C	150	Total 150	O 150	0	0

Continued on next page...



Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	D	132	Total 132	O 132	0	0
7	E	180	Total 180	O 180	0	0
7	F	133	Total 133	O 133	0	0
7	G	126	Total 126	O 126	0	0
7	H	109	Total 109	O 109	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hydratase ChsH1





- Molecule 1: Hydratase ChsH1



- Molecule 1: Hydratase ChsH1




- Molecule 1: Hydratase ChsH1



- Molecule 2: Hydratase ChsH2







- Molecule 2: Hydratase ChsH2



- Molecule 2: Hydratase ChsH2



- Molecule 2: Hydratase ChsH2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	51.78Å 134.10Å 90.42Å 90.00° 91.58° 90.00°	Depositor
Resolution (Å)	37.60 – 1.54 37.60 – 1.54	Depositor EDS
% Data completeness (in resolution range)	99.6 (37.60-1.54) 96.3 (37.60-1.54)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.53 (at 1.54Å)	Xtriage
Refinement program	PHENIX (phenix.refine: dev_1391)	Depositor
R, R_{free}	0.177 , 0.205 0.178 , 0.206	Depositor DCC
R_{free} test set	2001 reflections (1.10%)	wwPDB-VP
Wilson B-factor (Å ²)	16.1	Xtriage
Anisotropy	0.316	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 42.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Estimated twinning fraction	0.000 for h,-k,-l	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	10103	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.10 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.6419e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, NA, CL, CD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/1323	0.71	0/1808
1	C	0.77	2/1360 (0.1%)	0.79	3/1862 (0.2%)
1	E	0.44	0/1295	0.69	0/1770
1	G	0.43	0/1243	0.78	0/1704
2	B	0.42	0/1001	0.69	0/1367
2	D	0.41	0/978	0.75	0/1338
2	F	0.72	1/979 (0.1%)	0.81	2/1337 (0.1%)
2	H	0.97	3/977 (0.3%)	0.76	2/1336 (0.1%)
All	All	0.60	6/9156 (0.1%)	0.75	7/12522 (0.1%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	25	ALA	C-N	22.35	1.61	1.33
2	H	68	GLY	C-N	-21.47	1.10	1.34
2	F	1	MET	C-N	-18.23	1.08	1.33
2	H	69	PRO	N-CD	15.57	1.69	1.47
1	C	26	ARG	C-N	8.92	1.47	1.33
2	H	69	PRO	C-N	-8.34	1.21	1.33

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	26	ARG	O-C-N	10.82	133.34	122.09
2	F	1	MET	O-C-N	-8.20	109.89	123.00
2	H	68	GLY	CA-C-N	7.36	127.53	119.87
2	H	68	GLY	C-N-CA	7.36	127.53	119.87
1	C	25	ALA	CA-C-N	-7.16	110.87	120.54
1	C	25	ALA	C-N-CA	-7.16	110.87	120.54
2	F	3	VAL	N-CA-C	5.62	116.39	108.36

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1287	0	1266	10	0
1	C	1322	0	1273	6	0
1	E	1259	0	1236	3	0
1	G	1204	0	1132	20	0
2	B	974	0	996	3	0
2	D	957	0	959	6	0
2	F	961	0	962	13	0
2	H	956	0	966	7	0
3	B	1	0	0	0	0
3	C	2	0	0	0	0
3	D	3	0	0	0	0
3	E	4	0	0	0	0
3	F	1	0	0	0	0
3	G	1	0	0	0	0
3	H	1	0	0	0	0
4	B	1	0	0	1	0
4	D	1	0	0	0	0
4	F	1	0	0	0	0
4	H	1	0	0	0	0
5	D	1	0	0	0	0
6	E	1	0	0	0	0
6	G	1	0	0	0	0
7	A	190	0	0	1	0
7	B	143	0	0	2	0
7	C	150	0	0	3	0
7	D	132	0	0	1	0
7	E	180	0	0	1	0
7	F	133	0	0	0	0
7	G	126	0	0	1	0
7	H	109	0	0	0	0
All	All	10103	0	8790	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including

hydrogen atoms). The all-atom clashscore for this structure is 3.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:H:69:PRO:CD	2:H:69:PRO:N	1.69	1.26
2:F:1:MET:SD	1:G:17:ALA:O	2.32	0.88
2:H:68:GLY:C	2:H:69:PRO:CD	2.47	0.86
4:B:202:CL:CL	7:B:441:HOH:O	2.32	0.83
2:H:1:MET:SD	2:H:2:THR:N	2.62	0.73
1:G:33[B]:MET:SD	7:G:337:HOH:O	2.47	0.71
2:F:1:MET:HG2	1:G:19:PRO:O	1.92	0.69
1:A:145:ASN:ND2	7:A:363:HOH:O	2.24	0.69
2:D:32[A]:ASP:OD1	7:D:365:HOH:O	2.11	0.69
1:C:161:ASN:ND2	7:C:424:HOH:O	2.26	0.68
2:F:104:ILE:HD13	2:F:126:MET:HE3	1.79	0.65
1:G:99:TYR:O	1:G:101:GLY:N	2.30	0.65
1:A:84:ASP:HB3	2:D:1:MET:HA	1.80	0.63
2:H:68:GLY:CA	2:H:69:PRO:CD	2.80	0.58
1:E:55:ARG:NH1	7:E:301:HOH:O	2.30	0.58
1:G:135:GLN:CB	1:G:141:GLY:H	2.18	0.57
2:F:1:MET:HB3	1:G:17:ALA:HB1	1.89	0.55
1:A:70:GLN:HB3	2:B:53[B]:LEU:HD21	1.88	0.55
2:F:1:MET:CG	1:G:17:ALA:O	2.55	0.55
2:F:1:MET:HE3	1:G:19:PRO:HD2	1.89	0.53
2:F:1:MET:CG	1:G:19:PRO:O	2.56	0.52
1:C:33[B]:MET:HE2	2:D:28:ARG:HD2	1.92	0.51
2:F:1:MET:N	1:G:84:ASP:OD2	2.42	0.51
2:F:1:MET:HE2	1:G:19:PRO:O	2.11	0.51
1:G:108:GLU:HA	2:H:77:GLY:HA2	1.94	0.49
1:C:13:GLN:OE1	7:C:301:HOH:O	2.20	0.48
1:A:9:GLU:O	1:A:13:GLN:HG2	2.13	0.48
1:E:132:GLY:HA2	1:E:134:LYS:HG2	1.95	0.48
1:A:77:LEU:HD12	1:E:39:GLU:HG2	1.97	0.46
1:G:33[B]:MET:HE3	2:H:28:ARG:HD2	1.98	0.46
2:F:98:ALA:HB3	2:F:105:THR:HB	1.97	0.45
1:A:107:CYS:SG	1:A:162:TRP:HZ3	2.40	0.44
1:A:84:ASP:CB	2:D:1:MET:HA	2.46	0.43
1:C:33[A]:MET:HE3	2:D:28:ARG:HD2	2.00	0.43
2:F:1:MET:HE3	1:G:19:PRO:CD	2.48	0.43
1:C:179:ASP:OD1	1:C:179:ASP:N	2.52	0.43
2:F:1:MET:CE	1:G:19:PRO:O	2.67	0.42
1:G:14:ILE:HD11	1:G:89:PRO:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:108:GLU:HA	2:B:77:GLY:HA2	2.02	0.41
1:G:33[A]:MET:HE3	2:H:28:ARG:HD2	2.02	0.41
1:A:92:LYS:HG2	2:D:7:VAL:HG11	2.03	0.41
1:A:107:CYS:SG	1:A:162:TRP:CZ3	3.15	0.40
1:G:135:GLN:CB	1:G:141:GLY:N	2.83	0.40
2:F:8:LEU:HG	2:F:66:TRP:CG	2.57	0.40
1:G:135:GLN:CB	1:G:140:GLU:HA	2.51	0.40
2:B:32:ASP:OD1	7:B:418:HOH:O	2.22	0.40
1:C:39:GLU:OE1	7:C:378:HOH:O	2.21	0.40
1:G:75:MET:HE1	1:G:82:PRO:HD3	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/178 (94%)	166 (99%)	2 (1%)	0	100	100
1	C	177/178 (99%)	174 (98%)	3 (2%)	0	100	100
1	E	164/178 (92%)	162 (99%)	2 (1%)	0	100	100
1	G	163/178 (92%)	155 (95%)	6 (4%)	2 (1%)	10	1
2	B	127/127 (100%)	122 (96%)	5 (4%)	0	100	100
2	D	125/127 (98%)	121 (97%)	4 (3%)	0	100	100
2	F	125/127 (98%)	122 (98%)	3 (2%)	0	100	100
2	H	125/127 (98%)	122 (98%)	3 (2%)	0	100	100
All	All	1174/1220 (96%)	1144 (97%)	28 (2%)	2 (0%)	43	25

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	G	100	ILE
1	G	80	VAL

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	132/141 (94%)	132 (100%)	0	100	100
1	C	133/141 (94%)	132 (99%)	1 (1%)	73	51
1	E	128/141 (91%)	127 (99%)	1 (1%)	73	51
1	G	114/141 (81%)	114 (100%)	0	100	100
2	B	107/106 (101%)	105 (98%)	2 (2%)	50	20
2	D	104/106 (98%)	104 (100%)	0	100	100
2	F	103/106 (97%)	102 (99%)	1 (1%)	68	42
2	H	104/106 (98%)	104 (100%)	0	100	100
All	All	925/988 (94%)	920 (100%)	5 (0%)	84	64

All (5) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	46[A]	LYS
2	B	46[B]	LYS
1	C	4	VAL
1	E	80	VAL
2	F	113	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	70	GLN
1	C	109	GLN
1	E	109	GLN
1	E	161	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	G	145	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 20 ligands modelled in this entry, 20 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

The following chains have linkage breaks:

Mol	Chain	Number of breaks
1	C	1
2	H	1
2	F	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	C	25:ALA	C	26:ARG	N	1.61
1	H	68:GLY	C	69:PRO	N	1.10
1	F	1:MET	C	2:THR	N	1.08

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	169/178 (94%)	0.38	11 (6%) 25 31	10, 22, 38, 52	1 (0%)
1	C	178/178 (100%)	0.78	24 (13%) 7 9	8, 27, 51, 92	1 (0%)
1	E	165/178 (92%)	0.35	12 (7%) 21 26	9, 21, 40, 54	1 (0%)
1	G	163/178 (91%)	1.05	39 (23%) 2 2	7, 26, 60, 69	2 (1%)
2	B	126/127 (99%)	0.16	6 (4%) 35 44	8, 20, 36, 49	3 (2%)
2	D	126/127 (99%)	0.17	5 (3%) 42 51	11, 20, 33, 44	1 (0%)
2	F	127/127 (100%)	0.21	12 (9%) 14 18	9, 19, 41, 61	0
2	H	126/127 (99%)	0.29	4 (3%) 50 59	11, 23, 38, 52	1 (0%)
All	All	1180/1220 (96%)	0.46	113 (9%) 13 18	7, 22, 46, 92	10 (0%)

All (113) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	F	1	MET	7.6
1	C	173	PRO	6.8
1	G	132	GLY	6.1
1	G	133	PRO	6.0
1	G	138	LEU	5.4
2	F	2	THR	5.2
2	D	1	MET	5.2
1	G	102	VAL	5.2
1	C	180	LEU	5.1
1	C	78	GLY	5.1
1	G	167	PHE	5.1
1	G	100	ILE	4.9
1	C	171	GLY	4.9
1	G	136	THR	4.6
1	C	80	VAL	4.5
1	C	172	SER	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	F	114	LEU	4.5
1	G	96	ASP	4.5
2	B	101	ASP	4.4
1	C	174	SER	4.4
1	G	7	ILE	4.3
1	C	170	ALA	4.3
1	G	103	VAL	4.2
2	H	126	MET	4.1
1	G	137	ALA	4.0
2	H	1	MET	4.0
1	G	80	VAL	3.9
1	A	170	ALA	3.9
1	E	7	ILE	3.9
1	C	77	LEU	3.8
1	G	79	GLY	3.8
1	C	176	VAL	3.8
1	C	162	TRP	3.7
1	A	4	VAL	3.7
1	G	165	LEU	3.7
2	F	101	ASP	3.6
2	B	1	MET	3.6
1	G	99	TYR	3.5
2	F	102	GLY	3.4
1	G	134	LYS	3.4
1	G	97	ALA	3.4
1	G	78	GLY	3.3
1	G	142	TRP	3.3
2	F	103	LEU	3.3
1	A	171	GLY	3.3
1	C	79	GLY	3.3
1	G	169	PRO	3.2
1	G	139	GLY	3.2
1	G	131	VAL	3.2
1	G	143	PHE	3.1
1	A	80	VAL	3.1
1	C	3	GLY	3.1
1	E	6	ASP	3.1
2	H	114	LEU	3.1
2	B	2	THR	3.0
1	G	77	LEU	3.0
2	F	113	THR	3.0
1	A	107	CYS	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	D	101	ASP	3.0
1	G	163	ARG	2.9
2	D	103	LEU	2.9
2	D	100	ASN	2.9
2	D	126	MET	2.9
1	E	170	ALA	2.8
1	A	162	TRP	2.8
1	C	175	SER	2.8
1	E	80	VAL	2.8
1	G	98	GLY	2.8
2	F	100	ASN	2.8
1	G	101	GLY	2.7
2	B	126	MET	2.7
1	G	168	LYS	2.7
2	F	115	GLY	2.7
2	B	103	LEU	2.6
1	E	93	LEU	2.6
2	F	3	VAL	2.6
1	C	4	VAL	2.5
1	G	162	TRP	2.4
1	G	130	VAL	2.4
1	C	154	ASP	2.4
1	E	142	TRP	2.4
1	G	140	GLU	2.4
2	B	100	ASN	2.3
1	G	87	LEU	2.3
1	A	172	SER	2.3
1	A	137	ALA	2.2
1	A	77	LEU	2.2
2	F	104	ILE	2.2
1	C	137	ALA	2.2
1	G	10	ALA	2.2
1	G	44	ARG	2.2
1	E	154	ASP	2.2
1	E	97	ALA	2.1
1	G	104	ALA	2.1
2	H	4	VAL	2.1
1	E	78	GLY	2.1
1	C	138	LEU	2.1
1	C	7	ILE	2.1
1	C	106	ASN	2.1
1	E	85	ASP	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	G	91	ILE	2.1
2	F	79	ARG	2.1
1	A	9	GLU	2.1
1	G	95	ASP	2.1
1	G	90	ILE	2.1
1	C	81	ARG	2.1
1	A	5	SER	2.1
1	G	83	LYS	2.0
1	E	131	VAL	2.0
1	E	135	GLN	2.0
1	C	178	ASP	2.0
1	C	128	GLY	2.0
1	C	179	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	CD	D	204	1/1	0.29	0.21	187,187,187,187	0
6	CA	G	202	1/1	0.83	0.14	55,55,55,55	0
3	CD	G	201	1/1	0.86	0.11	79,79,79,79	0
3	CD	D	202	1/1	0.86	0.23	94,94,94,94	0
3	CD	E	204	1/1	0.90	0.20	65,65,65,65	0
5	NA	D	205	1/1	0.91	0.08	20,20,20,20	0
6	CA	E	205	1/1	0.92	0.15	35,35,35,35	0
3	CD	C	201	1/1	0.93	0.09	28,28,28,28	0
4	CL	H	202	1/1	0.94	0.09	27,27,27,27	0
3	CD	E	202	1/1	0.95	0.11	33,33,33,33	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CD	C	202	1/1	0.96	0.10	34,34,34,34	0
4	CL	B	202	1/1	0.96	0.07	18,18,18,18	0
3	CD	E	201	1/1	0.97	0.09	20,20,20,20	0
4	CL	D	203	1/1	0.98	0.05	19,19,19,19	0
4	CL	F	202	1/1	0.98	0.05	14,14,14,14	0
3	CD	E	203	1/1	0.99	0.26	37,37,37,37	0
3	CD	H	201	1/1	1.00	0.07	21,21,21,21	0
3	CD	D	201	1/1	1.00	0.03	16,16,16,16	0
3	CD	F	201	1/1	1.00	0.02	12,12,12,12	0
3	CD	B	201	1/1	1.00	0.05	13,13,13,13	0

6.5 Other polymers [i](#)

There are no such residues in this entry.