



wwPDB X-ray Structure Validation Summary Report ⓘ

Apr 25, 2026 – 10:29 PM EDT

PDB ID : 4W7R / pdb_00004w7r
Title : Crystal Structure of Full-Length Split GFP Mutant E124H/K126H Copper Mediated Dimer, P 21 Space Group
Authors : Leibly, D.J.; Waldo, G.S.; Yeates, T.O.
Deposited on : 2014-08-22
Resolution : 1.80 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7662 (1.80-1.80)
Clashscore	190562	8479 (1.80-1.80)
Ramachandran outliers	187476	8391 (1.80-1.80)
Sidechain outliers	187428	8390 (1.80-1.80)
RSRZ outliers	180081	7663 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	229	
1	B	229	
1	C	229	
1	D	229	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	EDO	B	304	-	-	X	-
3	EDO	D	304	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 7625 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called fluorescent protein E124H/K126C.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	226	Total	C	N	O	S	0	0	0
			1810	1149	312	345	4			
1	B	225	Total	C	N	O	S	0	0	0
			1809	1151	311	344	3			
1	C	226	Total	C	N	O	S	0	0	0
			1821	1159	313	346	3			
1	D	226	Total	C	N	O	S	0	0	0
			1814	1154	313	344	3			

- Molecule 2 is COPPER (II) ION (CCD ID: CU) (formula: Cu).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Cu	0	0
			1	1		
2	C	1	Total	Cu	0	0
			1	1		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	C	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		
3	D	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	80	Total 80	O 80	0	0
4	B	84	Total 84	O 84	0	0
4	C	82	Total 82	O 82	0	0
4	D	67	Total 67	O 67	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: fluorescent protein E124H/K126C




- Molecule 1: fluorescent protein E124H/K126C



- Molecule 1: fluorescent protein E124H/K126C



- Molecule 1: fluorescent protein E124H/K126C



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	62.67Å 87.19Å 92.07Å 90.00° 90.01° 90.00°	Depositor
Resolution (Å)	92.07 – 1.80 92.07 – 1.80	Depositor EDS
% Data completeness (in resolution range)	98.6 (92.07-1.80) 98.9 (92.07-1.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.12	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.16 (at 1.80Å)	Xtriage
Refinement program	PHENIX (PHENIX.REFINE: DEV_1555)	Depositor
R, R_{free}	0.223 , 0.253 0.224 , 0.253	Depositor DCC
R_{free} test set	9079 reflections (10.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.3	Xtriage
Anisotropy	1.013	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 24.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	0.458 for h,-k,-l	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7625	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.13% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: EDO, CU, CRO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.63	1/1828 (0.1%)	0.80	1/2472 (0.0%)
1	B	0.62	1/1828 (0.1%)	0.82	2/2472 (0.1%)
1	C	0.58	1/1840 (0.1%)	0.81	1/2488 (0.0%)
1	D	0.65	1/1833 (0.1%)	0.77	0/2479
All	All	0.62	4/7329 (0.1%)	0.80	4/9911 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	1	MET	C-N	-8.02	1.22	1.33
1	B	68	VAL	N-CA	-7.28	1.32	1.46
1	D	3	LYS	N-CA	-5.86	1.39	1.46
1	C	100	PHE	C-N	-5.07	1.26	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	190	ASP	N-CA-C	-7.33	96.96	108.90
1	A	197	ASP	O-C-N	6.37	130.05	122.79
1	B	95	GLU	O-C-N	5.78	130.74	123.23
1	B	190	ASP	N-CA-C	5.73	118.21	110.88

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	227	ALA	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1810	0	1751	23	0
1	B	1809	0	1752	18	0
1	C	1821	0	1768	15	0
1	D	1814	0	1755	35	0
2	A	1	0	0	0	0
2	C	1	0	0	0	0
3	A	12	0	18	5	0
3	B	16	0	24	9	0
3	C	8	0	12	4	0
3	D	20	0	30	9	0
4	A	80	0	0	0	0
4	B	84	0	0	2	0
4	C	82	0	0	1	0
4	D	67	0	0	12	0
All	All	7625	0	7110	91	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 91 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:228:GLY:C	3:B:304:EDO:H11	1.29	1.52
1:B:228:GLY:C	3:B:304:EDO:C1	2.05	1.29
1:D:135:ASN:CB	4:D:465:HOH:O	1.84	1.22
1:D:135:ASN:HB2	4:D:465:HOH:O	1.42	1.10
1:B:228:GLY:N	3:B:303:EDO:H21	1.68	1.06

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	221/229 (96%)	218 (99%)	3 (1%)	0	100	100
1	B	220/229 (96%)	216 (98%)	4 (2%)	0	100	100
1	C	221/229 (96%)	219 (99%)	2 (1%)	0	100	100
1	D	221/229 (96%)	219 (99%)	2 (1%)	0	100	100
All	All	883/916 (96%)	872 (99%)	11 (1%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/197 (98%)	191 (99%)	2 (1%)	68	64
1	B	193/197 (98%)	192 (100%)	1 (0%)	81	80
1	C	195/197 (99%)	193 (99%)	2 (1%)	68	64
1	D	193/197 (98%)	192 (100%)	1 (0%)	81	80
All	All	774/788 (98%)	768 (99%)	6 (1%)	73	70

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	C	9	THR
1	C	123	ILE
1	D	123	ILE
1	A	207	LEU
1	A	123	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	ASN
1	A	164	ASN
1	C	121	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CRO	B	66	1	22,23,24	3.18	7 (31%)	30,32,34	3.16	11 (36%)
1	CRO	D	66	1	22,23,24	3.13	6 (27%)	30,32,34	2.77	8 (26%)
1	CRO	A	66	1	22,23,24	3.15	8 (36%)	30,32,34	3.19	11 (36%)
1	CRO	C	66	1	22,23,24	3.10	6 (27%)	30,32,34	2.99	8 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CRO	B	66	1	-	0/12/31/32	0/2/2/2
1	CRO	D	66	1	-	1/12/31/32	0/2/2/2
1	CRO	A	66	1	-	1/12/31/32	0/2/2/2
1	CRO	C	66	1	-	0/12/31/32	0/2/2/2

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	66	CRO	CB2-CA2	12.22	1.46	1.35
1	B	66	CRO	CB2-CA2	12.10	1.46	1.35
1	A	66	CRO	CB2-CA2	10.57	1.45	1.35
1	D	66	CRO	CB2-CA2	9.94	1.44	1.35
1	D	66	CRO	C2-N3	-6.17	1.25	1.40

The worst 5 of 38 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	66	CRO	O2-C2-CA2	-10.61	124.25	131.02
1	C	66	CRO	O2-C2-CA2	-10.02	124.62	131.02
1	A	66	CRO	O2-C2-CA2	-9.97	124.66	131.02
1	B	66	CRO	O2-C2-CA2	-8.13	125.83	131.02
1	B	66	CRO	CA2-C2-N3	7.80	110.05	103.50

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	66	CRO	N3-C1-CA1-CB1
1	D	66	CRO	N3-C1-CA1-CB1

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	C	66	CRO	3	0

5.5 Carbohydrates

There are no oligosaccharides in this entry.

5.6 Ligand geometry

Of 16 ligands modelled in this entry, 2 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	A	303	-	3,3,3	0.35	0	2,2,2	0.08	0
3	EDO	D	304	-	3,3,3	0.27	0	2,2,2	0.43	0
3	EDO	C	302	-	3,3,3	0.41	0	2,2,2	0.22	0
3	EDO	D	302	-	3,3,3	0.47	0	2,2,2	0.31	0
3	EDO	B	303	-	3,3,3	0.23	0	2,2,2	0.34	0
3	EDO	D	305	-	3,3,3	0.36	0	2,2,2	0.29	0
3	EDO	C	303	-	3,3,3	0.30	0	2,2,2	0.67	0
3	EDO	B	301	-	3,3,3	0.33	0	2,2,2	0.63	0
3	EDO	B	304	-	3,3,3	0.38	0	2,2,2	0.59	0
3	EDO	D	301	-	3,3,3	0.23	0	2,2,2	0.27	0
3	EDO	A	304	-	3,3,3	0.27	0	2,2,2	0.59	0
3	EDO	A	302	-	3,3,3	0.32	0	2,2,2	0.55	0
3	EDO	B	302	-	3,3,3	0.21	0	2,2,2	0.63	0
3	EDO	D	303	-	3,3,3	0.25	0	2,2,2	0.99	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	303	-	-	0/1/1/1	-
3	EDO	D	304	-	-	1/1/1/1	-
3	EDO	C	302	-	-	0/1/1/1	-
3	EDO	D	302	-	-	1/1/1/1	-
3	EDO	B	303	-	-	1/1/1/1	-
3	EDO	D	305	-	-	1/1/1/1	-
3	EDO	C	303	-	-	1/1/1/1	-
3	EDO	B	301	-	-	1/1/1/1	-
3	EDO	B	304	-	-	1/1/1/1	-
3	EDO	D	301	-	-	1/1/1/1	-
3	EDO	A	304	-	-	0/1/1/1	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	A	302	-	-	1/1/1/1	-
3	EDO	B	302	-	-	1/1/1/1	-
3	EDO	D	303	-	-	1/1/1/1	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

5 of 11 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	302	EDO	O1-C1-C2-O2
3	B	303	EDO	O1-C1-C2-O2
3	A	302	EDO	O1-C1-C2-O2
3	B	301	EDO	O1-C1-C2-O2
3	B	304	EDO	O1-C1-C2-O2

There are no ring outliers.

11 monomers are involved in 27 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	D	304	EDO	6	0
3	C	302	EDO	1	0
3	D	302	EDO	1	0
3	B	303	EDO	3	0
3	D	305	EDO	1	0
3	C	303	EDO	3	0
3	B	301	EDO	2	0
3	B	304	EDO	4	0
3	A	304	EDO	3	0
3	A	302	EDO	2	0
3	D	303	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	225/229 (98%)	-0.96	0 100 100	26, 31, 44, 64	0
1	B	224/229 (97%)	-1.00	0 100 100	26, 31, 45, 88	0
1	C	225/229 (98%)	-0.93	0 100 100	27, 32, 44, 88	0
1	D	225/229 (98%)	-0.90	0 100 100	27, 32, 43, 69	0
All	All	899/916 (98%)	-0.95	0 100 100	26, 32, 44, 88	0

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CRO	C	66	22/23	0.98	0.05	26,28,35,36	0
1	CRO	B	66	22/23	0.99	0.04	26,30,39,50	0
1	CRO	A	66	22/23	0.99	0.04	26,26,30,32	0
1	CRO	D	66	22/23	0.99	0.04	30,31,34,34	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	EDO	C	303	4/4	0.96	0.10	49,49,50,57	0
3	EDO	D	302	4/4	0.97	0.08	37,41,44,46	0
3	EDO	D	304	4/4	0.97	0.08	49,52,52,55	0
3	EDO	D	305	4/4	0.97	0.07	41,45,50,57	0
3	EDO	B	304	4/4	0.98	0.06	50,51,55,56	0
3	EDO	C	302	4/4	0.98	0.07	42,42,48,50	0
3	EDO	A	302	4/4	0.98	0.06	35,36,41,41	0
3	EDO	A	303	4/4	0.98	0.06	35,36,42,42	0
3	EDO	D	303	4/4	0.98	0.06	43,45,48,51	0
3	EDO	A	304	4/4	0.98	0.06	50,51,52,56	0
3	EDO	B	303	4/4	0.98	0.09	49,49,57,60	0
3	EDO	B	302	4/4	0.99	0.06	46,50,51,54	0
3	EDO	D	301	4/4	0.99	0.07	43,44,46,52	0
3	EDO	B	301	4/4	0.99	0.06	39,45,47,50	0
2	CU	A	301	1/1	1.00	0.02	30,30,30,30	0
2	CU	C	301	1/1	1.00	0.03	33,33,33,33	0

6.5 Other polymers

There are no such residues in this entry.