



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 6, 2026 – 06:19 PM UTC

PDB ID : 6W7J / pdb_00006w7j
Title : Structure of Tdp1 catalytic domain in complex with inhibitor XZ635p
Authors : Lountos, G.T.; Zhao, X.Z.; Kiselev, E.; Tropea, J.E.; Needle, D.; Pommier, Y.; Burke, T.R.; Waugh, D.S.
Deposited on : 2020-03-19
Resolution : 1.49 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.49 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6779 (1.50-1.46)
Clashscore	190562	7025 (1.50-1.46)
Ramachandran outliers	187476	6917 (1.50-1.46)
Sidechain outliers	187428	6914 (1.50-1.46)
RSRZ outliers	180081	6781 (1.50-1.46)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	461	3% 88% 7% 5%
1	B	461	5% 88% 6% 5%

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 8059 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Tyrosyl-DNA phosphodiesterase 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	437	Total	C	N	O	S	0	19	0
			3640	2356	615	655	14			
1	B	436	Total	C	N	O	S	0	15	0
			3597	2332	607	644	14			

- Molecule 2 is 4-{{2-(2-hydroxyphenyl)imidazo[1,2-a]pyridin-3-yl}amino}benzene-1,2-dicarboxylic acid (CCD ID: TGM) (formula: C₂₁H₁₅N₃O₅) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			29	21	3	5		
2	B	1	Total	C	N	O	0	0
			13	8	1	4		

- Molecule 3 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		
3	B	1	Total	C	O	0	0
			4	2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	467	Total	O	0	0
			467	467		
4	B	297	Total	O	0	0
			297	297		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Tyrosyl-DNA phosphodiesterase 1



- Molecule 1: Tyrosyl-DNA phosphodiesterase 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	50.00Å 105.47Å 193.57Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	46.31 – 1.49 46.31 – 1.49	Depositor EDS
% Data completeness (in resolution range)	98.6 (46.31-1.49) 98.6 (46.31-1.49)	Depositor EDS
R_{merge}	0.04	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.93 (at 1.49Å)	Xtriage
Refinement program	PHENIX 1.16_3549	Depositor
R, R_{free}	0.151 , 0.181 0.152 , 0.181	Depositor DCC
R_{free} test set	8294 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	16.8	Xtriage
Anisotropy	0.540	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 43.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	8059	wwPDB-VP
Average B, all atoms (Å ²)	26.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.94% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: TGM, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.23	0/3758	0.47	0/5105
1	B	0.20	0/3714	0.43	0/5046
All	All	0.21	0/7472	0.45	0/10151

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3640	0	3554	22	0
1	B	3597	0	3526	24	0
2	A	29	0	0	1	0
2	B	13	0	0	0	0
3	A	4	0	6	0	0
3	B	12	0	18	0	0
4	A	467	0	0	4	0
4	B	297	0	0	4	0
All	All	8059	0	7104	46	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:498:MET:HE2	1:B:510:PHE:CE2	2.14	0.83
1:B:380:LYS:NZ	4:B:801:HOH:O	2.18	0.77
1:A:558:LYS:NZ	4:A:801:HOH:O	2.19	0.74
1:B:498:MET:HE2	1:B:510:PHE:CD2	2.25	0.72
1:A:469:LYS:NZ	4:A:803:HOH:O	2.25	0.68
1:B:510:PHE:HE1	1:B:512[B]:VAL:HG22	1.59	0.67
1:B:498:MET:HE1	1:B:572:VAL:HG13	1.78	0.63
1:A:420:MET:HE1	1:A:497:TYR:CD2	2.34	0.62
1:B:498:MET:HE2	1:B:510:PHE:HE2	1.65	0.61
1:A:324:ILE:O	1:A:328[B]:MET:HG3	2.01	0.60
1:A:558:LYS:HB2	1:A:561:ALA:HB3	1.84	0.59
1:B:584:LYS:N	1:B:584:LYS:HD2	2.18	0.57
1:B:506:LYS:NZ	4:B:802:HOH:O	2.18	0.57
1:A:408:GLU:OE1	1:A:476:TYR:OH	2.19	0.52
1:B:510:PHE:CE1	1:B:512[B]:VAL:HG22	2.44	0.50
1:B:377:LYS:HD2	1:B:380:LYS:HE2	1.95	0.48
1:B:509:TRP:HB3	1:B:544:LEU:HD23	1.96	0.47
1:B:519:LYS:NZ	4:B:811:HOH:O	2.42	0.47
1:A:559:PHE:CE1	1:A:571:PRO:HB2	2.50	0.47
1:A:457:GLY:HA3	1:A:588:TRP:CZ2	2.50	0.47
1:B:361[B]:ARG:NH1	1:B:526:GLU:OE2	2.47	0.46
1:A:417:LYS:O	1:A:421[A]:LEU:HG	2.15	0.46
1:B:498:MET:HE1	1:B:572:VAL:CG1	2.44	0.46
1:A:263[A]:HIS:ND1	1:A:514:SER:HB2	2.30	0.46
1:A:186:LYS:NZ	4:A:805:HOH:O	2.30	0.44
1:B:498:MET:CE	1:B:510:PHE:HE2	2.30	0.43
1:B:461:PRO:HB3	1:B:590:TRP:CZ3	2.52	0.43
1:A:324:ILE:HG22	1:A:328[B]:MET:HE2	2.00	0.43
1:B:457:GLY:HA3	1:B:588:TRP:CZ2	2.54	0.43
1:A:557:GLN:C	1:A:558:LYS:HD2	2.44	0.43
1:B:302:TYR:CD1	1:B:347:LEU:HA	2.53	0.43
1:A:457:GLY:HA3	1:A:588:TRP:CE2	2.54	0.43
1:A:263[A]:HIS:NE2	2:A:701:TGM:O01	2.51	0.42
1:B:203:ASN:OD1	1:B:283:ASN:HA	2.20	0.42
1:A:302:TYR:CD1	1:A:347:LEU:HA	2.55	0.42
1:A:318:HIS:HE1	4:A:1176:HOH:O	2.02	0.41
1:A:421[A]:LEU:HD22	1:A:435:VAL:HG11	2.02	0.41
1:A:509:TRP:HB3	1:A:544:LEU:HD23	2.02	0.41
1:B:162:ASN:HB2	4:B:1060:HOH:O	2.20	0.41
1:B:416:PHE:CZ	1:B:420:MET:HE3	2.56	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:319:PHE:CG	1:B:350:THR:HG21	2.55	0.41
1:B:457:GLY:HA3	1:B:588:TRP:CE2	2.55	0.41
1:A:452:GLU:HB2	1:A:456:ALA:HB2	2.02	0.41
1:A:495:LYS:HE3	1:A:516:ASN:OD1	2.20	0.40
1:A:319:PHE:CG	1:A:350:THR:HG21	2.56	0.40
1:B:454:TYR:CD2	1:B:599:ASP:HB3	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	450/461 (98%)	443 (98%)	7 (2%)	0	100	100
1	B	445/461 (96%)	436 (98%)	8 (2%)	1 (0%)	43	22
All	All	895/922 (97%)	879 (98%)	15 (2%)	1 (0%)	48	23

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	411	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	398/400 (100%)	398 (100%)	0	100	100
1	B	393/400 (98%)	393 (100%)	0	100	100
All	All	791/800 (99%)	791 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	214	GLN
1	A	388	ASN
1	A	471	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	EDO	B	702	-	3,3,3	0.44	0	2,2,2	0.37	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	TGM	A	701	-	32,32,32	2.09	10 (31%)	43,46,46	1.54	10 (23%)
2	TGM	B	701	-	13,13,32	1.41	3 (23%)	18,18,46	1.00	0
3	EDO	A	702	-	3,3,3	0.50	0	2,2,2	0.39	0
3	EDO	B	704	-	3,3,3	0.43	0	2,2,2	0.42	0
3	EDO	B	703	-	3,3,3	0.40	0	2,2,2	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	EDO	B	702	-	-	0/1/1/1	-
2	TGM	A	701	-	-	2/16/16/16	0/4/4/4
2	TGM	B	701	-	-	6/8/8/16	0/1/1/4
3	EDO	A	702	-	-	0/1/1/1	-
3	EDO	B	704	-	-	0/1/1/1	-
3	EDO	B	703	-	-	0/1/1/1	-

All (13) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	701	TGM	C09-N08	5.50	1.49	1.37
2	A	701	TGM	C19-N24	-5.12	1.32	1.39
2	A	701	TGM	C11-C10	-3.61	1.42	1.48
2	A	701	TGM	C09-N24	-3.53	1.35	1.39
2	A	701	TGM	C04-C02	2.97	1.56	1.49
2	A	701	TGM	C10-N18	-2.85	1.33	1.38
2	B	701	TGM	C04-C02	2.76	1.55	1.49
2	A	701	TGM	O17-C16	2.64	1.41	1.36
2	A	701	TGM	C07-N08	2.54	1.46	1.41
2	B	701	TGM	C07-N08	2.45	1.46	1.38
2	A	701	TGM	C23-C22	2.34	1.40	1.35
2	A	701	TGM	C26-C27	2.13	1.54	1.49
2	B	701	TGM	C26-C27	2.07	1.54	1.49

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	TGM	C09-C10-N18	-4.10	106.43	111.34
2	A	701	TGM	C10-N18-C19	3.77	108.96	105.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	701	TGM	C20-C19-N24	3.33	121.55	118.41
2	A	701	TGM	C10-C09-N24	2.68	106.78	104.96
2	A	701	TGM	C11-C10-N18	2.67	124.22	119.01
2	A	701	TGM	N24-C19-N18	-2.52	108.99	111.26
2	A	701	TGM	O29-C27-C26	2.30	121.81	115.28
2	A	701	TGM	C09-N24-C19	2.08	108.83	107.26
2	A	701	TGM	C22-C21-C20	-2.04	118.44	121.20
2	A	701	TGM	O29-C27-O28	-2.02	119.01	123.35

There are no chirality outliers.

All (8) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	B	701	TGM	C25-C26-C27-O29
2	B	701	TGM	C25-C26-C27-O28
2	B	701	TGM	O03-C02-C04-C05
2	A	701	TGM	C25-C26-C27-O29
2	B	701	TGM	C04-C26-C27-O29
2	B	701	TGM	O01-C02-C04-C05
2	B	701	TGM	C04-C26-C27-O28
2	A	701	TGM	C25-C26-C27-O28


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	701	TGM	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	437/461 (94%)	-0.20	13 (2%)	52 57	7, 18, 40, 73	19 (4%)
1	B	436/461 (94%)	0.16	21 (4%)	35 39	7, 26, 50, 71	15 (3%)
All	All	873/922 (94%)	-0.02	34 (3%)	43 47	7, 22, 48, 73	34 (3%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	568	ALA	6.1
1	A	566	PRO	5.0
1	B	566	PRO	4.9
1	B	306	ALA	4.5
1	B	162	ASN	4.4
1	A	163	PRO	3.8
1	B	163	PRO	3.8
1	A	162	ASN	3.7
1	B	567	MET	3.7
1	A	427	SER	3.5
1	A	365	SER	3.4
1	A	561	ALA	3.3
1	A	388	ASN	3.2
1	B	427	SER	3.1
1	B	559	PHE	3.1
1	B	309	THR	3.0
1	B	560	PHE	3.0
1	A	567	MET	2.9
1	B	205[A]	CYS	2.8
1	B	365	SER	2.8
1	B	561	ALA	2.6
1	B	558	LYS	2.6
1	B	307	ASP	2.6
1	A	384	SER	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	414	SER	2.4
1	B	568	ALA	2.3
1	A	180	SER	2.3
1	B	387	PRO	2.3
1	B	569	THR	2.2
1	B	607	PRO	2.2
1	B	193	PHE	2.2
1	B	434	SER	2.1
1	A	558	LYS	2.0
1	A	560	PHE	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	TGM	A	701	29/29	0.88	0.14	24,47,58,59	0
3	EDO	B	703	4/4	0.88	0.17	31,39,41,73	0
3	EDO	B	704	4/4	0.91	0.16	28,31,41,58	0
2	TGM	B	701	13/29	0.94	0.09	31,33,36,36	0
3	EDO	A	702	4/4	0.97	0.06	14,15,16,19	0
3	EDO	B	702	4/4	0.97	0.07	18,19,21,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around TGM A 701:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)





6.5 Other polymers [i](#)

There are no such residues in this entry.