



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2026 – 08:15 PM UTC

PDB ID : 2W8D / pdb_00002w8d
Title : Distinct and essential morphogenic functions for wall- and lipo- teichoic acids in *Bacillus subtilis*
Authors : Schirner, K.; Marles-Wright, J.; Lewis, R.J.; Errington, J.
Deposited on : 2009-01-15
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1596 (2.36-2.36)
Clashscore	190562	1663 (2.36-2.36)
Ramachandran outliers	187476	1646 (2.36-2.36)
Sidechain outliers	187428	1646 (2.36-2.36)
RSRZ outliers	180081	1598 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	436	% 80% 15% ...
1	B	436	3% 82% 12% ...

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	PEG	A	1638	-	-	X	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 7060 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called PROCESSED GLYCEROL PHOSPHATE LIPOTEICHOIC ACID SYNTHASE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	420	Total	C	N	O	P	Se	0	0	0
			3400	2169	541	677	1	12			
1	B	421	Total	C	N	O	P	Se	0	0	0
			3405	2172	542	678	1	12			

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		
2	B	1	Total	Mg	0	0
			1	1		

- Molecule 3 is TETRAETHYLENE GLYCOL (CCD ID: PG4) (formula: C₈H₁₈O₅).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			13	8	5		

- Molecule 4 is DI(HYDROXYETHYL)ETHER (CCD ID: PEG) (formula: $C_4H_{10}O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			7	4	3		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	145	Total	O	0	0
			145	145		
5	B	88	Total	O	0	0
			88	88		


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: PROCESSED GLYCEROL PHOSPHATE LIPOTEICHOIC ACID SYNTHASE 2



- Molecule 1: PROCESSED GLYCEROL PHOSPHATE LIPOTEICHOIC ACID SYNTHASE 2



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	56.63Å 54.41Å 140.78Å 90.00° 90.97° 90.00°	Depositor
Resolution (Å)	141.42 – 2.35 140.76 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.9 (141.42-2.35) 99.9 (140.76-2.35)	Depositor EDS
R_{merge}	0.12	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.07 (at 2.34Å)	Xtriage
Refinement program	REFMAC 5.5.0070	Depositor
R, R_{free}	0.170 , 0.235 0.196 , 0.257	Depositor DCC
R_{free} test set	1797 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	31.3	Xtriage
Anisotropy	0.293	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 47.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.015 for k,h,-l 0.015 for -k,-h,-l 0.027 for h,-k,-l	Xtriage
F_o, F_c correlation	0.91	EDS
Total number of atoms	7060	wwPDB-VP
Average B, all atoms (Å ²)	21.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MG, TPO, PEG, PG4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.84	0/3461	0.99	5/4649 (0.1%)
1	B	0.74	0/3466	0.93	4/4656 (0.1%)
All	All	0.79	0/6927	0.96	9/9305 (0.1%)

There are no bond length outliers.

All (9) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	536	VAL	N-CA-C	7.37	119.51	108.23
1	B	557	ILE	N-CA-C	6.33	113.81	107.55
1	A	464	SER	N-CA-C	5.67	117.81	109.24
1	A	536	VAL	N-CA-C	5.48	115.75	107.80
1	B	217	SER	N-CA-C	5.31	117.48	111.11
1	A	328	VAL	CA-C-N	-5.14	113.51	119.47
1	A	328	VAL	C-N-CA	-5.14	113.51	119.47
1	A	622	PHE	N-CA-C	5.08	117.58	109.96
1	B	487	LEU	N-CA-C	-5.01	101.54	109.96

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3400	0	3225	45	0
1	B	3405	0	3230	35	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	13	0	18	2	0
4	A	7	0	10	6	0
5	A	145	0	0	3	0
5	B	88	0	0	1	0
All	All	7060	0	6483	81	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (81) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:304:MSE:HE2	1:B:469:TYR:CD2	1.89	1.07
1:B:499:GLN:HE22	1:B:570:THR:HG21	1.49	0.77
1:A:311:LEU:CD2	4:A:1638:PEG:H11	2.16	0.76
1:B:256:GLN:O	1:B:259:ILE:HG13	1.91	0.71
1:B:304:MSE:HE2	1:B:469:TYR:CG	2.26	0.68
1:A:256:GLN:HE21	1:A:474:GLY:H	1.43	0.67
4:A:1638:PEG:H21	5:A:2134:HOH:O	1.95	0.67
1:A:560:ARG:CD	1:A:560:ARG:O	2.43	0.67
1:B:256:GLN:NE2	1:B:475:ILE:H	1.93	0.66
1:B:405:LYS:HE2	5:B:2057:HOH:O	1.95	0.66
1:A:311:LEU:HD22	4:A:1638:PEG:H11	1.77	0.66
1:A:560:ARG:O	1:A:560:ARG:HD2	1.96	0.66
1:A:256:GLN:HE22	1:A:475:ILE:H	1.47	0.63
1:B:304:MSE:CE	1:B:469:TYR:CD2	2.75	0.61
1:A:256:GLN:NE2	1:A:475:ILE:H	1.98	0.61
1:A:311:LEU:HD23	4:A:1638:PEG:H11	1.83	0.61
1:B:499:GLN:NE2	1:B:570:THR:HG21	2.17	0.58
1:A:499:GLN:HE22	1:A:570:THR:HG21	1.69	0.58
1:A:610:ASN:OD1	4:A:1638:PEG:H12	2.04	0.57
1:A:256:GLN:NE2	1:A:474:GLY:H	2.03	0.56
1:A:560:ARG:CD	1:A:560:ARG:C	2.79	0.56
1:A:347:GLN:HE21	1:A:353:ASN:HD22	1.55	0.54
1:B:392:MSE:HE1	1:B:455:LEU:HD23	1.89	0.54
1:B:256:GLN:HE21	1:B:474:GLY:H	1.55	0.54
1:B:305:GLU:OE1	1:B:405:LYS:HE3	2.07	0.54
1:B:560:ARG:O	1:B:560:ARG:HD2	2.06	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:378:LYS:HE3	1:A:416:GLY:O	2.07	0.53
1:B:560:ARG:O	1:B:560:ARG:CD	2.57	0.53
1:B:560:ARG:NH2	1:B:606:ASP:OD1	2.36	0.53
1:B:484:ALA:O	1:B:487:LEU:O	2.26	0.52
1:A:568:LYS:HD3	1:A:569:TYR:CZ	2.45	0.52
1:A:560:ARG:O	1:A:560:ARG:HD3	2.10	0.52
1:B:337:TYR:HA	1:B:401:PRO:O	2.10	0.52
1:A:329:PRO:HG3	1:A:361:ILE:CD1	2.41	0.51
1:A:250:VAL:HB	1:A:468:MSE:HG2	1.92	0.51
1:A:369:TYR:CE1	1:A:634:LYS:HD2	2.46	0.51
1:B:340:ALA:HA	1:B:363:LYS:O	2.11	0.50
1:B:256:GLN:NE2	1:B:474:GLY:H	2.09	0.50
1:A:422:THR:HG23	1:A:424:PHE:H	1.77	0.49
1:A:313:GLN:H	4:A:1638:PEG:H42	1.76	0.49
1:A:560:ARG:NH2	1:A:606:ASP:OD1	2.41	0.48
1:B:233:VAL:HG13	1:B:234:PRO:HD2	1.94	0.48
1:B:373:ASN:HB2	1:B:376:ASN:HD22	1.80	0.47
1:A:226:TYR:CD2	1:A:604:MSE:HG3	2.49	0.47
1:B:599:LYS:HE2	1:B:599:LYS:HB3	1.70	0.47
1:A:560:ARG:HD3	1:A:560:ARG:HA	1.49	0.47
1:A:570:THR:HG22	1:A:579:THR:OG1	2.14	0.47
1:B:259:ILE:HD12	1:B:260:ILE:N	2.30	0.47
1:B:241:LYS:HB3	1:B:512:VAL:HG22	1.97	0.46
1:A:620:LYS:HD2	1:A:620:LYS:HA	1.72	0.46
1:A:489:LYS:HD3	5:A:2095:HOH:O	2.16	0.46
1:A:419:GLU:HA	1:A:420:GLY:HA2	1.67	0.46
1:A:339:SER:HA	1:A:403:TYR:O	2.16	0.45
1:B:256:GLN:HE22	1:B:475:ILE:H	1.62	0.45
1:B:227:MSE:HE2	1:B:542:LEU:HB3	1.97	0.45
1:A:392:MSE:HE1	1:A:455:LEU:HD23	1.99	0.45
1:B:392:MSE:HB2	1:B:393:PRO:HD3	1.97	0.45
1:A:305:GLU:OE1	1:A:403:TYR:OH	2.24	0.44
1:A:571:LYS:HE2	1:A:574:GLY:HA2	1.98	0.44
1:A:499:GLN:NE2	1:A:570:THR:HG21	2.33	0.44
1:A:329:PRO:HG3	1:A:361:ILE:HD11	2.00	0.44
1:B:249:TYR:HE2	1:B:304:MSE:HE3	1.82	0.43
1:B:304:MSE:HE2	1:B:469:TYR:CE2	2.44	0.43
1:A:329:PRO:CG	1:A:361:ILE:CD1	2.97	0.43
1:B:317:PHE:CE2	1:B:352:ARG:HG2	2.53	0.43
1:A:230:ASN:O	1:A:231:TYR:C	2.61	0.43
1:A:363:LYS:NZ	3:A:1637:PG4:H71	2.33	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:560:ARG:CD	1:B:560:ARG:C	2.92	0.42
1:A:223:VAL:HG11	1:A:608:ILE:HD11	2.02	0.42
1:A:363:LYS:HE3	3:A:1637:PG4:H71	2.02	0.42
1:B:289:PHE:CE1	1:B:502:ARG:HG3	2.55	0.42
1:A:524:VAL:O	1:A:545:GLY:HA3	2.20	0.42
1:B:241:LYS:HD3	1:B:512:VAL:HG22	2.01	0.41
1:A:478:ASN:ND2	5:A:2087:HOH:O	2.50	0.41
1:A:343:HIS:ND1	1:A:407:ILE:HB	2.36	0.41
1:B:260:ILE:HG12	1:B:284:TYR:OH	2.21	0.41
1:B:487:LEU:HD12	1:B:487:LEU:N	2.36	0.41
1:A:374:GLU:H	1:A:374:GLU:HG3	1.71	0.41
1:A:487:LEU:HD11	1:A:500:LEU:HD21	2.03	0.41
1:B:304:MSE:HE1	1:B:526:VAL:HG21	2.02	0.40
1:A:340:ALA:HA	1:A:363:LYS:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	417/436 (96%)	400 (96%)	16 (4%)	1 (0%)	43	52
1	B	418/436 (96%)	399 (96%)	19 (4%)	0	100	100
All	All	835/872 (96%)	799 (96%)	35 (4%)	1 (0%)	48	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	374	GLU

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	373/376 (99%)	352 (94%)	21 (6%)	19	23
1	B	373/376 (99%)	344 (92%)	29 (8%)	11	12
All	All	746/752 (99%)	696 (93%)	50 (7%)	15	17

All (50) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	333	LYS
1	A	374	GLU
1	A	389	LYS
1	A	412	HIS
1	A	419	GLU
1	A	422	THR
1	A	517	VAL
1	A	531	LEU
1	A	536	VAL
1	A	560	ARG
1	A	570	THR
1	A	580	LYS
1	A	585	LEU
1	A	587	GLU
1	A	592	LYS
1	A	609	ILE
1	A	618	GLU
1	A	620	LYS
1	A	623	LYS
1	A	628	SER
1	A	635	HIS
1	B	220	VAL
1	B	227	MSE
1	B	236	ASN
1	B	241	LYS
1	B	252	LEU
1	B	260	ILE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	276	LEU
1	B	304	MSE
1	B	322	GLN
1	B	374	GLU
1	B	396	GLU
1	B	405	LYS
1	B	412	HIS
1	B	447	SER
1	B	458	ASP
1	B	517	VAL
1	B	531	LEU
1	B	536	VAL
1	B	540	ASP
1	B	552	GLU
1	B	560	ARG
1	B	570	THR
1	B	575	LYS
1	B	583	LYS
1	B	585	LEU
1	B	587	GLU
1	B	589	GLU
1	B	623	LYS
1	B	633	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (14) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	230	ASN
1	A	256	GLN
1	A	319	ASN
1	A	347	GLN
1	A	379	ASN
1	A	446	GLN
1	A	478	ASN
1	A	497	ASN
1	B	256	GLN
1	B	319	ASN
1	B	376	ASN
1	B	379	ASN
1	B	439	GLN
1	B	508	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	297	2,1	8,10,11	0.94	0	10,14,16	1.27	1 (10%)
1	TPO	B	297	2,1	8,10,11	1.02	0	10,14,16	1.58	1 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	297	2,1	-	1/9/11/13	-
1	TPO	B	297	2,1	-	1/9/11/13	-

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	297	TPO	OG1-P-O1P	-3.69	96.17	109.33
1	A	297	TPO	OG1-P-O1P	-2.84	99.22	109.33

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	297	TPO	O-C-CA-CB
1	B	297	TPO	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	PG4	A	1637	-	12,12,12	0.61	0	11,11,11	0.53	0
4	PEG	A	1638	-	6,6,6	0.65	0	5,5,5	0.82	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	PG4	A	1637	-	-	5/10/10/10	-
4	PEG	A	1638	-	-	3/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (8) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	1638	PEG	C1-C2-O2-C3
3	A	1637	PG4	O3-C5-C6-O4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
4	A	1638	PEG	O2-C3-C4-O4
3	A	1637	PG4	O1-C1-C2-O2
4	A	1638	PEG	O1-C1-C2-O2
3	A	1637	PG4	C4-C3-O2-C2
3	A	1637	PG4	C3-C4-O3-C5
3	A	1637	PG4	C5-C6-O4-C7

There are no ring outliers.

2 monomers are involved in 8 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1637	PG4	2	0
4	A	1638	PEG	6	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	407/436 (93%)	0.04	3 (0%) 84 87	12, 19, 36, 44	0
1	B	408/436 (93%)	0.58	13 (3%) 50 57	9, 19, 36, 47	0
All	All	815/872 (93%)	0.31	16 (1%) 65 70	9, 19, 36, 47	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	462	ASP	3.0
1	B	215	ALA	2.6
1	B	423	ASP	2.6
1	A	416	GLY	2.4
1	A	226	TYR	2.3
1	B	614	LEU	2.3
1	B	226	TYR	2.3
1	B	236	ASN	2.3
1	B	280	ASN	2.2
1	A	635	HIS	2.2
1	B	220	VAL	2.2
1	B	318	VAL	2.1
1	B	635	HIS	2.1
1	B	223	VAL	2.0
1	B	229	ALA	2.0
1	B	414	PRO	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TPO	B	297	11/12	0.93	0.10	15,18,23,23	0
1	TPO	A	297	11/12	0.98	0.05	14,17,21,21	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	PG4	A	1637	13/13	0.82	0.17	54,64,66,67	0
4	PEG	A	1638	7/7	0.85	0.14	53,56,58,58	0
2	MG	A	1636	1/1	0.96	0.05	14,14,14,14	0
2	MG	B	1636	1/1	0.99	0.13	12,12,12,12	0

6.5 Other polymers [i](#)

There are no such residues in this entry.