



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 01:01 PM UTC

PDB ID : 5W84 / pdb_00005w84
Title : CRYSTAL STRUCTURE OF IRAK-4 WITH A 4,6-DIAMINONICOTINAMIDE INHIBITOR (COMPOUND NUMBER 4)
Authors : Sack, J.S.
Deposited on : 2017-06-21
Resolution : 2.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	2481 (2.90-2.90)
Clashscore	190562	2690 (2.90-2.90)
Ramachandran outliers	187476	2623 (2.90-2.90)
Sidechain outliers	187428	2625 (2.90-2.90)
RSRZ outliers	180081	2481 (2.90-2.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	305	4% 78% 13% 8%
1	B	305	7% 72% 14% 13%

2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 4324 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Interleukin-1 receptor-associated kinase 4.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	282	Total	C	N	O	P	S	0	0	0
			2197	1376	366	438	3	14			
1	B	264	Total	C	N	O	P	S	0	0	0
			2044	1277	344	406	3	14			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	156	GLY	-	expression tag	UNP Q9NWZ3
A	157	ALA	-	expression tag	UNP Q9NWZ3
A	158	MET	-	expression tag	UNP Q9NWZ3
A	159	GLY	-	expression tag	UNP Q9NWZ3
B	156	GLY	-	expression tag	UNP Q9NWZ3
B	157	ALA	-	expression tag	UNP Q9NWZ3
B	158	MET	-	expression tag	UNP Q9NWZ3
B	159	GLY	-	expression tag	UNP Q9NWZ3

- Molecule 2 is 4-(benzylamino)-6-[(2-oxo-2,3-dihydro-1H-indol-5-yl)amino]pyridine-3-carboxamide (CCD ID: 9YY) (formula: C₂₁H₁₉N₅O₂).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			28	21	5	2		
2	B	1	Total	C	N	O	0	0
			28	21	5	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	18	Total	O	0	0
			18	18		
3	B	9	Total	O	0	0
			9	9		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Interleukin-1 receptor-associated kinase 4



- Molecule 1: Interleukin-1 receptor-associated kinase 4



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	88.19Å 125.20Å 141.68Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.82 – 2.90 93.82 – 2.90	Depositor EDS
% Data completeness (in resolution range)	95.5 (93.82-2.90) 95.5 (93.82-2.90)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.91Å)	Xtriage
Refinement program	BUSTER 2.11.7	Depositor
R, R_{free}	0.200 , 0.248 0.208 , 0.266	Depositor DCC
R_{free} test set	813 reflections (4.79%)	wwPDB-VP
Wilson B-factor (Å ²)	67.6	Xtriage
Anisotropy	0.625	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 76.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	4324	wwPDB-VP
Average B, all atoms (Å ²)	83.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 32.68 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 9.0417e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality ⓘ

5.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, SEP, 9YY

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.89	0/2200	1.35	9/2970 (0.3%)
1	B	0.89	2/2043 (0.1%)	1.36	11/2758 (0.4%)
All	All	0.89	2/4243 (0.0%)	1.35	20/5728 (0.3%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	418	MET	SD-CE	-5.48	1.65	1.79
1	B	287	MET	SD-CE	-5.18	1.66	1.79

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	231	ASP	CA-CB-CG	6.06	118.66	112.60
1	B	351	THR	CA-C-N	5.67	129.33	120.82
1	B	351	THR	C-N-CA	5.67	129.33	120.82
1	A	320	ASP	CA-C-N	5.45	127.58	120.28
1	A	320	ASP	C-N-CA	5.45	127.58	120.28
1	B	320	ASP	CA-C-N	5.38	127.48	120.28
1	B	320	ASP	C-N-CA	5.38	127.48	120.28
1	B	398	ASP	CA-CB-CG	5.37	117.97	112.60
1	A	237	MET	N-CA-C	-5.35	103.63	110.53
1	B	231	ASP	CA-C-N	5.28	127.30	120.44
1	B	231	ASP	C-N-CA	5.28	127.30	120.44
1	A	456	GLU	CA-C-N	5.25	127.57	120.38
1	A	456	GLU	C-N-CA	5.25	127.57	120.38
1	B	369	ASP	CA-CB-CG	5.17	117.77	112.60
1	A	293	GLN	CA-C-N	5.10	125.60	119.94
1	A	293	GLN	C-N-CA	5.10	125.60	119.94
1	B	185	ILE	CA-C-N	5.06	127.32	120.38

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	185	ILE	C-N-CA	5.06	127.32	120.38
1	A	185	ILE	CA-C-N	5.05	127.30	120.38
1	A	185	ILE	C-N-CA	5.05	127.30	120.38

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2197	0	2122	14	0
1	B	2044	0	1973	13	0
2	A	28	0	0	0	0
2	B	28	0	0	0	0
3	A	18	0	0	0	0
3	B	9	0	0	0	0
All	All	4324	0	4095	27	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (27) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:352:THR:O	1:A:353:ALA:HB3	1.89	0.71
1:B:284:SER:H	1:B:287:MET:HE3	1.55	0.71
1:A:266:PRO:HG3	1:A:321:GLU:HG3	1.77	0.67
1:B:176:VAL:HG22	1:B:204:TYR:H	1.60	0.66
1:B:266:PRO:HG3	1:B:321:GLU:HG3	1.79	0.63
1:A:234:ILE:HD11	1:A:251:PHE:HB3	1.80	0.63
1:A:265:MET:HG3	1:A:318:LEU:HB3	1.82	0.62
1:A:352:THR:O	1:A:353:ALA:CB	2.51	0.58
1:A:237:MET:O	1:A:238:ALA:HB3	2.05	0.56
1:A:439:GLU:H	1:A:439:GLU:CD	2.16	0.53
1:A:415:ASP:HB3	1:A:418:MET:HE2	1.89	0.53

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:212:VAL:HG22	1:B:261:VAL:HG22	1.92	0.52
1:B:452:GLN:HA	1:B:455:GLN:HE21	1.76	0.49
1:A:300:ASN:HA	1:A:447:ILE:HG21	1.94	0.49
1:B:300:ASN:HA	1:B:447:ILE:HG21	1.94	0.49
1:B:176:VAL:HG11	1:B:205:VAL:HG22	1.94	0.48
1:A:356:ALA:HA	1:A:371:TYR:CD2	2.51	0.45
1:A:232:GLN:O	1:A:236:VAL:HG23	2.16	0.45
1:B:240:CYS:HB3	1:B:301:PHE:HE1	1.81	0.45
1:B:388:ASP:HB3	1:B:391:ARG:HB3	1.98	0.45
1:B:356:ALA:HA	1:B:371:TYR:CD2	2.52	0.45
1:A:168:PHE:HE1	1:A:205:VAL:HG11	1.82	0.44
1:B:176:VAL:CG2	1:B:204:TYR:H	2.29	0.42
1:A:439:GLU:CD	1:A:439:GLU:N	2.77	0.42
1:A:237:MET:O	1:A:238:ALA:CB	2.68	0.42
1:B:265:MET:SD	1:B:326:LYS:HG3	2.60	0.40
1:B:321:GLU:H	1:B:321:GLU:CD	2.30	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	272/305 (89%)	248 (91%)	21 (8%)	3 (1%)	11	36
1	B	254/305 (83%)	240 (94%)	11 (4%)	3 (1%)	10	34
All	All	526/610 (86%)	488 (93%)	32 (6%)	6 (1%)	11	36

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	166	HIS
1	A	228	GLN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	169	SER
1	B	406	GLU
1	A	406	GLU
1	B	441	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	236/260 (91%)	219 (93%)	17 (7%)	13	39
1	B	216/260 (83%)	206 (95%)	10 (5%)	24	57
All	All	452/520 (87%)	425 (94%)	27 (6%)	17	47

All (27) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	169	SER
1	A	186	SER
1	A	233	GLU
1	A	239	LYS
1	A	257	ASP
1	A	277	LEU
1	A	318	LEU
1	A	327	ILE
1	A	329	ASP
1	A	344	MET
1	A	365	THR
1	A	399	ILE
1	A	401	GLU
1	A	434	SER
1	A	438	HIS
1	A	439	GLU
1	A	456	GLU
1	B	233	GLU
1	B	259	CYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	277	LEU
1	B	318	LEU
1	B	327	ILE
1	B	329	ASP
1	B	344	MET
1	B	347	ARG
1	B	401	GLU
1	B	434	SER

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	178	ASN
1	A	297	ASN
1	A	435	GLN
1	A	451	GLN
1	B	206	ASN
1	B	293	GLN
1	B	297	ASN
1	B	419	ASN
1	B	435	GLN
1	B	451	GLN
1	B	455	GLN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	TPO	A	342	1	8,10,11	1.08	0	10,14,16	1.71	2 (20%)
1	SEP	A	346	1	8,9,10	0.85	0	7,12,14	3.48	2 (28%)
1	TPO	B	345	1	8,10,11	1.32	2 (25%)	10,14,16	1.35	2 (20%)
1	SEP	B	346	1	8,9,10	0.81	0	7,12,14	2.69	3 (42%)
1	TPO	B	342	1	8,10,11	0.99	0	10,14,16	1.84	3 (30%)
1	TPO	A	345	1	8,10,11	1.37	2 (25%)	10,14,16	1.09	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	342	1	-	1/9/11/13	-
1	SEP	A	346	1	-	6/6/8/10	-
1	TPO	B	345	1	-	5/9/11/13	-
1	SEP	B	346	1	-	2/6/8/10	-
1	TPO	B	342	1	-	2/9/11/13	-
1	TPO	A	345	1	-	2/9/11/13	-

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	345	TPO	CB-CA	2.65	1.59	1.53
1	B	345	TPO	P-OG1	-2.28	1.55	1.59
1	A	345	TPO	CG2-CB	2.25	1.57	1.51
1	B	345	TPO	CB-CA	2.19	1.58	1.53

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	346	SEP	OG-CB-CA	8.59	116.51	108.14
1	B	346	SEP	OG-CB-CA	6.00	113.98	108.14
1	B	342	TPO	P-OG1-CB	-3.63	113.47	123.33
1	A	342	TPO	P-OG1-CB	-3.58	113.61	123.33
1	B	342	TPO	O2P-P-OG1	3.37	118.99	105.85
1	A	342	TPO	O2P-P-OG1	2.95	117.36	105.85
1	A	346	SEP	O3P-P-OG	2.86	114.13	106.67
1	B	345	TPO	O2P-P-OG1	2.86	116.98	105.85

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	346	SEP	OG-P-O1P	2.58	113.42	106.44
1	B	346	SEP	O2P-P-O1P	-2.28	101.95	110.83
1	B	342	TPO	CG2-CB-CA	2.17	117.48	113.26
1	B	345	TPO	P-OG1-CB	-2.11	117.60	123.33

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	A	345	TPO	N-CA-CB-OG1
1	A	346	SEP	C-CA-CB-OG
1	A	346	SEP	CA-CB-OG-P
1	A	346	SEP	CB-OG-P-O1P
1	B	345	TPO	N-CA-CB-OG1
1	B	346	SEP	N-CA-CB-OG
1	B	346	SEP	C-CA-CB-OG
1	A	346	SEP	CB-OG-P-O2P
1	B	342	TPO	CB-OG1-P-O1P
1	A	346	SEP	N-CA-CB-OG
1	B	345	TPO	CB-OG1-P-O1P
1	A	346	SEP	CB-OG-P-O3P
1	B	345	TPO	CB-OG1-P-O2P
1	B	345	TPO	CB-OG1-P-O3P
1	A	345	TPO	O-C-CA-CB
1	B	342	TPO	C-CA-CB-CG2
1	B	345	TPO	O-C-CA-CB
1	A	342	TPO	CB-OG1-P-O1P

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	9YY	B	4000	-	31,31,31	0.41	0	42,43,43	0.46	0
2	9YY	A	4000	-	31,31,31	0.40	0	42,43,43	0.48	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9YY	B	4000	-	-	0/13/21/21	0/4/4/4
2	9YY	A	4000	-	-	0/13/21/21	0/4/4/4

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	279/305 (91%)	0.25	13 (4%) 36 28	42, 77, 112, 162	0
1	B	261/305 (85%)	0.34	21 (8%) 18 15	46, 76, 145, 167	0
All	All	540/610 (88%)	0.29	34 (6%) 26 20	42, 77, 134, 167	0

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	168	PHE	5.0
1	B	252	SER	4.8
1	A	221	ILE	3.9
1	B	170	PHE	3.8
1	A	187	VAL	3.6
1	B	231	ASP	3.5
1	A	237	MET	3.2
1	A	459	ALA	3.1
1	A	223	THR	3.0
1	A	222	THR	2.9
1	B	173	LEU	2.9
1	B	234	ILE	2.9
1	B	213	LYS	2.7
1	A	399	ILE	2.6
1	B	192	MET	2.5
1	B	343	VAL	2.5
1	A	168	PHE	2.5
1	B	251	PHE	2.5
1	B	169	SER	2.4
1	B	171	TYR	2.4
1	B	186	SER	2.4
1	B	185	ILE	2.4
1	A	175	ASN	2.3
1	B	330	PHE	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	343	VAL	2.3
1	B	332	LEU	2.2
1	A	232	GLN	2.2
1	A	196	GLY	2.2
1	B	205	VAL	2.1
1	A	215	LEU	2.1
1	B	236	VAL	2.0
1	B	459	ALA	2.0
1	B	344	MET	2.0
1	B	264	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	TPO	A	342	11/12	0.81	0.16	110,113,122,122	0
1	TPO	B	342	11/12	0.82	0.15	118,120,125,125	0
1	SEP	A	346	10/11	0.83	0.10	111,116,123,124	0
1	SEP	B	346	10/11	0.84	0.10	116,119,125,125	0
1	TPO	A	345	11/12	0.88	0.10	104,108,110,111	0
1	TPO	B	345	11/12	0.95	0.08	111,112,118,118	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
2	9YY	B	4000	28/28	0.83	0.18	101,106,118,118	0
2	9YY	A	4000	28/28	0.90	0.14	84,90,96,96	0

6.5 Other polymers [i](#)

There are no such residues in this entry.