



wwPDB NMR Structure Validation Summary Report ⓘ

Mar 7, 2026 – 04:31 AM UTC

PDB ID : 5W88 / pdb_00005w88
BMRB ID : 30308
Title : NMR structure of the N-domain of troponin C bound to switch region of troponin I and 3-methyldiphenylamine (peptide mode)
Authors : Cai, F.; Hwang, P.M.; Sykes, B.D.
Deposited on : 2017-06-21

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
BMRB Restraints Analysis : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 91%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	229148	14424
Ramachandran outliers	224038	12848
Sidechain outliers	223484	12823

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	120	35% 39% . 22%

2 Ensemble composition and analysis

This entry contains 8 models. Model 4 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *closest to the average*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:2-A:84, A:149-A:158 (93)	0.52	4

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters. No single-model clusters were found.

Cluster number	Models
1	1, 2, 3, 4, 5
2	6, 7, 8

3 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 1901 atoms, of which 949 are hydrogens and 0 are deuteriums.


- Molecule 1 is a protein called Troponin C, Troponin I.

Mol	Chain	Residues	Atoms						Trace
1	A	120	Total	C	H	N	O	S	0
			1874	582	936	160	187	9	

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	35	SER	CYS	engineered mutation	UNP P63316
A	84	SER	CYS	engineered mutation	UNP P63316
A	136	GLY	-	linker	UNP P63316
A	137	LYS	-	linker	UNP P63316
A	164	GLY	-	expression tag	UNP P19429
A	165	HIS	-	expression tag	UNP P19429

- Molecule 2 is 3-methyl-N-phenylaniline (CCD ID: 9XG) (formula: C₁₃H₁₃N).




Mol	Chain	Residues	Atoms			
2	A	1	Total	C	H	N
			27	13	13	1

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.


- Molecule 1: Troponin C, Troponin I



4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 4. Colouring as in section 4.1 above.

- Molecule 1: Troponin C, Troponin I



5 Refinement protocol and experimental data overview

The models were refined using the following method: *molecular dynamics*.

Of the 80 calculated structures, 8 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
ARIA2	structure calculation	
ARIA2	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	1420
Number of shifts mapped to atoms	1420
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	91%

6 Model quality

6.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: 9XG

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	720	692	692	46±19
2	A	14	13	0	0±0
All	All	5872	5640	4844	370

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 35.

5 of 194 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:44:VAL:O	1:A:48:LEU:HG	0.78	1.78	7	3
1:A:5:TYR:HB3	1:A:82:VAL:HG13	0.76	1.57	7	2
1:A:61:ILE:HG23	1:A:72:VAL:HG23	0.76	1.57	5	5
1:A:20:PHE:CE2	1:A:81:MET:HE2	0.74	2.18	2	3
1:A:20:PHE:CE1	1:A:81:MET:HE2	0.73	2.18	3	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	93/120 (78%)	88±1 (94±1%)	4±1 (5±2%)	1±0 (1±1%)	20	70
All	All	744/960 (78%)	703 (94%)	36 (5%)	5 (1%)	20	70

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	65	ASP	5

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/101 (78%)	69±2 (87±3%)	10±2 (13±3%)	6	47
All	All	632/808 (78%)	552 (87%)	80 (13%)	6	47

5 of 23 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	4	ILE	7
1	A	27	PHE	7
1	A	79	VAL	7
1	A	39	LYS	6
1	A	40	GLU	6

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mol	Type	Chain	Res	Link	Bond lengths		
					Counts	RMSZ	#Z>2
2	9XG	A	201	-	15,15,15	1.63±0.01	3±0 (20±0%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mol	Type	Chain	Res	Link	Bond angles		
					Counts	RMSZ	#Z>2
2	9XG	A	201	-	19,19,19	0.48±0.03	0±0 (0±0%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	9XG	A	201	-	-	0±0,4,4,4	0±0,2,2,2

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
2	A	201	9XG	CAA-CAL	4.02	1.39	1.51	1	8
2	A	201	9XG	CAM-NAK	3.50	1.33	1.40	3	8
2	A	201	9XG	CAN-NAK	3.45	1.33	1.40	5	8

There are no bond-angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 91% for the well-defined parts and 87% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	1420
Number of shifts mapped to atoms	1420
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	1

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction $\pm$ precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	119	-0.13 ± 0.10	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	110	0.09 ± 0.08	None needed (< 0.5 ppm)
$^{13}\text{C}'$	114	-0.40 ± 0.12	None needed (< 0.5 ppm)
^{15}N	111	0.03 ± 0.36	None needed (< 0.5 ppm)

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 91%, i.e. 1110 atoms were assigned a chemical shift out of a possible 1218. 0 out of 15 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	463/467 (99%)	188/190 (99%)	186/186 (100%)	89/91 (98%)
Sidechain	617/692 (89%)	422/448 (94%)	195/226 (86%)	0/18 (0%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	30/59 (51%)	15/29 (52%)	15/30 (50%)	0/0 (—%)
Overall	1110/1218 (91%)	625/667 (94%)	396/442 (90%)	89/109 (82%)

7.1.4 Statistically unusual chemical shifts [i](#)


The following table lists the statistically unusual chemical shifts. These are statistical measures, and large deviations from the mean do not necessarily imply incorrect assignments. Molecules containing paramagnetic centres or hemes are expected to give rise to anomalous chemical shifts.

List Id	Chain	Res	Type	Atom	Shift, ppm	Expected range, ppm	Z-score
1	A	21	LYS	HD3	0.46	0.54 – 2.65	-5.4

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:



8 NMR restraints analysis

8.1 Conformationally restricting restraints

The following table provides the summary of experimentally observed NMR restraints in different categories. Restraints are classified into different categories based on the sequence separation of the atoms involved.

Description	Value
Total distance restraints	1983
Intra-residue ($ i-j =0$)	785
Sequential ($ i-j =1$)	315
Medium range ($ i-j >1$ and $ i-j <5$)	363
Long range ($ i-j \geq 5$)	496
Inter-chain	24
Hydrogen bond restraints	0
Disulfide bond restraints	0
Total dihedral-angle restraints	0
Number of unmapped restraints	0
Number of restraints per residue	16.4
Number of long range restraints per residue ¹	4.1

¹Long range hydrogen bonds and disulfide bonds are counted as long range restraints while calculating the number of long range restraints per residue

8.2 Residual restraint violations

This section provides the overview of the restraint violations analysis. The violations are binned as small, medium and large violations based on its absolute value. Average number of violations per model is calculated by dividing the total number of violations in each bin by the size of the ensemble.

8.2.1 Average number of distance violations per model

Distance violations less than 0.1 Å are not included in the calculation.

Bins (Å)	Average number of violations per model	Max (Å)
0.1-0.2 (Small)	75.2	0.2
0.2-0.5 (Medium)	145.8	0.5
>0.5 (Large)	134.9	8.22

8.2.2 Average number of dihedral-angle violations per model [i](#)

Dihedral-angle violations less than 1° are not included in the calculation. There are no dihedral-angle violations

9 Distance violation analysis ⓘ


9.1 Summary of distance violations ⓘ

The following table shows the summary of distance violations in different restraint categories based on the sequence separation of the atoms involved. Each category is further sub-divided into three sub-categories based on the atoms involved. Violations less than 0.1 Å are not included in the statistics.

Restrains type	Count	% ¹	Violated ³			Consistently Violated ⁴		
			Count	% ²	% ¹	Count	% ²	% ¹
Intra-residue (i-j =0)	785	39.6	182	23.2	9.2	41	5.2	2.1
Backbone-Backbone	74	3.7	2	2.7	0.1	1	1.4	0.1
Backbone-Sidechain	406	20.5	112	27.6	5.6	20	4.9	1.0
Sidechain-Sidechain	305	15.4	68	22.3	3.4	20	6.6	1.0
Sequential (i-j =1)	315	15.9	92	29.2	4.6	16	5.1	0.8
Backbone-Backbone	99	5.0	3	3.0	0.2	0	0.0	0.0
Backbone-Sidechain	140	7.1	43	30.7	2.2	6	4.3	0.3
Sidechain-Sidechain	76	3.8	46	60.5	2.3	10	13.2	0.5
Medium range (i-j >1 & i-j <5)	363	18.3	152	41.9	7.7	33	9.1	1.7
Backbone-Backbone	27	1.4	2	7.4	0.1	0	0.0	0.0
Backbone-Sidechain	187	9.4	67	35.8	3.4	14	7.5	0.7
Sidechain-Sidechain	149	7.5	83	55.7	4.2	19	12.8	1.0
Long range (i-j ≥5)	496	25.0	220	44.4	11.1	46	9.3	2.3
Backbone-Backbone	7	0.4	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	171	8.6	69	40.4	3.5	14	8.2	0.7
Sidechain-Sidechain	318	16.0	151	47.5	7.6	32	10.1	1.6
Inter-chain	24	1.2	7	29.2	0.4	1	4.2	0.1
Backbone-Backbone	0	0.0	0	0.0	0.0	0	0.0	0.0
Backbone-Sidechain	1	0.1	0	0.0	0.0	0	0.0	0.0
Sidechain-Sidechain	23	1.2	7	30.4	0.4	1	4.3	0.1
Hydrogen bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Disulfide bond	0	0.0	0	0.0	0.0	0	0.0	0.0
Total	1983	100.0	653	32.9	32.9	137	6.9	6.9
Backbone-Backbone	207	10.4	7	3.4	0.4	1	0.5	0.1
Backbone-Sidechain	905	45.6	291	32.2	14.7	54	6.0	2.7
Sidechain-Sidechain	871	43.9	355	40.8	17.9	82	9.4	4.1

¹ percentage calculated with respect to the total number of distance restraints, ² percentage calculated with respect to the number of restraints in a particular restraint category, ³ violated in at least one model, ⁴ violated in all the models

9.1.1 Bar chart : Distribution of distance restraints and violations [i](#)



Violated and consistently violated restraints are shown using different hatch patterns in their respective categories. The hydrogen bonds and disulfied bonds are counted in their appropriate category on the x-axis

9.2 Distance violation statistics for each model [i](#)


The following table provides the distance violation statistics for each model in the ensemble. Violations less than 0.1 Å are not included in the statistics.

Model ID	Number of violations						Mean (Å)	Max (Å)	SD ⁶ (Å)	Median (Å)
	IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total				
1	96	46	75	124	3	344	0.49	3.35	0.42	0.38
2	100	46	79	117	3	345	0.49	2.72	0.38	0.4
3	105	39	88	115	2	349	0.49	3.5	0.44	0.35
4	108	38	86	115	2	349	0.51	3.67	0.44	0.38
5	109	49	81	103	2	344	0.47	2.77	0.38	0.36
6	102	56	100	126	3	387	0.66	8.22	0.77	0.43
7	100	49	100	131	2	382	0.65	6.99	0.74	0.42
8	104	49	94	132	2	381	0.66	7.7	0.74	0.42

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶Standard deviation

9.2.1 Bar graph : Distance Violation statistics for each model [i](#)



The mean(dot),median(x) and the standard deviation are shown in blue with respect to the y axis on the right

9.3 Distance violation statistics for the ensemble [i](#)

Violation analysis may find that some restraints are violated in few models and some are violated in most of models. The following table provides this information as number of violated restraints for a given fraction of the ensemble. In total, 1330(IR:603, SQ:223, MR:211, LR:276, IC:17) restraints are not violated in the ensemble.

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
25	22	24	34	3	108	1	12.5
26	10	20	24	1	81	2	25.0
26	12	21	34	2	95	3	37.5
20	5	14	27	0	66	4	50.0
19	13	20	26	0	78	5	62.5
10	8	7	15	0	40	6	75.0

Continued on next page...


Continued from previous page...

Number of violated restraints						Fraction of the ensemble	
IR ¹	SQ ²	MR ³	LR ⁴	IC ⁵	Total	Count ⁶	%
15	6	13	14	0	48	7	87.5
41	16	33	46	1	137	8	100.0

¹Intra-residue restraints, ²Sequential restraints, ³Medium range restraints, ⁴Long range restraints,

⁵Inter-chain restraints, ⁶ Number of models with violations


9.3.1 Bar graph : Distance violation statistics for the ensemble ⓘ



9.4 Most violated distance restraints in the ensemble ⓘ

9.4.1 Histogram : Distribution of mean distance violations ⓘ

The following histogram shows the distribution of the average value of the violation. The average is calculated for each restraint that is violated in more than one model over all the violated models in the ensemble



9.4.2 Table: Most violated distance restraints [i](#)

The following table provides the mean and the standard deviation of the violations for the 10 worst performing restraints, sorted by number of violated models and the mean violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(2,1063)	1:145:A:ARG:HD3	1:142:A:THR:HA	8	3.24	3.42	0.73
(2,1063)	1:145:A:ARG:HD2	1:142:A:THR:HA	8	3.24	3.42	0.73
(1,166)	1:158:A:LEU:HD13	1:78:A:LEU:HD13	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD11	1:44:A:VAL:HG13	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD13	1:44:A:VAL:HG12	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD11	1:78:A:LEU:HD12	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD12	1:44:A:VAL:HG12	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD12	1:44:A:VAL:HG13	8	2.05	0.84	2.45
(1,166)	1:158:A:LEU:HD12	1:78:A:LEU:HD13	8	2.05	0.84	2.45
(1,125)	1:160:A:ALA:HB1	1:26:A:ILE:HB	8	1.91	1.46	0.92
(1,125)	1:160:A:ALA:HB3	1:26:A:ILE:HB	8	1.91	1.46	0.92
(1,125)	1:160:A:ALA:HB2	1:26:A:ILE:HB	8	1.91	1.46	0.92
(1,104)	1:4:A:ILE:HG23	1:80:A:MET:HB2	8	1.87	0.28	1.92
(1,104)	1:4:A:ILE:HG22	1:80:A:MET:HB2	8	1.87	0.28	1.92
(1,104)	1:4:A:ILE:HG21	1:80:A:MET:HB2	8	1.87	0.28	1.92
(1,104)	1:4:A:ILE:HG23	1:139:A:LYS:HB3	8	1.87	0.28	1.92

Continued on next page...

Continued from previous page...


Key	Atom-1	Atom-2	Models ¹	Mean (Å)	SD ¹ (Å)	Median (Å)
(1,297)	1:4:A:ILE:HD11	1:2:A:ASP:HB2	8	1.78	0.74	1.32
(1,297)	1:4:A:ILE:HD13	1:2:A:ASP:HB2	8	1.78	0.74	1.32
(1,297)	1:4:A:ILE:HD12	1:2:A:ASP:HB2	8	1.78	0.74	1.32
(1,262)	1:151:A:ASP:HB2	1:148:A:ILE:HG22	8	1.7	0.15	1.72
(1,262)	1:151:A:ASP:HB3	1:158:A:LEU:HD22	8	1.7	0.15	1.72
(1,262)	1:151:A:ASP:HB3	1:148:A:ILE:HG23	8	1.7	0.15	1.72
(1,262)	1:151:A:ASP:HB3	1:148:A:ILE:HG21	8	1.7	0.15	1.72
(1,279)	1:60:A:MET:HE1	1:147:A:ARG:HD3	8	1.66	1.0	1.6
(1,279)	1:60:A:MET:HE1	1:145:A:ARG:HD2	8	1.66	1.0	1.6
(1,279)	1:60:A:MET:HE3	1:145:A:ARG:HD2	8	1.66	1.0	1.6
(1,279)	1:60:A:MET:HE3	1:147:A:ARG:HD2	8	1.66	1.0	1.6
(1,279)	1:60:A:MET:HE1	1:147:A:ARG:HD2	8	1.66	1.0	1.6
(1,279)	1:60:A:MET:HE2	1:147:A:ARG:HD2	8	1.66	1.0	1.6
(1,281)	1:154:A:MET:HG3	1:152:A:ALA:HB2	8	1.54	0.37	1.32
(1,281)	1:154:A:MET:HG3	1:152:A:ALA:HB3	8	1.54	0.37	1.32
(1,281)	1:154:A:MET:HG3	1:152:A:ALA:HB1	8	1.54	0.37	1.32
(1,153)	1:41:A:LEU:HD12	1:52:A:PRO:HB3	8	1.44	0.36	1.48
(1,153)	1:41:A:LEU:HD13	1:72:A:VAL:HB	8	1.44	0.36	1.48
(1,153)	1:41:A:LEU:HD12	1:72:A:VAL:HB	8	1.44	0.36	1.48
(1,153)	1:41:A:LEU:HD11	1:72:A:VAL:HB	8	1.44	0.36	1.48
(1,153)	1:41:A:LEU:HD13	1:52:A:PRO:HB3	8	1.44	0.36	1.48
(1,153)	1:41:A:LEU:HD11	1:50:A:GLN:HG3	8	1.44	0.36	1.48
(1,152)	1:41:A:LEU:HD11	1:153:A:MET:HE1	8	1.35	0.46	1.49

¹Number of violated models, ²Standard deviation

9.5 All violated distance restraints [i](#)

9.5.1 Histogram : Distribution of distance violations [i](#)

The following histogram shows the distribution of the absolute value of the violation for all violated restraints in the ensemble.



9.5.2 Table : All distance violations [i](#)

The following table provides the 10 worst performing restraints, sorted by the violation value. The Key (restraint list ID, restraint ID) is the unique identifier for a given restraint. Rows with same key represent combinatorial or ambiguous restraints and are counted as a single restraint.

Key	Atom-1	Atom-2	Model ID	Violation (Å)
(2,1063)	1:145:A:ARG:HD3	1:142:A:THR:HA	6	8.22
(2,1063)	1:145:A:ARG:HD3	1:142:A:THR:HA	8	7.7
(2,1063)	1:145:A:ARG:HD3	1:142:A:THR:HA	7	6.99
(2,1102)	1:145:A:ARG:HD3	1:143:A:LEU:HD13	7	5.16
(2,105)	1:84:A:SER:HA	1:88:A:ASP:HB3	6	5.08
(2,1102)	1:145:A:ARG:HD2	1:143:A:LEU:HD13	6	5.0
(2,497)	1:81:A:MET:HE1	1:85:A:MET:HB3	8	4.54
(2,1102)	1:145:A:ARG:HD3	1:143:A:LEU:HD12	8	4.42
(2,772)	1:7:A:ALA:HB2	1:11:A:GLN:HG2	6	4.06
(1,125)	1:160:A:ALA:HB2	1:26:A:ILE:HB	7	4.06

10 Dihedral-angle violation analysis ⓘ

No dihedral-angle restraints found