



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 18, 2026 – 12:01 AM UTC

PDB ID : 6W87 / pdb_00006w87
Title : K2P2.1 (TREK-1):ML335 complex, 50 mM K+
Authors : Lolicato, M.; Minor, D.L.
Deposited on : 2020-03-20
Resolution : 3.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1466 (3.20-3.20)
Clashscore	190562	1573 (3.20-3.20)
Ramachandran outliers	187476	1548 (3.20-3.20)
Sidechain outliers	187428	1547 (3.20-3.20)
RSRZ outliers	180081	1466 (3.20-3.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	312	% 79% 9% 12%
1	B	312	2% 78% 12% 10%

2 Entry composition

There are 10 unique types of molecules in this entry. The entry contains 4568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Potassium channel subfamily K member 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	276	Total	C	N	O	S	0	1	0
			2167	1446	341	375	5			
1	B	280	Total	C	N	O	S	0	0	0
			2176	1450	344	377	5			

There are 46 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	20	MET	-	expression tag	UNP P97438
A	84	ARG	LYS	engineered mutation	UNP P97438
A	85	GLU	GLN	engineered mutation	UNP P97438
A	86	LYS	THR	engineered mutation	UNP P97438
A	88	LEU	ILE	engineered mutation	UNP P97438
A	89	ARG	ALA	engineered mutation	UNP P97438
A	90	ALA	GLN	engineered mutation	UNP P97438
A	92	PRO	ALA	engineered mutation	UNP P97438
A	95	SER	ASN	engineered mutation	UNP P97438
A	96	ASP	SER	engineered mutation	UNP P97438
A	97	GLN	THR	engineered mutation	UNP P97438
A	119	ALA	ASN	engineered mutation	UNP P97438
A	300	ALA	SER	engineered mutation	UNP P97438
A	306	ALA	GLU	engineered mutation	UNP P97438
A	323	SER	-	expression tag	UNP P97438
A	324	ASN	-	expression tag	UNP P97438
A	325	SER	-	expression tag	UNP P97438
A	326	LEU	-	expression tag	UNP P97438
A	327	GLU	-	expression tag	UNP P97438
A	328	VAL	-	expression tag	UNP P97438
A	329	LEU	-	expression tag	UNP P97438
A	330	PHE	-	expression tag	UNP P97438
A	331	GLN	-	expression tag	UNP P97438
B	20	MET	-	expression tag	UNP P97438
B	84	ARG	LYS	engineered mutation	UNP P97438

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	85	GLU	GLN	engineered mutation	UNP P97438
B	86	LYS	THR	engineered mutation	UNP P97438
B	88	LEU	ILE	engineered mutation	UNP P97438
B	89	ARG	ALA	engineered mutation	UNP P97438
B	90	ALA	GLN	engineered mutation	UNP P97438
B	92	PRO	ALA	engineered mutation	UNP P97438
B	95	SER	ASN	engineered mutation	UNP P97438
B	96	ASP	SER	engineered mutation	UNP P97438
B	97	GLN	THR	engineered mutation	UNP P97438
B	119	ALA	ASN	engineered mutation	UNP P97438
B	300	ALA	SER	engineered mutation	UNP P97438
B	306	ALA	GLU	engineered mutation	UNP P97438
B	323	SER	-	expression tag	UNP P97438
B	324	ASN	-	expression tag	UNP P97438
B	325	SER	-	expression tag	UNP P97438
B	326	LEU	-	expression tag	UNP P97438
B	327	GLU	-	expression tag	UNP P97438
B	328	VAL	-	expression tag	UNP P97438
B	329	LEU	-	expression tag	UNP P97438
B	330	PHE	-	expression tag	UNP P97438
B	331	GLN	-	expression tag	UNP P97438

- Molecule 2 is POTASSIUM ION (CCD ID: K) (formula: K).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	5	Total K 5 5	0	0

- Molecule 3 is N-[(2,4-dichlorophenyl)methyl]-4-[(methylsulfonyl)amino]benzamide (CCD ID: Q6F) (formula: C₁₅H₁₄Cl₂N₂O₃S).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
3	A	1	Total	C	Cl	N	O	S	0	0
			23	15	2	2	3	1		
3	B	1	Total	C	Cl	N	O	S	0	0
			23	15	2	2	3	1		

- Molecule 4 is DODECANE (CCD ID: D12) (formula: $C_{12}H_{26}$).




Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	C	0	0
			12	12		
4	B	1	Total	C	0	0
			12	12		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total C 12 12	0	0
4	B	1	Total C 12 12	0	0

- Molecule 5 is UNDECANE (CCD ID: UND) (formula: $C_{11}H_{24}$).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C 11 11	0	0
5	B	1	Total C 11 11	0	0
5	B	1	Total C 11 11	0	0

- Molecule 6 is DECANE (CCD ID: D10) (formula: $C_{10}H_{22}$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	1	Total C 10 10	0	0

- Molecule 7 is N-OCTANE (CCD ID: OCT) (formula: C_8H_{18}).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total C 8 8	0	0
7	A	1	Total C 8 8	0	0

- Molecule 8 is CADMIUM ION (CCD ID: CD) (formula: Cd).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total Cd 1 1	0	0
8	B	3	Total Cd 3 3	0	0

- Molecule 9 is HEXADECANE (CCD ID: R16) (formula: C₁₆H₃₄).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
9	B	1	Total C 16 16	0	0
9	B	1	Total C 16 16	0	0
9	B	1	Total C 16 16	0	0

- Molecule 10 is PENTANE (CCD ID: LNK) (formula: C₅H₁₂).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	B	1	Total C 5 5	0	0
10	B	1	Total C 5 5	0	0
10	B	1	Total C 5 5	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Potassium channel subfamily K member 2

Chain A: 



- Molecule 1: Potassium channel subfamily K member 2

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	66.94Å 119.48Å 128.39Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	14.95 – 3.20 14.95 – 3.20	Depositor EDS
% Data completeness (in resolution range)	100.0 (14.95-3.20) 98.8 (14.95-3.20)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.15 (at 3.19Å)	Xtriage
Refinement program	BUSTER 2.10.3 (6-FEB-2020)	Depositor
R, R_{free}	0.246 , 0.296 0.258 , 0.310	Depositor DCC
R_{free} test set	870 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	125.5	Xtriage
Anisotropy	0.092	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.24 , 94.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4568	wwPDB-VP
Average B, all atoms (Å ²)	143.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.29% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: K, D12, UND, D10, R16, CD, LNK, Q6F, OCT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.69	0/2224	0.92	1/3024 (0.0%)
1	B	0.67	0/2228	0.91	1/3029 (0.0%)
All	All	0.68	0/4452	0.92	2/6053 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	A	143	ILE	N-CA-C	-5.76	105.00	110.42
1	B	143	ILE	N-CA-C	-5.28	105.57	110.53

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	268	ASP	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2167	0	2217	14	0
1	B	2176	0	2237	20	0
2	A	5	0	0	0	0
3	A	23	0	0	1	0
3	B	23	0	0	1	0
4	A	12	0	26	0	0
4	B	36	0	78	1	0
5	A	11	0	24	0	0
5	B	22	0	48	1	0
6	A	10	0	22	0	0
7	A	16	0	36	0	0
8	A	1	0	0	0	0
8	B	3	0	0	0	0
9	B	48	0	102	1	0
10	B	15	0	36	0	0
All	All	4568	0	4826	31	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (31) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:294:ASP:HA	1:B:297:ARG:HD2	1.64	0.78
1:B:197:ILE:HA	1:B:201:VAL:HG11	1.79	0.65
1:B:210:SER:O	1:B:213:ILE:HG22	1.98	0.63
1:A:197:ILE:HD11	1:A:206:ILE:HG21	1.82	0.62
1:A:88:LEU:HD11	1:A:99:LEU:HD22	1.84	0.60
1:B:238:ALA:HB3	9:B:409:R16:H422	1.82	0.60
1:B:85:GLU:HG2	1:B:88:LEU:HD12	1.86	0.56
1:B:107:VAL:O	1:B:110:ILE:HG22	2.12	0.49
1:A:243:TYR:HA	1:B:160:ILE:HD11	1.95	0.49
1:B:278:ILE:CD1	3:B:404:Q6F:C21	2.91	0.48
1:A:77:ARG:NH1	1:B:123:GLN:OE1	2.47	0.47
1:A:70:GLN:HB3	1:A:71:PRO:HD3	1.96	0.47
1:B:110:ILE:HA	1:B:114:ILE:HG12	1.95	0.47
1:A:278:ILE:CD1	3:A:403:Q6F:C21	2.93	0.46
1:A:135:PHE:CZ	1:A:150:PRO:HD3	2.50	0.46
1:A:111:ASN:O	1:A:151:ARG:NE	2.36	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:167:ILE:HB	1:B:168:PRO:HD3	1.99	0.45
1:A:167:ILE:HB	1:A:168:PRO:HD3	1.98	0.44
1:B:273:VAL:HG12	4:B:407:D12:H22	1.99	0.44
1:A:197:ILE:HG13	1:A:206:ILE:HD13	1.99	0.44
1:B:60:ILE:O	1:B:64:VAL:HG23	2.18	0.44
1:B:124:VAL:HG12	1:B:125:SER:N	2.33	0.44
1:B:152:THR:HG22	1:B:154:GLY:H	1.82	0.44
1:A:60:ILE:O	1:A:64:VAL:HG23	2.18	0.44
1:A:110:ILE:HD13	1:B:110:ILE:HG21	2.00	0.44
1:B:268:ASP:OD1	5:B:411:UND:H112	2.18	0.43
1:A:152:THR:HG22	1:A:154:GLY:H	1.83	0.43
1:B:135:PHE:CZ	1:B:150:PRO:HD3	2.54	0.42
1:B:70:GLN:HB3	1:B:71:PRO:HD3	2.00	0.42
1:B:271:LYS:N	1:B:272:PRO:HD2	2.34	0.42
1:A:271:LYS:N	1:A:272:PRO:HD2	2.35	0.41

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/312 (88%)	256 (94%)	15 (6%)	2 (1%)	18	52
1	B	276/312 (88%)	255 (92%)	18 (6%)	3 (1%)	11	43
All	All	549/624 (88%)	511 (93%)	33 (6%)	5 (1%)	14	47

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	201	VAL
1	B	264	ILE
1	A	199	TRP

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	B	95	SER
1	B	266	TYR

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	228/260 (88%)	224 (98%)	4 (2%)	51	74
1	B	229/260 (88%)	226 (99%)	3 (1%)	61	78
All	All	457/520 (88%)	450 (98%)	7 (2%)	57	76

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LEU
1	A	98	GLU
1	A	129	LEU
1	A	265	GLU
1	B	93	CYS
1	B	202	SER
1	B	269	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	70	GLN
1	B	91	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 27 ligands modelled in this entry, 9 are monoatomic - leaving 18 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	D12	B	405	-	11,11,11	0.51	0	10,10,10	0.50	0
7	OCT	A	411	-	7,7,7	0.43	0	6,6,6	0.44	0
9	R16	B	409	-	15,15,15	0.50	0	14,14,14	0.53	0
9	R16	B	410	-	15,15,15	0.60	0	14,14,14	0.30	0
6	D10	A	409	-	9,9,9	0.47	0	8,8,8	0.61	0
10	LNK	B	414	-	4,4,4	0.46	0	3,3,3	0.34	0
3	Q6F	A	403	-	24,24,24	1.95	6 (25%)	34,34,34	2.83	9 (26%)
4	D12	B	406	-	11,11,11	0.58	0	10,10,10	0.41	0
7	OCT	A	410	-	7,7,7	0.44	0	6,6,6	0.53	0
9	R16	B	408	-	15,15,15	0.42	0	14,14,14	0.76	0
5	UND	A	408	-	10,10,10	0.47	0	9,9,9	0.52	0
5	UND	B	412	-	10,10,10	0.50	0	9,9,9	0.49	0
10	LNK	B	413	-	4,4,4	0.48	0	3,3,3	0.29	0
4	D12	A	407	-	11,11,11	0.54	0	10,10,10	0.42	0
10	LNK	B	415	-	4,4,4	0.43	0	3,3,3	0.39	0
5	UND	B	411	-	10,10,10	0.47	0	9,9,9	0.54	0
3	Q6F	B	404	-	24,24,24	1.77	5 (20%)	34,34,34	2.89	11 (32%)
4	D12	B	407	-	11,11,11	0.53	0	10,10,10	0.45	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns.

'-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	D12	B	405	-	-	3/9/9/9	-
7	OCT	A	411	-	-	2/5/5/5	-
9	R16	B	409	-	-	4/13/13/13	-
9	R16	B	410	-	-	8/13/13/13	-
6	D10	A	409	-	-	1/7/7/7	-
10	LNK	B	414	-	-	0/2/2/2	-
3	Q6F	A	403	-	-	2/14/14/14	0/2/2/2
4	D12	B	406	-	-	0/9/9/9	-
7	OCT	A	410	-	-	0/5/5/5	-
9	R16	B	408	-	-	4/13/13/13	-
5	UND	A	408	-	-	0/8/8/8	-
5	UND	B	412	-	-	1/8/8/8	-
10	LNK	B	413	-	-	1/2/2/2	-
4	D12	A	407	-	-	4/9/9/9	-
10	LNK	B	415	-	-	0/2/2/2	-
5	UND	B	411	-	-	4/8/8/8	-
3	Q6F	B	404	-	-	1/14/14/14	0/2/2/2
4	D12	B	407	-	-	3/9/9/9	-

All (11) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	403	Q6F	C13-N07	6.17	1.47	1.33
3	B	404	Q6F	C13-N07	5.13	1.45	1.33
3	A	403	Q6F	C10-C13	3.75	1.58	1.50
3	B	404	Q6F	C10-C13	3.13	1.57	1.50
3	B	404	Q6F	C11-N08	2.90	1.47	1.43
3	A	403	Q6F	C17-C15	2.78	1.43	1.38
3	A	403	Q6F	C20-S03	2.73	1.81	1.75
3	A	403	Q6F	C11-N08	2.67	1.47	1.43
3	B	404	Q6F	C18-CL1	2.65	1.79	1.73
3	B	404	Q6F	C20-S03	2.55	1.81	1.75
3	A	403	Q6F	C18-CL1	2.10	1.78	1.73

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	404	Q6F	O06-S03-O05	-12.95	101.51	118.87
3	A	403	Q6F	O06-S03-O05	-12.64	101.93	118.87

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	404	Q6F	C20-S03-N08	4.97	112.16	106.56
3	A	403	Q6F	C12-C09-N07	-4.91	103.26	113.15
3	A	403	Q6F	O06-S03-C20	4.24	114.97	108.26
3	B	404	Q6F	C12-C09-N07	-3.61	105.88	113.15
3	B	404	Q6F	C17-C11-N08	-3.59	112.45	120.02
3	B	404	Q6F	C16-C11-N08	3.44	127.25	120.02
3	B	404	Q6F	O06-S03-C20	3.38	113.61	108.26
3	A	403	Q6F	O05-S03-C20	3.24	113.39	108.26
3	B	404	Q6F	C21-C18-C12	-2.95	119.83	122.41
3	A	403	Q6F	O05-S03-N08	2.87	112.83	107.10
3	A	403	Q6F	C12-C18-CL1	2.86	124.31	119.75
3	A	403	Q6F	C09-C12-C18	2.65	124.57	121.08
3	B	404	Q6F	O05-S03-C20	2.62	112.41	108.26
3	B	404	Q6F	C21-C23-CL2	2.48	122.28	119.17
3	A	403	Q6F	C21-C18-C12	-2.38	120.33	122.41
3	B	404	Q6F	C22-C23-C21	-2.31	118.51	121.53
3	A	403	Q6F	C09-N07-C13	2.13	126.87	121.89
3	B	404	Q6F	C18-C21-C23	2.10	121.09	118.72

There are no chirality outliers.

All (38) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	403	Q6F	C11-N08-S03-C20
3	A	403	Q6F	C11-N08-S03-O05
9	B	408	R16	C37-C38-C39-C40
5	B	411	UND	C4-C5-C6-C7
9	B	410	R16	C34-C35-C36-C37
4	B	407	D12	C6-C7-C8-C9
4	A	407	D12	C7-C8-C9-C10
9	B	408	R16	C36-C37-C38-C39
9	B	410	R16	C30-C31-C32-C33
9	B	408	R16	C35-C36-C37-C38
9	B	410	R16	C33-C34-C35-C36
4	A	407	D12	C2-C3-C4-C5
9	B	410	R16	C31-C32-C33-C34
5	B	412	UND	C7-C8-C9-C10
9	B	410	R16	C37-C38-C39-C40
9	B	410	R16	C38-C39-C40-C41
4	B	405	D12	C4-C5-C6-C7
4	A	407	D12	C9-C10-C11-C12
4	B	407	D12	C9-C10-C11-C12

Continued on next page...

Continued from previous page...


Mol	Chain	Res	Type	Atoms
9	B	409	R16	C36-C37-C38-C39
4	B	405	D12	C6-C7-C8-C9
7	A	411	OCT	C1-C2-C3-C4
7	A	411	OCT	C5-C6-C7-C8
5	B	411	UND	C2-C3-C4-C5
6	A	409	D10	C5-C6-C7-C8
3	B	404	Q6F	C11-N08-S03-O05
9	B	409	R16	C35-C36-C37-C38
9	B	409	R16	C32-C33-C34-C35
9	B	410	R16	C29-C30-C31-C32
5	B	411	UND	C7-C8-C9-C10
4	A	407	D12	C6-C7-C8-C9
10	B	413	LNK	C1-C2-C3-C4
4	B	405	D12	C3-C4-C5-C6
9	B	410	R16	C39-C40-C41-C42
9	B	409	R16	C29-C30-C31-C32
9	B	408	R16	C28-C29-C30-C31
5	B	411	UND	C1-C2-C3-C4
4	B	407	D12	C4-C5-C6-C7

There are no ring outliers.

5 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
9	B	409	R16	1	0
3	A	403	Q6F	1	0
5	B	411	UND	1	0
3	B	404	Q6F	1	0
4	B	407	D12	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	276/312 (88%)	-0.08	4 (1%)	73 54	78, 131, 224, 275	1 (0%)
1	B	280/312 (89%)	-0.12	5 (1%)	67 48	85, 132, 233, 290	0
All	All	556/624 (89%)	-0.10	9 (1%)	70 51	78, 132, 230, 290	1 (0%)

All (9) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	71	PRO	3.2
1	A	197	ILE	2.9
1	B	132	SER	2.7
1	B	129	LEU	2.6
1	A	255	GLY	2.6
1	B	268	ASP	2.5
1	B	93	CYS	2.4
1	A	72	GLN	2.2
1	B	254	PHE	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
10	LNK	B	415	5/5	0.09	0.31	140,140,140,141	0
4	D12	B	405	12/12	0.60	0.10	138,141,141,142	0
10	LNK	B	414	5/5	0.62	0.25	124,124,124,124	0
4	D12	A	407	12/12	0.67	0.20	118,122,126,126	0
5	UND	A	408	11/11	0.68	0.18	128,136,145,146	0
5	UND	B	412	11/11	0.68	0.19	135,136,139,139	0
5	UND	B	411	11/11	0.70	0.30	125,129,132,132	0
4	D12	B	406	12/12	0.72	0.15	102,113,126,127	0
7	OCT	A	411	8/8	0.72	0.24	130,133,138,138	0
2	K	A	405	1/1	0.80	0.10	110,110,110,110	0
8	CD	A	412	1/1	0.81	0.12	171,171,171,171	1
9	R16	B	409	16/16	0.82	0.11	123,134,145,146	0
9	R16	B	410	16/16	0.82	0.10	139,151,162,163	0
6	D10	A	409	10/10	0.83	0.19	114,120,124,125	0
7	OCT	A	410	8/8	0.84	0.33	127,132,142,144	0
10	LNK	B	413	5/5	0.84	0.31	108,108,108,108	0
8	CD	B	403	1/1	0.85	0.14	196,196,196,196	1
4	D12	B	407	12/12	0.85	0.17	117,133,142,143	0
9	R16	B	408	16/16	0.87	0.14	105,109,119,120	0
2	K	A	406	1/1	0.87	0.06	148,148,148,148	0
3	Q6F	A	403	23/23	0.90	0.09	107,109,132,134	0
2	K	A	404	1/1	0.91	0.06	119,119,119,119	0
3	Q6F	B	404	23/23	0.93	0.08	96,103,125,130	0
2	K	A	401	1/1	0.95	0.06	115,115,115,115	0
8	CD	B	401	1/1	0.97	0.04	234,234,234,234	0
2	K	A	402	1/1	0.98	0.03	101,101,101,101	0
8	CD	B	402	1/1	0.99	0.05	160,160,160,160	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around Q6F A 403:

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around Q6F B 404:**

$2mF_o - DF_c$ (at 0.7 rmsd) in gray
 $mF_o - DF_c$ (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.