



# wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 7, 2026 – 02:22 AM UTC

PDB ID : 7W8A / pdb\_00007w8a  
Title : Babesia orientalis lactate dehydrogenase, BoLDH apo  
Authors : Yu, L.  
Deposited on : 2021-12-07  
Resolution : 2.67 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0  
Xtriage (Phenix) : 2.0  
EDS : 3.0  
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)  
CCP4 : 9.0.010 (Gargrove)  
Density-Fitness : 1.0.12  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.49


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.67 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	180053	5070 (2.70-2.66)
Clashscore	190562	5409 (2.70-2.66)
Ramachandran outliers	187476	5324 (2.70-2.66)
Sidechain outliers	187428	5324 (2.70-2.66)
RSRZ outliers	180081	5070 (2.70-2.66)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	330	 2% 90% 6%
1	B	330	 2% 87% 11%
1	C	330	 2% 89% 9%
1	D	330	 2% 87% 12%
1	E	330	 2% 84% 5% 11%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
1	F	330	 2% 88% 12%
1	G	330	 % 87% 12%
1	H	330	 2% 88% 9%

## 2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 35845 atoms, of which 18104 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called lactate dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	311	4719	1483	2382	393	445	16	3	0	0
1	B	294	4433	1393	2240	368	416	16	0	0	0
1	C	301	4531	1425	2288	375	426	17	3	0	0
1	D	292	4410	1386	2228	366	414	16	0	0	0
1	E	293	4416	1388	2230	367	415	16	4	0	0
1	F	292	4402	1383	2223	366	414	16	0	0	0
1	G	292	4411	1386	2229	366	414	16	0	1	0
1	H	300	4523	1423	2284	374	425	17	0	0	0



### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



- Molecule 1: lactate dehydrogenase





- Molecule 1: lactate dehydrogenase





- Molecule 1: lactate dehydrogenase



- Molecule 1: lactate dehydrogenase





- Molecule 1: lactate dehydrogenase









- Molecule 1: lactate dehydrogenase



- Molecule 1: lactate dehydrogenase



- Molecule 1: lactate dehydrogenase



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.85Å 92.30Å 109.89Å 77.47° 70.69° 64.36°	Depositor
Resolution (Å)	45.00 – 2.67 45.00 – 2.67	Depositor EDS
% Data completeness (in resolution range)	93.5 (45.00-2.67) 93.5 (45.00-2.67)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.81 (at 2.69Å)	Xtrriage
Refinement program	PHENIX 1.12_2829	Depositor
R, $R_{free}$	0.236 , 0.268 0.237 , 0.267	Depositor DCC
$R_{free}$ test set	3683 reflections (4.76%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	27.3	Xtrriage
Anisotropy	0.394	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 16.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.011 for h,h-k,h-l	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	35845	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	26.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.08% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.10	0/2374	0.27	0/3212
1	B	0.10	0/2226	0.27	0/3011
1	C	0.10	0/2276	0.24	0/3076
1	D	0.11	0/2212	0.24	0/2988
1	E	0.11	0/2219	0.26	0/3001
1	F	0.10	0/2210	0.25	0/2986
1	G	0.11	0/2223	0.26	0/3004
1	H	0.10	0/2273	0.26	0/3073
All	All	0.10	0/18013	0.26	0/24351

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2337	2382	2381	8	0
1	B	2193	2240	2238	3	1
1	C	2243	2288	2286	3	0
1	D	2182	2228	2226	4	0
1	E	2186	2230	2229	9	1
1	F	2179	2223	2221	2	0
1	G	2182	2229	2212	2	0
1	H	2239	2284	2281	5	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	17741	18104	18074	30	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

The worst 5 of 30 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:46:ASP:OD1	1:E:47:ILE:N	2.39	0.56
1:E:306:LYS:NZ	1:F:216:GLU:OE1	2.41	0.54
1:A:135:CYS:HB3	1:A:144:VAL:HG22	1.97	0.46
1:D:30:MET:HE2	1:D:256:ALA:HB3	1.98	0.45
1:A:101:ASP:O	1:A:102:GLU:HB2	2.16	0.45

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:127:TYR:OH	1:E:83:ASP:OD2[1_456]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	309/330 (94%)	297 (96%)	11 (4%)	1 (0%)	36 57
1	B	290/330 (88%)	280 (97%)	10 (3%)	0	100 100
1	C	295/330 (89%)	285 (97%)	9 (3%)	1 (0%)	36 57
1	D	284/330 (86%)	269 (95%)	15 (5%)	0	100 100
1	E	289/330 (88%)	277 (96%)	12 (4%)	0	100 100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	F	286/330 (87%)	273 (96%)	13 (4%)	0	100	100
1	G	287/330 (87%)	271 (94%)	16 (6%)	0	100	100
1	H	294/330 (89%)	284 (97%)	10 (3%)	0	100	100
All	All	2334/2640 (88%)	2236 (96%)	96 (4%)	2 (0%)	48	71

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	102	GLU
1	C	162	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	250/265 (94%)	248 (99%)	2 (1%)	73	87
1	B	234/265 (88%)	233 (100%)	1 (0%)	84	93
1	C	240/265 (91%)	240 (100%)	0	100	100
1	D	233/265 (88%)	233 (100%)	0	100	100
1	E	233/265 (88%)	232 (100%)	1 (0%)	84	93
1	F	232/265 (88%)	231 (100%)	1 (0%)	84	93
1	G	234/265 (88%)	233 (100%)	1 (0%)	84	93
1	H	240/265 (91%)	240 (100%)	0	100	100
All	All	1896/2120 (89%)	1890 (100%)	6 (0%)	86	94

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	66	TYR
1	F	17	LYS
1	G	290	THR
1	A	106	ARG

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	66	TYR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
1	D	222	GLN
1	F	130	ASN
1	H	222	GLN
1	G	245	ASN
1	H	113	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	311/330 (94%)	0.16	8 (2%) 57 53	21, 26, 33, 46	0
1	B	294/330 (89%)	0.12	5 (1%) 69 66	20, 25, 31, 36	0
1	C	301/330 (91%)	0.09	5 (1%) 69 66	18, 24, 30, 33	0
1	D	292/330 (88%)	0.14	6 (2%) 63 60	21, 27, 32, 38	0
1	E	293/330 (88%)	0.23	7 (2%) 59 56	21, 28, 32, 38	0
1	F	292/330 (88%)	0.19	8 (2%) 56 52	21, 27, 32, 42	0
1	G	292/330 (88%)	0.12	4 (1%) 73 71	21, 26, 32, 36	0
1	H	300/330 (90%)	0.20	6 (2%) 65 61	21, 26, 30, 34	0
All	All	2375/2640 (89%)	0.16	49 (2%) 63 60	18, 26, 32, 46	0

The worst 5 of 49 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	G	12	ALA	4.8
1	A	162	GLY	4.5
1	A	101	ASP	4.3
1	A	245	ASN	3.8
1	F	48	ALA	3.6

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.