



wwPDB X-ray Structure Validation Summary Report ⓘ

Mar 8, 2026 – 05:21 PM UTC

PDB ID : 8W8O / pdb_00008w8o
Title : Thermus thermophilus initiation complex in the half-translocated state
Authors : Li, L.; Zhang, Y.
Deposited on : 2023-09-04
Resolution : 2.51 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4-5-2 with Phenix2.0
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 2.0
EDS : 3.0
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4 : 9.0.010 (Gargrove)
Density-Fitness : 1.0.12
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	7383 (2.54-2.50)
Clashscore	190562	8079 (2.54-2.50)
Ramachandran outliers	187476	7944 (2.54-2.50)
Sidechain outliers	187428	7946 (2.54-2.50)
RSRZ outliers	180081	7387 (2.54-2.50)
RNA backbone	3983	1161 (2.80-2.24)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	315	58% 13% • 28%
1	B	315	2% 53% 17% • 29%
2	C	1119	2% 80% 18% ••
3	D	1524	5% 77% 19% ••

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	E	99	2% 90% 6%
5	F	443	4% 61% 16% 22%
6	G	21	10% 48% 29% 14%
7	H	27	4% 33% 56% 11%
8	I	3	100%

2 Entry composition

There are 11 unique types of molecules in this entry. The entry contains 29261 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-directed RNA polymerase subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	228	Total	C	N	O	S	0	0	0
			1792	1144	312	334	2			
1	B	224	Total	C	N	O	S	0	0	0
			1767	1130	307	328	2			

- Molecule 2 is a protein called DNA-directed RNA polymerase subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	C	1107	Total	C	N	O	S	0	0	0
			8728	5523	1560	1621	24			

- Molecule 3 is a protein called DNA-directed RNA polymerase subunit beta'.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	D	1481	Total	C	N	O	S	0	3	0
			11682	7412	2056	2178	36			

- Molecule 4 is a protein called DNA-directed RNA polymerase subunit omega.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	E	96	Total	C	N	O	S	0	0	0
			781	497	137	143	4			

- Molecule 5 is a protein called RNA polymerase sigma factor SigA.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
5	F	346	Total	C	N	O	S	0	0	0
			2804	1767	509	524	4			

There are 20 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
F	-19	MET	-	expression tag	UNP Q5SKW1
F	-18	GLY	-	expression tag	UNP Q5SKW1
F	-17	SER	-	expression tag	UNP Q5SKW1
F	-16	SER	-	expression tag	UNP Q5SKW1
F	-15	HIS	-	expression tag	UNP Q5SKW1
F	-14	HIS	-	expression tag	UNP Q5SKW1
F	-13	HIS	-	expression tag	UNP Q5SKW1
F	-12	HIS	-	expression tag	UNP Q5SKW1
F	-11	HIS	-	expression tag	UNP Q5SKW1
F	-10	HIS	-	expression tag	UNP Q5SKW1
F	-9	SER	-	expression tag	UNP Q5SKW1
F	-8	SER	-	expression tag	UNP Q5SKW1
F	-7	GLY	-	expression tag	UNP Q5SKW1
F	-6	LEU	-	expression tag	UNP Q5SKW1
F	-5	VAL	-	expression tag	UNP Q5SKW1
F	-4	PRO	-	expression tag	UNP Q5SKW1
F	-3	ARG	-	expression tag	UNP Q5SKW1
F	-2	GLY	-	expression tag	UNP Q5SKW1
F	-1	SER	-	expression tag	UNP Q5SKW1
F	0	HIS	-	expression tag	UNP Q5SKW1

- Molecule 6 is a DNA chain called DNA (21-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
6	G	18	Total	C	N	O	P	0	0	0
			369	175	71	106	17			

- Molecule 7 is a DNA chain called DNA (27-MER).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
7	H	24	Total	C	N	O	P	0	0	0
			495	236	94	142	23			

- Molecule 8 is a RNA chain called RNA (5'-(GTP)GA-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
8	I	3	Total	C	N	O	P	0	0	0
			77	30	15	27	5			

- Molecule 9 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
9	B	1	Total 1	Mg 1	0	0
9	D	3	Total 3	Mg 3	0	0
9	F	1	Total 1	Mg 1	0	0
9	I	2	Total 2	Mg 2	0	0

- Molecule 10 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
10	D	2	Total 2	Zn 2	0	0


- Molecule 11 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
11	A	44	Total 44	O 44	0	0
11	B	29	Total 29	O 29	0	0
11	C	270	Total 270	O 270	0	0
11	D	304	Total 304	O 304	0	0
11	E	32	Total 32	O 32	0	0
11	F	37	Total 37	O 37	0	0
11	G	14	Total 14	O 14	0	0
11	H	5	Total 5	O 5	0	0
11	I	22	Total 22	O 22	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 1: DNA-directed RNA polymerase subunit alpha




• Molecule 2: DNA-directed RNA polymerase subunit beta





• Molecule 3: DNA-directed RNA polymerase subunit beta'





- Molecule 8: RNA (5'-(GTP)GA-3')

Chain I:  100%

There are no outlier residues recorded for this chain.

4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	183.40Å 103.80Å 295.04Å 90.00° 99.19° 90.00°	Depositor
Resolution (Å)	39.88 – 2.51 39.88 – 2.51	Depositor EDS
% Data completeness (in resolution range)	99.0 (39.88-2.51) 99.0 (39.88-2.51)	Depositor EDS
R_{merge}	0.18	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.79 (at 2.51Å)	Xtriage
Refinement program	PHENIX (1.12_2829: ???)	Depositor
R, R_{free}	0.187 , 0.226 0.189 , 0.228	Depositor DCC
R_{free} test set	9212 reflections (4.93%)	wwPDB-VP
Wilson B-factor (Å ²)	51.4	Xtriage
Anisotropy	0.345	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.29 , 35.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.012 for $1/2^*h-3/2^*k,-1/2^*h-1/2^*k,-1/2^*h+1/2^*k-l$ 0.012 for $1/2^*h+3/2^*k,1/2^*h-1/2^*k,-1/2^*h-1/2^*k-l$	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	29261	wwPDB-VP
Average B, all atoms (Å ²)	65.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.93% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GTP, MG, ZN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.39	0/1824	0.62	0/2480
1	B	0.41	0/1799	0.58	0/2446
2	C	0.42	0/8892	0.62	0/12022
3	D	0.41	0/11895	0.60	1/16084 (0.0%)
4	E	0.42	0/795	0.60	0/1071
5	F	0.36	0/2849	0.56	0/3833
6	G	0.57	0/414	1.07	7/638 (1.1%)
7	H	0.46	0/556	0.61	0/858
8	I	0.52	0/50	0.86	0/76
All	All	0.41	0/29074	0.62	8/39508 (0.0%)

There are no bond length outliers.

The worst 5 of 8 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
6	G	19	DG	P-O3'-C3'	-7.89	108.36	120.20
6	G	6	DA	P-O3'-C3'	-7.67	108.69	120.20
6	G	18	DA	P-O3'-C3'	-7.56	108.86	120.20
6	G	15	DT	P-O3'-C3'	-7.32	109.23	120.20
6	G	17	DC	P-O3'-C3'	-7.05	109.62	120.20

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1792	0	1844	31	0
1	B	1767	0	1821	39	0
2	C	8728	0	8834	140	0
3	D	11682	0	11899	192	0
4	E	781	0	797	5	0
5	F	2804	0	2873	51	0
6	G	369	0	203	22	0
7	H	495	0	272	14	0
8	I	77	0	33	0	0
9	B	1	0	0	0	0
9	D	3	0	0	0	0
9	F	1	0	0	0	0
9	I	2	0	0	0	0
10	D	2	0	0	0	0
11	A	44	0	0	4	0
11	B	29	0	0	0	0
11	C	270	0	0	8	0
11	D	304	0	0	17	0
11	E	32	0	0	1	0
11	F	37	0	0	3	0
11	G	14	0	0	1	0
11	H	5	0	0	0	0
11	I	22	0	0	0	0
All	All	29261	0	28576	443	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

The worst 5 of 443 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:D:1380:GLU:HB3	3:D:1418:LYS:HD3	1.56	0.87
6:G:9:DC:H2'	6:G:10:DG:C8	2.13	0.84
2:C:853:LEU:HB2	2:C:858:MET:HE1	1.60	0.82
1:A:39:PRO:HG3	1:B:39:PRO:HG3	1.65	0.79
3:D:238:PRO:HD3	3:D:318:ARG:HG3	1.63	0.78

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	226/315 (72%)	225 (100%)	1 (0%)	0	100	100
1	B	222/315 (70%)	214 (96%)	8 (4%)	0	100	100
2	C	1099/1119 (98%)	1066 (97%)	33 (3%)	0	100	100
3	D	1476/1524 (97%)	1440 (98%)	36 (2%)	0	100	100
4	E	94/99 (95%)	92 (98%)	2 (2%)	0	100	100
5	F	344/443 (78%)	336 (98%)	8 (2%)	0	100	100
All	All	3461/3815 (91%)	3373 (98%)	88 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	199/273 (73%)	189 (95%)	10 (5%)	22	42
1	B	197/273 (72%)	187 (95%)	10 (5%)	21	41
2	C	929/941 (99%)	881 (95%)	48 (5%)	21	40
3	D	1242/1279 (97%)	1173 (94%)	69 (6%)	19	37
4	E	85/88 (97%)	83 (98%)	2 (2%)	43	69
5	F	300/388 (77%)	279 (93%)	21 (7%)	14	27
All	All	2952/3242 (91%)	2792 (95%)	160 (5%)	20	39

5 of 160 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
3	D	1128	VAL
5	F	96	LEU
3	D	1207	TYR
3	D	1424	VAL
5	F	295	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 13 such sidechains are listed below:

Mol	Chain	Res	Type
2	C	991	GLN
2	C	1006	HIS
3	D	1333	HIS
3	D	709	HIS
3	D	994	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
8	I	1/3 (33%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 9 ligands modelled in this entry, 9 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	228/315 (72%)	-0.07	1 (0%) 88 87	43, 58, 81, 99	0
1	B	224/315 (71%)	0.34	7 (3%) 51 48	46, 68, 95, 116	0
2	C	1107/1119 (98%)	-0.00	27 (2%) 59 56	31, 52, 109, 134	0
3	D	1481/1524 (97%)	0.23	74 (4%) 34 31	31, 59, 116, 142	3 (0%)
4	E	96/99 (96%)	-0.06	2 (2%) 63 60	36, 52, 98, 110	0
5	F	346/443 (78%)	0.45	19 (5%) 30 27	42, 71, 111, 128	0
6	G	18/21 (85%)	0.26	0 100 100	50, 77, 165, 166	0
7	H	24/27 (88%)	0.14	1 (4%) 40 37	70, 85, 134, 166	0
8	I	2/3 (66%)	-0.95	0 100 100	43, 43, 43, 43	0
All	All	3526/3866 (91%)	0.16	131 (3%) 45 41	31, 59, 112, 166	3 (0%)

The worst 5 of 131 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
3	D	142	LEU	6.0
3	D	1207	TYR	4.7
3	D	203	ALA	4.7
5	F	139	ALA	4.7
3	D	420	VAL	4.6

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
9	MG	D	1605	1/1	0.82	0.08	56,56,56,56	0
9	MG	B	2001	1/1	0.92	0.09	73,73,73,73	0
9	MG	F	501	1/1	0.95	0.09	56,56,56,56	0
9	MG	D	1604	1/1	0.99	0.04	64,64,64,64	0
9	MG	I	101	1/1	0.99	0.04	43,43,43,43	0
9	MG	I	102	1/1	0.99	0.03	44,44,44,44	0
9	MG	D	1603	1/1	1.00	0.03	33,33,33,33	0
10	ZN	D	1601	1/1	1.00	0.03	37,37,37,37	0
10	ZN	D	1602	1/1	1.00	0.02	76,76,76,76	0

6.5 Other polymers [i](#)

There are no such residues in this entry.