



wwPDB EM Validation Summary Report ⓘ

Jul 14, 2026 – 06:13 AM JST

PDB ID : 9W8W / pdb_00009w8w
EMDB ID : EMD-65754
Title : Structure of backtracked yeast Pol II in complex with a cisplatin-induced inter-strand crosslink lesion at the +2 position.
Authors : Xu, J.; Zhu, L.
Deposited on : 2025-08-08
Resolution : 2.91 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : 0.0.1.dev133
Mogul : 1.8.5 (274361), CSD as541be (2020)
MolProbity : 4-5-2 with Phenix2.0
Buster-report : wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics : 20250101.v01 (using entries in the PDB archive January 1st 2025)
EM percentile statistics : 202505.v01 (Using data in the EMDB archive up until May 2025)
MapQ : 1.9.13
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.50



1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.91 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	EM structures (#Entries)	Similar EM resolution (#Entries, resolution range(Å))
Clashscore	229148	23984	-
Ramachandran outliers	224038	23583	-
Sidechain outliers	223484	23102	-
RNA backbone	8273	3508	-
Q-score	-	25397	12972 (2.41 - 3.41)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the EM map (all-atom inclusion $< 40\%$). The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	N	10	
2	P	21	
3	T	20	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	C	318	
5	E	215	
6	F	155	
7	G	171	
8	I	122	
9	J	70	
10	K	120	
11	L	70	
12	A	1733	
13	B	1224	
14	D	221	
15	H	146	

2 Entry composition

There are 18 unique types of molecules in this entry. The entry contains 32389 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a DNA chain called DNA (NTS, non-template strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
1	N	6	Total	C	N	O	P	0	0
			128	59	28	35	6		

- Molecule 2 is a RNA chain called RNA (5'-R(*UP*AP*CP*AP*AP*UP*GP*UP*UP*GP*GP*AP*AP*AP*GP*GP*AP*GP*AP*GP*A)-3').

Mol	Chain	Residues	Atoms					AltConf	Trace
2	P	12	Total	C	N	O	P	0	0
			270	120	60	78	12		

- Molecule 3 is a DNA chain called DNA (TS, template strand).

Mol	Chain	Residues	Atoms					AltConf	Trace
3	T	16	Total	C	N	O	P	0	0
			313	151	44	102	16		

- Molecule 4 is a protein called DNA-directed RNA polymerase II subunit RPB3.

Mol	Chain	Residues	Atoms					AltConf	Trace
4	C	265	Total	C	N	O	S	0	0
			2086	1312	347	414	13		

- Molecule 5 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC1.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	E	214	Total	C	N	O	S	0	0
			1752	1111	309	321	11		

- Molecule 6 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC2.

Mol	Chain	Residues	Atoms					AltConf	Trace
6	F	87	Total	C	N	O	S	0	0
			705	451	119	132	3		

- Molecule 7 is a protein called DNA-directed RNA polymerase II subunit RPB7.

Mol	Chain	Residues	Atoms					AltConf	Trace
7	G	171	Total	C	N	O	S	0	0
			1339	861	222	248	8		

- Molecule 8 is a protein called DNA-directed RNA polymerase II subunit RPB9.

Mol	Chain	Residues	Atoms					AltConf	Trace
8	I	116	Total	C	N	O	S	0	0
			944	581	172	181	10		

- Molecule 9 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC5.

Mol	Chain	Residues	Atoms					AltConf	Trace
9	J	69	Total	C	N	O	S	0	0
			569	362	101	100	6		

- Molecule 10 is a protein called DNA-directed RNA polymerase II subunit RPB11.

Mol	Chain	Residues	Atoms					AltConf	Trace
10	K	115	Total	C	N	O	S	0	0
			924	593	157	172	2		

- Molecule 11 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC4.

Mol	Chain	Residues	Atoms					AltConf	Trace
11	L	45	Total	C	N	O	S	0	0
			359	221	71	63	4		

- Molecule 12 is a protein called DNA-directed RNA polymerase II subunit RPB1.

Mol	Chain	Residues	Atoms					AltConf	Trace
12	A	1418	Total	C	N	O	S	0	0
			11153	7029	1947	2115	62		

- Molecule 13 is a protein called DNA-directed RNA polymerase II subunit RPB2.

Mol	Chain	Residues	Atoms					AltConf	Trace
13	B	1177	Total	C	N	O	S	0	0
			9383	5932	1639	1757	55		


- Molecule 14 is a protein called DNA-directed RNA polymerase II subunit RPB4.

Mol	Chain	Residues	Atoms					AltConf	Trace
14	D	166	Total	C	N	O	S	0	0
			1332	823	238	269	2		

- Molecule 15 is a protein called DNA-directed RNA polymerases I, II, and III subunit RPABC3.

Mol	Chain	Residues	Atoms					AltConf	Trace
15	H	140	Total	C	N	O	S	0	0
			1120	704	188	224	4		

- Molecule 16 is Cisplatin (CCD ID: CPT) (formula: $\text{Cl}_2\text{H}_6\text{N}_2\text{Pt}$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			AltConf
16	T	1	Total	N	Pt	0
			3	2	1	

- Molecule 17 is ZINC ION (CCD ID: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		AltConf
17	C	1	Total	Zn	0
			1	1	
17	I	2	Total	Zn	0
			2	2	
17	J	1	Total	Zn	0
			1	1	
17	L	1	Total	Zn	0
			1	1	

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		AltConf
17	A	2	Total 2	Zn 2	0
17	B	1	Total 1	Zn 1	0


- Molecule 18 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		AltConf
18	A	1	Total 1	Mg 1	0

3 Residue-property plots


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and atom inclusion in map density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red diamond above a residue indicates a poor fit to the EM map for this residue (all-atom inclusion < 40%). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA (NTS, non-template strand)

Chain N: 


DS DA DG DA GS GS GS GS GS GS GS

- Molecule 2: RNA (5'-R(*UP*AP*CP*AP*AP*UP*GP*UP*UP*GP*GP*AP*AP*AP*GP*GP*AP*GP*AP*GP*A)-3')

Chain P: 

U A C A U A U C U G10 G11 A12 A13 A14 G15 G16 A17 G18 A19 C20 A21

- Molecule 3: DNA (TS, template strand)

Chain T: 

G1 C2 T3 C4 G5 C6 T7 C8 T9 C10 C11 T12 T13 T14 C15 C16 DC DC DT DC

- Molecule 4: DNA-directed RNA polymerase II subunit RPB3

Chain C: 


MET SER GLU E4 G5 P6 Q7 V8 K9 I10 R11 D16 D19 F20 L21 L22 V25 M29 S32 R35 V36 A39 E40 L44 S48 N54 D60 E61 P71 I77 S83 E89 D90 L101 Q102 T111 Y114 S115 L118 N140

G141 V142 L147 R148 K154 K160 A164 A174 Y180 T189 T226 T227 F228 S234 V245 I248 I258 L262 M265 D266 Q267 D268 LYS VAL ASN PHE SER GLY ASP ASN THR ALA SER ASN MET LEU GLY SER ASN GLU ASP VAL MET

THR GLY ALA GLU ASP PRO TYR SER ASN ALA SER MET GLY ASN THR GLY SER GLY TYR ASP ASN ALA TRP


- Molecule 5: DNA-directed RNA polymerases I, II, and III subunit RPABC1

Chain E: 




- Molecule 6: DNA-directed RNA polymerases I, II, and III subunit RPABC2

Chain F: 50% 6% 44%




- Molecule 7: DNA-directed RNA polymerase II subunit RPB7

Chain G: 37% 75% 25%




- Molecule 8: DNA-directed RNA polymerase II subunit RPB9

Chain I: 7% 67% 27% 5%



- Molecule 9: DNA-directed RNA polymerases I, II, and III subunit RPABC5

Chain J: 80% 19%




- Molecule 10: DNA-directed RNA polymerase II subunit RPB1

Chain K:  74% 22%




- Molecule 11: DNA-directed RNA polymerases I, II, and III subunit RPABC4

Chain L:  54% 10% 36%







- Molecule 12: DNA-directed RNA polymerase II subunit RPB1

Chain A:  59% 23% 18%






• Molecule 14: DNA-directed RNA polymerase II subunit RPB4



• Molecule 15: DNA-directed RNA polymerases I, II, and III subunit RPABC3



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	144198	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	TFS KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	50	Depositor
Minimum defocus (nm)	1200	Depositor
Maximum defocus (nm)	1800	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor
Maximum map value	0.695	Depositor
Minimum map value	-0.315	Depositor
Average map value	0.002	Depositor
Map value standard deviation	0.023	Depositor
Recommended contour level	0.09	Depositor
Map size (Å)	297.6, 297.6, 297.6	wwPDB
Map dimensions	320, 320, 320	wwPDB
Map angles (°)	90.0, 90.0, 90.0	wwPDB
Pixel spacing (Å)	0.93, 0.93, 0.93	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CPT, ZN, MG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	N	0.27	0/144	0.54	0/221
2	P	0.58	0/305	0.75	0/476
3	T	0.42	0/345	0.66	0/527
4	C	0.11	0/2124	0.27	0/2879
5	E	0.10	0/1788	0.28	0/2406
6	F	0.11	0/717	0.27	0/967
7	G	0.11	0/1367	0.30	0/1844
8	I	0.45	0/962	0.57	0/1295
9	J	0.09	0/578	0.25	0/775
10	K	0.11	0/942	0.24	0/1272
11	L	0.09	0/361	0.27	0/478
12	A	0.45	0/11352	0.61	1/15350 (0.0%)
13	B	0.14	0/9568	0.30	0/12907
14	D	0.27	0/1340	0.40	0/1797
15	H	0.09	0/1139	0.24	0/1544
All	All	0.31	0/33032	0.45	1/44738 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
12	A	0	1

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
12	A	807	GLY	CA-C-O	-5.24	118.05	122.29

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
12	A	447	GLN	Peptide

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	N	128	0	67	18	0
2	P	270	0	133	21	0
3	T	313	0	183	27	0
4	C	2086	0	2045	38	0
5	E	1752	0	1776	23	0
6	F	705	0	731	8	0
7	G	1339	0	1357	30	0
8	I	944	0	899	29	0
9	J	569	0	585	9	0
10	K	924	0	934	30	0
11	L	359	0	381	7	0
12	A	11153	0	11228	327	0
13	B	9383	0	9371	151	0
14	D	1332	0	1353	54	0
15	H	1120	0	1086	33	0
16	T	3	0	0	0	0
17	A	2	0	0	0	0
17	B	1	0	0	0	0
17	C	1	0	0	0	0
17	I	2	0	0	0	0
17	J	1	0	0	0	0
17	L	1	0	0	0	0
18	A	1	0	0	0	0
All	All	32389	0	32129	678	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

The worst 5 of 678 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
10:K:47:ARG:NH1	12:A:548:ASN:HD21	1.40	1.19
10:K:47:ARG:HH12	12:A:548:ASN:ND2	1.52	1.07
12:A:598:LEU:HD11	15:H:124:ARG:HB2	1.36	1.06
12:A:568:PRO:HD2	15:H:46:LEU:HG	1.31	1.06
10:K:47:ARG:NH1	12:A:548:ASN:ND2	2.04	1.04

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	C	263/318 (83%)	255 (97%)	8 (3%)	0	100	100
5	E	212/215 (99%)	206 (97%)	6 (3%)	0	100	100
6	F	85/155 (55%)	85 (100%)	0	0	100	100
7	G	169/171 (99%)	165 (98%)	4 (2%)	0	100	100
8	I	114/122 (93%)	109 (96%)	5 (4%)	0	100	100
9	J	67/70 (96%)	66 (98%)	1 (2%)	0	100	100
10	K	113/120 (94%)	113 (100%)	0	0	100	100
11	L	43/70 (61%)	42 (98%)	1 (2%)	0	100	100
12	A	1408/1733 (81%)	1373 (98%)	35 (2%)	0	100	100
13	B	1165/1224 (95%)	1132 (97%)	33 (3%)	0	100	100
14	D	162/221 (73%)	154 (95%)	8 (5%)	0	100	100
15	H	136/146 (93%)	134 (98%)	2 (2%)	0	100	100
All	All	3937/4565 (86%)	3834 (97%)	103 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
4	C	233/274 (85%)	233 (100%)	0	100	100
5	E	196/197 (100%)	196 (100%)	0	100	100
6	F	77/137 (56%)	77 (100%)	0	100	100
7	G	152/152 (100%)	152 (100%)	0	100	100
8	I	110/116 (95%)	109 (99%)	1 (1%)	70	88
9	J	64/65 (98%)	64 (100%)	0	100	100
10	K	99/102 (97%)	99 (100%)	0	100	100
11	L	40/57 (70%)	40 (100%)	0	100	100
12	A	1240/1520 (82%)	1238 (100%)	2 (0%)	87	96
13	B	1022/1061 (96%)	1022 (100%)	0	100	100
14	D	148/200 (74%)	145 (98%)	3 (2%)	48	77
15	H	123/128 (96%)	123 (100%)	0	100	100
All	All	3504/4009 (87%)	3498 (100%)	6 (0%)	85	96

5 of 6 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
14	D	14	ARG
14	D	16	LYS
14	D	18	VAL
12	A	445	ASN
8	I	7	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 32 such sidechains are listed below:

Mol	Chain	Res	Type
13	B	1104	HIS
15	H	11	GLN
12	A	118	HIS
12	A	92	HIS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
15	H	35	GLN

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	P	11/21 (52%)	2 (18%)	0

All (2) RNA backbone outliers are listed below:

Mol	Chain	Res	Type
2	P	12	A
2	P	21	A

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 10 ligands modelled in this entry, 9 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
16	CPT	T	101	3,1	0,2,4	-	-	-		

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.


6 Map visualisation [i](#)

This section contains visualisations of the EMDB entry EMD-65754. These allow visual inspection of the internal detail of the map and identification of artifacts.


Images derived from a raw map, generated by summing the deposited half-maps, are presented below the corresponding image components of the primary map to allow further visual inspection and comparison with those of the primary map.

6.1 Orthogonal projections [i](#)


6.1.1 Primary map



X




Y




Z


6.1.2 Raw map



X



Y




Z


The images above show the map projected in three orthogonal directions.

6.2 Central slices [i](#)


6.2.1 Primary map



X Index: 160




Y Index: 160




Z Index: 160


6.2.2 Raw map



X Index: 160



Y Index: 160




Z Index: 160


The images above show central slices of the map in three orthogonal directions.

6.3 Largest variance slices [i](#)


6.3.1 Primary map



X Index: 148




Y Index: 161




Z Index: 187


6.3.2 Raw map



X Index: 148



Y Index: 161




Z Index: 187


The images above show the largest variance slices of the map in three orthogonal directions.

6.4 Orthogonal standard-deviation projections (False-color) ⓘ


6.4.1 Primary map



X




Y




Z


6.4.2 Raw map



X



Y



Z

The images above show the map standard deviation projections with false color in three orthogonal directions. Minimum values are shown in green, max in blue, and dark to light orange shades represent small to large values respectively.


6.5 Orthogonal surface views [i](#)

6.5.1 Primary map



The images above show the 3D surface view of the map at the recommended contour level 0.09. These images, in conjunction with the slice images, may facilitate assessment of whether an appropriate contour level has been provided.

6.5.2 Raw map



These images show the 3D surface of the raw map. The raw map's contour level was selected so that its surface encloses the same volume as the primary map does at its recommended contour level.


6.6 Mask visualisation [i](#)

This section shows the 3D surface view of the primary map at 50% transparency overlaid with the specified mask at 0% transparency


A mask typically either:

- Encompasses the whole structure
- Separates out a domain, a functional unit, a monomer or an area of interest from a larger structure


6.6.1 emd_65754_msk_1.map [i](#)



X



Y




Z

7 Map analysis [i](#)


This section contains the results of statistical analysis of the map.

7.1 Map-value distribution [i](#)



The map-value distribution is plotted in 128 intervals along the x-axis. The y-axis is logarithmic. A spike in this graph at zero usually indicates that the volume has been masked.


7.2 Volume estimate [i](#)



The volume at the recommended contour level is 270 nm^3 ; this corresponds to an approximate mass of 244 kDa.

The volume estimate graph shows how the enclosed volume varies with the contour level. The recommended contour level is shown as a vertical line and the intersection between the line and the curve gives the volume of the enclosed surface at the given level.

7.3 Rotationally averaged power spectrum ⓘ




*Reported resolution corresponds to spatial frequency of 0.344 Å⁻¹

8 Fourier-Shell correlation [i](#)

Fourier-Shell Correlation (FSC) is the most commonly used method to estimate the resolution of single-particle and subtomogram-averaged maps. The shape of the curve depends on the imposed symmetry, mask and whether or not the two 3D reconstructions used were processed from a common reference. The reported resolution is shown as a black line. A curve is displayed for the half-bit criterion in addition to lines showing the 0.143 gold standard cut-off and 0.5 cut-off.

8.1 FSC [i](#)



*Reported resolution corresponds to spatial frequency of 0.344 Å⁻¹

8.2 Resolution estimates [i](#)


Resolution estimate (Å)	Estimation criterion (FSC cut-off)		
	0.143	0.5	Half-bit
Reported by author	2.91	-	-
Author-provided FSC curve	2.91	3.36	2.98
Unmasked-calculated*	3.42	4.16	3.52

*Resolution estimate based on FSC curve calculated by comparison of deposited half-maps. The value from deposited half-maps intersecting FSC 0.143 CUT-OFF 3.42 differs from the reported value 2.91 by more than 10 %


9 Map-model fit [i](#)

This section contains information regarding the fit between EMDB map EMD-65754 and PDB model 9W8W. Per-residue inclusion information can be found in section 3 on page 8.


9.1 Map-model overlay [i](#)



X




Y



Z


The images above show the 3D surface view of the map at the recommended contour level 0.09 at 50% transparency in yellow overlaid with a ribbon representation of the model coloured in blue. These images allow for the visual assessment of the quality of fit between the atomic model and the map.

9.2 Q-score mapped to coordinate model [i](#)




The images above show the model with each residue coloured according to its Q-score. This shows their resolvability in the map with higher Q-score values reflecting better resolvability. Please note: Q-score is calculating the resolvability of atoms, and thus high values are only expected at resolutions at which atoms can be resolved. Low Q-score values may therefore be expected for many entries.

9.3 Atom inclusion mapped to coordinate model [i](#)



The images above show the model with each residue coloured according to its atom inclusion. This shows to what extent they are inside the map at the recommended contour level (0.09).

9.4 Atom inclusion [i](#)



At the recommended contour level, 93% of all backbone atoms, 85% of all non-hydrogen atoms, are inside the map.

9.5 Map-model fit summary ⓘ

The table lists the average atom inclusion at the recommended contour level (0.09) and Q-score for the entire model and for each chain.

Chain	Atom inclusion	Q-score
All	0.8540	0.4720
A	0.8820	0.4900
B	0.8870	0.5020
C	0.9090	0.5100
D	0.4670	0.2040
E	0.8880	0.4430
F	0.9230	0.5330
G	0.5340	0.2930
H	0.8650	0.5110
I	0.7630	0.3900
J	0.9110	0.5210
K	0.9130	0.5260
L	0.8730	0.4540
N	0.9610	0.3210
P	0.9820	0.4890
T	0.9780	0.5120

1.0

0.0

<0.0