



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 5, 2026 – 03:53 PM UTC

PDB ID : 1W9I / pdb_00001w9i
Title : Myosin II Dictyostelium discoideum motor domain S456Y bound with MgADP-BeFx
Authors : Morris, C.A.; Coureux, P.-D.; Wells, A.L.; Houdusse, A.; Sweeney, H.L.
Deposited on : 2004-10-13
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	3183 (1.76-1.76)
Clashscore	190562	3299 (1.76-1.76)
Ramachandran outliers	187476	3274 (1.76-1.76)
Sidechain outliers	187428	3274 (1.76-1.76)
RSRZ outliers	180081	3183 (1.76-1.76)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	770	19% 89% 9%

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 6489 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called MYOSIN II HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	703	Total	C	N	O	S	0	29	1
			5663	3614	969	1063	17			

There are 13 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	760	LEU	-	expression tag	UNP P08799
A	761	ARG	-	expression tag	UNP P08799
A	762	GLY	-	expression tag	UNP P08799
A	763	ASP	-	expression tag	UNP P08799
A	764	TYR	-	expression tag	UNP P08799
A	765	LYS	-	expression tag	UNP P08799
A	766	ASP	-	expression tag	UNP P08799
A	767	ASP	-	expression tag	UNP P08799
A	768	ASP	-	expression tag	UNP P08799
A	769	ASP	-	expression tag	UNP P08799
A	770	LYS	-	expression tag	UNP P08799
A	456	TYR	SER	engineered mutation	UNP P08799
A	737	PHE	TYR	engineered mutation	UNP P08799

- Molecule 2 is MAGNESIUM ION (CCD ID: MG) (formula: Mg).


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mg	0	0
			1	1		

- Molecule 3 is ADENOSINE-5'-DIPHOSPHATE (CCD ID: ADP) (formula: C₁₀H₁₅N₅O₁₀P₂).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			27	10	5	10	2		

- Molecule 4 is BERYLLIUM TRIFLUORIDE ION (CCD ID: BEF) (formula: BeF₃).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	Be	F	0	0
			4	1	3		

- Molecule 5 is 1,2-ETHANEDIOL (CCD ID: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	734	Total	O	0	0
			734	734		

3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: MYOSIN II HEAVY CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	105.02Å 186.51Å 54.71Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	40.00 – 1.75 40.00 – 1.75	Depositor EDS
% Data completeness (in resolution range)	96.1 (40.00-1.75) 96.1 (40.00-1.75)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.65 (at 1.75Å)	Xtriage
Refinement program	REFMAC 5.1.24	Depositor
R, R_{free}	0.182 , 0.207 0.195 , 0.217	Depositor DCC
R_{free} test set	5227 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	16.5	Xtriage
Anisotropy	0.489	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 60.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	6489	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.31% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: BEF, MG, ADP, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.65	1/5887 (0.0%)	0.78	0/7948

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	754	ILE	C-N	-6.90	1.23	1.33

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	5663	0	5563	20	0
2	A	1	0	0	0	0
3	A	27	0	12	0	0
4	A	4	0	0	0	0
5	A	60	0	90	7	0
6	A	734	0	0	3	1
All	All	6489	0	5665	20	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (20) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:193[B]:ILE:HD12	1:A:245[B]:ILE:HD11	1.52	0.89
1:A:187:GLU:HB3	5:A:1763:EDO:H12	1.69	0.74
1:A:582:GLN:H	1:A:582:GLN:HE21	1.38	0.71
1:A:402:ARG:O	6:A:2418:HOH:O	2.09	0.70
1:A:637[B]:GLN:NE2	6:A:2643:HOH:O	2.24	0.68
1:A:142:ILE:CD1	5:A:1767:EDO:H21	2.29	0.62
1:A:193[A]:ILE:HD13	5:A:1764:EDO:H22	1.82	0.61
1:A:193[B]:ILE:CD1	1:A:245[B]:ILE:HD11	2.27	0.60
1:A:562[B]:ARG:HD3	6:A:2560:HOH:O	2.02	0.59
1:A:87:GLY:H	1:A:105:ASN:ND2	2.03	0.57
1:A:193[A]:ILE:HD11	1:A:243:ILE:HD13	1.87	0.55
1:A:142:ILE:HD12	5:A:1767:EDO:H21	1.90	0.52
1:A:582:GLN:HE21	1:A:582:GLN:N	2.06	0.52
1:A:142:ILE:HD11	5:A:1767:EDO:H21	1.93	0.49
1:A:641:LEU:HA	5:A:1770:EDO:H12	1.94	0.48
1:A:499:ILE:CG2	1:A:499:ILE:O	2.64	0.45
1:A:187:GLU:HB3	5:A:1763:EDO:C1	2.44	0.44
1:A:261:TYR:HE1	1:A:637[B]:GLN:HG2	1.84	0.43
1:A:688:THR:O	1:A:693:PRO:HB3	2.19	0.42
1:A:35:ILE:HD11	1:A:62:PHE:CD2	2.56	0.41

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2062:HOH:O	6:A:2720:HOH:O[1_556]	2.16	0.04

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	724/770 (94%)	710 (98%)	14 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	619/673 (92%)	616 (100%)	3 (0%)	81	75

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	443	GLN
1	A	502	THR
1	A	582	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	5	HIS
1	A	78	ASN
1	A	105	ASN
1	A	188	ASN
1	A	219	ASN
1	A	234	ASN
1	A	283	GLN
1	A	309	GLN
1	A	439	ASN
1	A	485	HIS
1	A	582	GLN
1	A	594	GLN
1	A	606	ASN
1	A	613	ASN
1	A	649	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	662	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 1 is monoatomic - leaving 17 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
5	EDO	A	1758	-	3,3,3	0.26	0	2,2,2	0.15	0
5	EDO	A	1764	-	3,3,3	0.28	0	2,2,2	0.35	0
5	EDO	A	1765	-	3,3,3	0.26	0	2,2,2	0.40	0
5	EDO	A	1761	-	3,3,3	0.26	0	2,2,2	0.46	0
5	EDO	A	1768	-	3,3,3	0.27	0	2,2,2	0.23	0
5	EDO	A	1762	-	3,3,3	0.25	0	2,2,2	0.39	0
5	EDO	A	1763	-	3,3,3	0.24	0	2,2,2	0.47	0
4	BEF	A	1757	2,3	0,3,3	-	-	-	-	-
5	EDO	A	1767	-	3,3,3	0.26	0	2,2,2	0.45	0
5	EDO	A	1766	-	3,3,3	0.26	0	2,2,2	0.39	0
5	EDO	A	1769	-	3,3,3	0.24	0	2,2,2	0.41	0
5	EDO	A	1772	-	3,3,3	0.27	0	2,2,2	0.35	0
5	EDO	A	1759	-	3,3,3	0.26	0	2,2,2	0.30	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	1760	-	3,3,3	0.26	0	2,2,2	0.36	0
3	ADP	A	1756	2,4	28,29,29	2.22	5 (17%)	43,45,45	2.02	13 (30%)
5	EDO	A	1770	-	3,3,3	0.24	0	2,2,2	0.42	0
5	EDO	A	1771	-	3,3,3	0.27	0	2,2,2	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	1758	-	-	0/1/1/1	-
5	EDO	A	1764	-	-	0/1/1/1	-
5	EDO	A	1765	-	-	0/1/1/1	-
5	EDO	A	1761	-	-	0/1/1/1	-
5	EDO	A	1768	-	-	0/1/1/1	-
5	EDO	A	1762	-	-	0/1/1/1	-
5	EDO	A	1763	-	-	0/1/1/1	-
5	EDO	A	1767	-	-	0/1/1/1	-
5	EDO	A	1769	-	-	0/1/1/1	-
5	EDO	A	1766	-	-	1/1/1/1	-
5	EDO	A	1772	-	-	0/1/1/1	-
5	EDO	A	1759	-	-	0/1/1/1	-
5	EDO	A	1760	-	-	0/1/1/1	-
3	ADP	A	1756	2,4	-	2/16/32/32	0/3/3/3
5	EDO	A	1770	-	-	0/1/1/1	-
5	EDO	A	1771	-	-	1/1/1/1	-

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	1756	ADP	C8-N7	6.74	1.44	1.31
3	A	1756	ADP	C8-N9	5.93	1.47	1.37
3	A	1756	ADP	PA-O3A	5.49	1.65	1.59
3	A	1756	ADP	C6-N6	2.58	1.40	1.34
3	A	1756	ADP	PB-O1B	2.35	1.57	1.50

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1756	ADP	N9-C8-N7	-7.15	103.80	113.94
3	A	1756	ADP	N3-C2-N1	-4.28	122.10	128.58

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1756	ADP	C5-C4-N3	-4.14	121.01	126.72
3	A	1756	ADP	C6-C5-C4	3.03	121.32	117.18
3	A	1756	ADP	C5-N7-C8	2.93	108.06	103.45
3	A	1756	ADP	C2'-C1'-N9	-2.77	106.41	113.30
3	A	1756	ADP	O3B-PB-O2B	2.48	117.10	107.80
3	A	1756	ADP	C2-N3-C4	2.33	117.51	111.83
3	A	1756	ADP	C4-N9-C8	2.26	108.11	105.74
3	A	1756	ADP	O3A-PB-O1B	-2.24	99.24	111.04
3	A	1756	ADP	N3-C4-N9	2.22	130.95	127.17
3	A	1756	ADP	C5-C4-N9	2.02	108.01	105.81
3	A	1756	ADP	C4-N9-C1'	-2.01	121.92	126.63

There are no chirality outliers.

All (4) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	1756	ADP	PA-O3A-PB-O2B
5	A	1766	EDO	O1-C1-C2-O2
5	A	1771	EDO	O1-C1-C2-O2
3	A	1756	ADP	PA-O3A-PB-O3B

There are no ring outliers.

4 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	1764	EDO	1	0
5	A	1763	EDO	2	0
5	A	1767	EDO	3	0
5	A	1770	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	703/770 (91%)	1.05	148 (21%) 2 2	9, 22, 61, 87	30 (4%)

All (148) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	752	ALA	9.0
1	A	698	TYR	8.2
1	A	746	PHE	7.9
1	A	744	ILE	7.6
1	A	399	LEU	6.8
1	A	754	ILE	6.4
1	A	741	ILE	6.3
1	A	692	PHE	6.0
1	A	696	ILE	6.0
1	A	697	ILE	5.9
1	A	749	GLY	5.9
1	A	699	ALA	5.8
1	A	624	GLY	5.6
1	A	737	PHE	5.6
1	A	751	LEU	5.6
1	A	687	ILE	5.5
1	A	693	PRO	5.3
1	A	745	PHE	5.3
1	A	738	ARG	5.3
1	A	295	ALA	5.2
1	A	743	LYS	5.0
1	A	739	PHE	4.9
1	A	495	LEU	4.9
1	A	625	ALA	4.8
1	A	499	ILE	4.7
1	A	498	LYS	4.6
1	A	740	GLY	4.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	621	ALA	4.4
1	A	21	ASP	4.4
1	A	290	ALA	4.4
1	A	742	THR	4.4
1	A	504	ILE	4.4
1	A	2	ASN	4.3
1	A	4	ILE	4.3
1	A	502	THR	4.3
1	A	400	ALA	4.3
1	A	58[A]	ASP	4.2
1	A	501	TRP	4.2
1	A	535	PHE	4.2
1	A	685	ILE	4.2
1	A	67	GLY	4.1
1	A	536	PRO	4.1
1	A	500	ASN	4.1
1	A	496	LYS	4.0
1	A	691	GLY	4.0
1	A	493	GLU	3.9
1	A	748	ALA	3.9
1	A	497	GLU	3.9
1	A	750	GLN	3.8
1	A	65[A]	VAL	3.7
1	A	626	ASN	3.7
1	A	443	GLN	3.7
1	A	72	VAL	3.7
1	A	534	VAL	3.7
1	A	506	PHE	3.6
1	A	289	THR	3.6
1	A	747	ARG	3.5
1	A	71	GLN	3.5
1	A	688	THR	3.5
1	A	627	PHE	3.5
1	A	753	ARG	3.5
1	A	694	ASN	3.4
1	A	444	GLU	3.4
1	A	59	SER	3.4
1	A	407	GLN	3.4
1	A	494	TYR	3.3
1	A	364	GLY	3.3
1	A	445	ARG	3.3
1	A	40	ASP	3.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	623	LYS	3.1
1	A	52	ILE	3.1
1	A	147	ARG	3.1
1	A	695	ARG	3.0
1	A	683	GLU	3.0
1	A	202	ARG	3.0
1	A	288	ALA	3.0
1	A	5	HIS	3.0
1	A	388	VAL	2.9
1	A	292	GLU	2.9
1	A	537	ASN	2.9
1	A	736	GLN	2.9
1	A	509	ASP	2.9
1	A	503	PHE	2.8
1	A	755	GLU	2.8
1	A	83[A]	ILE	2.8
1	A	8	THR	2.8
1	A	20	GLY	2.8
1	A	57	SER	2.8
1	A	298	LEU	2.8
1	A	32	LYS	2.8
1	A	75	ASP	2.8
1	A	686	ARG	2.7
1	A	3	PRO	2.7
1	A	380	THR	2.7
1	A	689	ARG	2.7
1	A	398	ILE	2.7
1	A	61	THR	2.6
1	A	252	PHE	2.6
1	A	291	GLU	2.6
1	A	41	PRO	2.5
1	A	406	ALA	2.5
1	A	684	GLY	2.5
1	A	405	VAL	2.5
1	A	293	LYS	2.4
1	A	362	GLY	2.4
1	A	46	SER	2.3
1	A	47	TYR	2.3
1	A	700	ASP	2.3
1	A	73	LYS	2.3
1	A	622	LYS	2.3
1	A	690	LYS	2.3

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	404	LEU	2.3
1	A	606	ASN	2.3
1	A	33	ARG	2.3
1	A	66[A]	ASP	2.3
1	A	69	ASP	2.3
1	A	294	LYS	2.3
1	A	200	ALA	2.3
1	A	43	GLU	2.3
1	A	96	TYR	2.3
1	A	93	GLU	2.2
1	A	150[A]	GLU	2.2
1	A	77	ALA	2.2
1	A	97	LEU	2.2
1	A	322	GLU	2.2
1	A	145	GLY	2.2
1	A	60	PHE	2.2
1	A	42	LYS	2.2
1	A	35	ILE	2.2
1	A	615	PRO	2.2
1	A	53	VAL	2.1
1	A	401	GLY	2.1
1	A	369	LEU	2.1
1	A	371	ASP	2.1
1	A	487	PHE	2.1
1	A	581	ILE	2.1
1	A	440	VAL	2.1
1	A	505	ASP	2.1
1	A	34	TYR	2.1
1	A	37	TYR	2.1
1	A	74	LYS	2.1
1	A	574	ALA	2.1
1	A	678[A]	CYS	2.1
1	A	6	ASP	2.1
1	A	619	SER	2.1
1	A	489	LEU	2.1
1	A	44	ARG	2.0
1	A	367	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

6.4 Ligands ⓘ




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	EDO	A	1763	4/4	0.75	0.19	47,47,48,49	0
5	EDO	A	1771	4/4	0.75	0.24	56,56,57,57	0
5	EDO	A	1770	4/4	0.81	0.23	40,40,40,41	0
5	EDO	A	1766	4/4	0.83	0.21	45,45,45,45	0
5	EDO	A	1768	4/4	0.84	0.20	27,31,34,36	0
5	EDO	A	1769	4/4	0.85	0.14	31,31,32,34	0
5	EDO	A	1765	4/4	0.88	0.15	22,26,28,31	0
5	EDO	A	1772	4/4	0.88	0.14	54,54,55,55	0
5	EDO	A	1762	4/4	0.89	0.12	35,36,36,38	0
5	EDO	A	1767	4/4	0.90	0.11	32,35,35,37	0
5	EDO	A	1758	4/4	0.91	0.12	25,27,27,29	0
5	EDO	A	1759	4/4	0.91	0.19	31,31,32,33	0
5	EDO	A	1761	4/4	0.94	0.10	34,35,35,37	0
5	EDO	A	1764	4/4	0.94	0.08	25,26,26,28	0
5	EDO	A	1760	4/4	0.96	0.07	28,28,30,30	0
3	ADP	A	1756	27/27	0.98	0.05	9,12,15,15	0
4	BEF	A	1757	4/4	0.99	0.04	10,11,11,12	0
2	MG	A	1755	1/1	1.00	0.01	10,10,10,10	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around ADP A 1756:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.