



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 7, 2026 – 02:16 AM UTC

PDB ID : 2W9D / pdb_00002w9d
Title : Structure of Fab fragment of the ICSM 18 - anti-Prp therapeutic antibody at 1.57 Å resolution.
Authors : Antonyuk, S.V.; Trevitt, C.R.; Strange, R.W.; Jackson, G.S.; Sangar, D.; Batchelor, M.; Jones, S.; Georgiou, T.; Cooper, S.; Fraser, C.; Khalili-Shirazi, A.; Clarke, A.R.; Hasnain, S.S.; Collinge, J.
Deposited on : 2009-01-23
Resolution : 1.57 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.57 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	1094 (1.58-1.58)
Clashscore	190562	1105 (1.58-1.58)
Ramachandran outliers	187476	1082 (1.58-1.58)
Sidechain outliers	187428	1081 (1.58-1.58)
RSRZ outliers	180081	1094 (1.58-1.58)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	H	215	14% 86% 11%
2	L	212	29% 87% 12%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4067 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called ICSM 18-ANTI-PRP THERAPEUTIC FAB HEAVY CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	H	215	Total	C	N	O	S	3	17	0
			1701	1087	269	335	10			

- Molecule 2 is a protein called ICSM 18-ANTI-PRP THERAPEUTIC FAB LIGHT CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	L	211	Total	C	N	O	S	0	16	0
			1698	1068	277	342	11			

- Molecule 3 is CALCIUM ION (CCD ID: CA) (formula: Ca).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	H	1	Total	Ca	0	0
			1	1		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	H	344	Total	O	0	0
			344	344		
4	L	323	Total	O	0	0
			323	323		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: ICSM 18-ANTI-PRP THERAPEUTIC FAB HEAVY CHAIN



- Molecule 2: ICSM 18-ANTI-PRP THERAPEUTIC FAB LIGHT CHAIN



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	36.78Å 84.71Å 127.85Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.00 – 1.57 27.00 – 1.57	Depositor EDS
% Data completeness (in resolution range)	94.9 (27.00-1.57) 95.0 (27.00-1.57)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 1.57Å)	Xtriage
Refinement program	REFMAC 5.3.0037	Depositor
R, R_{free}	0.184 , 0.217 0.191 , 0.222	Depositor DCC
R_{free} test set	2725 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	15.9	Xtriage
Anisotropy	0.523	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 34.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	4067	wwPDB-VP
Average B, all atoms (Å ²)	25.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.35% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	H	0.93	3/1797 (0.2%)	0.69	4/2458 (0.2%)
2	L	0.39	0/1782	0.60	0/2419
All	All	0.71	3/3579 (0.1%)	0.64	4/4877 (0.1%)

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	H	135	THR	CA-CB	32.75	1.97	1.53
1	H	52	TYR	C-N	9.21	1.46	1.34
1	H	57	VAL	C-N	-6.98	1.24	1.33

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	H	135	THR	N-CA-CB	8.78	126.67	110.87
1	H	52	TYR	CA-C-N	5.48	125.13	119.05
1	H	52	TYR	C-N-CA	5.48	125.13	119.05
1	H	56	GLY	N-CA-C	-5.11	101.02	110.73

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	H	1701	0	1700	26	0
2	L	1698	0	1658	29	0
3	H	1	0	0	0	0
4	H	344	0	0	5	0
4	L	323	0	0	2	0
All	All	4067	0	3358	51	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:12[B]:VAL:CG1	1:H:16[B]:SER:OG	1.96	1.13
2:L:47:ILE:HD12	2:L:72[A]:LEU:HD13	1.37	1.05
2:L:31[B]:TYR:HE2	2:L:33[B]:HIS:CE1	1.77	1.01
1:H:12[B]:VAL:HG12	1:H:16[B]:SER:OG	1.65	0.94
2:L:31[B]:TYR:CE2	2:L:33[B]:HIS:CE1	2.56	0.93
2:L:31[B]:TYR:HE2	2:L:33[B]:HIS:HE1	1.20	0.86
2:L:47:ILE:HD12	2:L:72[A]:LEU:CD1	2.12	0.79
2:L:31[B]:TYR:CE2	2:L:33[B]:HIS:HE1	2.03	0.73
2:L:47:ILE:CD1	2:L:72[A]:LEU:CD1	2.69	0.71
1:H:98[A]:LEU:HD12	1:H:99:TYR:N	2.09	0.67
2:L:47:ILE:CD1	2:L:72[A]:LEU:HD13	2.18	0.66
1:H:171[A]:VAL:CG2	2:L:159[A]:LEU:HD13	2.29	0.62
1:H:171[B]:VAL:HB	2:L:159[B]:LEU:HD11	1.80	0.61
2:L:204[A]:ILE:HD11	4:L:2311:HOH:O	1.99	0.61
2:L:89:GLN:HE22	2:L:92:SER:H	1.49	0.60
1:H:12[B]:VAL:HG12	1:H:16[B]:SER:HG	1.64	0.59
1:H:12[B]:VAL:HG13	1:H:16[B]:SER:OG	1.98	0.57
1:H:32:TYR:CD1	1:H:98[A]:LEU:HD13	2.40	0.55
1:H:12[B]:VAL:HG11	1:H:16[B]:SER:OG	1.98	0.55
1:H:32:TYR:HD1	1:H:98[A]:LEU:HD13	1.72	0.54
2:L:88:HIS:HD2	2:L:96:THR:O	1.89	0.54
2:L:88:HIS:CE1	2:L:90:TRP:HE1	2.25	0.54
4:H:2293:HOH:O	2:L:159[A]:LEU:HD11	2.11	0.51
2:L:80[A]:GLU:HG3	4:L:2287:HOH:O	2.11	0.50
1:H:171[A]:VAL:HG23	2:L:159[A]:LEU:HD13	1.93	0.49
2:L:31[B]:TYR:CD2	2:L:33[B]:HIS:CE1	3.01	0.49
1:H:185:VAL:HG12	4:H:2305:HOH:O	2.13	0.47
1:H:34:LEU:HD21	1:H:98[B]:LEU:HD12	1.97	0.47
2:L:11[B]:MET:HG3	2:L:103:LEU:HD12	1.96	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:H:6:GLN:HE21	1:H:107:GLY:HA3	1.80	0.47
1:H:60:TYR:HD1	4:H:2112:HOH:O	1.97	0.47
2:L:47:ILE:HD11	2:L:72[A]:LEU:CD1	2.45	0.47
1:H:171[A]:VAL:HG21	2:L:159[A]:LEU:HD13	1.96	0.46
1:H:131:SER:O	1:H:132:ALA:HB2	2.13	0.46
1:H:12[B]:VAL:HG21	1:H:18:VAL:HB	1.99	0.45
2:L:89:GLN:NE2	2:L:91:ARG:H	2.16	0.44
2:L:11[B]:MET:HG3	2:L:103:LEU:CD1	2.47	0.44
1:H:179:LEU:C	1:H:179:LEU:HD12	2.43	0.43
2:L:135:LEU:N	2:L:135:LEU:HD12	2.34	0.43
2:L:149:ILE:CD1	2:L:180:LEU:HD11	2.49	0.43
1:H:153:THR:HG23	4:H:2123:HOH:O	2.17	0.43
2:L:18:LYS:HE3	2:L:73:THR:HG21	2.01	0.42
1:H:31[A]:ASP:OD1	4:H:2076:HOH:O	2.21	0.42
1:H:52:TYR:HB2	1:H:56:GLY:HA3	2.01	0.42
1:H:12[B]:VAL:CG1	1:H:16[B]:SER:HG	2.17	0.42
2:L:210:ARG:O	2:L:211:ASN:ND2	2.54	0.41
2:L:211:ASN:HD22	2:L:211:ASN:HA	1.67	0.41
2:L:160:ASN:ND2	2:L:176:SER:OG	2.54	0.41
1:H:137:MET:HE3	1:H:184:THR:HG22	2.03	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	H	230/215 (107%)	220 (96%)	6 (3%)	4 (2%)	7	0
2	L	225/212 (106%)	222 (99%)	3 (1%)	0	100	100
All	All	455/427 (107%)	442 (97%)	9 (2%)	4 (1%)	14	3

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	H	55	ASN
1	H	133	ALA
1	H	132	ALA
1	H	134	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	H	205/189 (108%)	197 (96%)	8 (4%)	28	4
2	L	200/185 (108%)	198 (99%)	2 (1%)	68	49
All	All	405/374 (108%)	395 (98%)	10 (2%)	43	12

All (10) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	H	23	LYS
1	H	74	LYS
1	H	75	SER
1	H	86	LEU
1	H	98[A]	LEU
1	H	98[B]	LEU
1	H	135	THR
1	H	179	LEU
2	L	89	GLN
2	L	184	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (8) such sidechains are listed below:

Mol	Chain	Res	Type
1	H	6	GLN
1	H	193	GLN
2	L	88	HIS
2	L	89	GLN
2	L	93	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	L	160	ASN
2	L	189	ASN
2	L	211	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

1 non-standard protein/DNA/RNA residue is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PCA	L	1	2	7,8,9	1.87	1 (14%)	9,10,12	1.96	4 (44%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	L	1	2	-	0/0/11/13	0/1/1/1

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	L	1	PCA	CD-N	4.79	1.46	1.34

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	1	PCA	CB-CA-N	2.75	110.81	103.24

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	L	1	PCA	CA-N-CD	-2.71	104.30	113.58
2	L	1	PCA	CG-CD-N	2.48	114.45	108.39
2	L	1	PCA	OE-CD-CG	-2.32	122.58	126.72

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	H	215/215 (100%)	1.35	30 (13%)	6 10	13, 23, 31, 38	26 (12%)
2	L	210/212 (99%)	1.57	62 (29%)	1 1	13, 23, 28, 35	19 (9%)
All	All	425/427 (99%)	1.46	92 (21%)	2 4	13, 23, 29, 38	45 (10%)

All (92) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	H	133	ALA	8.2
1	H	131	SER	6.6
1	H	132	ALA	5.6
1	H	56	GLY	5.5
2	L	139	TYR	5.3
2	L	211	ASN	5.3
1	H	57	VAL	5.3
1	H	130	GLY	5.2
2	L	124	LEU	5.0
2	L	172	TYR	4.8
2	L	108	ALA	4.8
2	L	156	ASN	4.5
2	L	180	LEU	4.4
1	H	74	LYS	4.3
2	L	127	GLY	4.3
1	H	134	GLN	4.2
2	L	183	ASP	4.1
2	L	171	THR	3.9
2	L	170	SER	3.8
2	L	128	GLY	3.8
1	H	60	TYR	3.7
2	L	155	GLN	3.7
2	L	140	PRO	3.6
2	L	138	PHE	3.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	149	ILE	3.5
1	H	41	HIS	3.5
1	H	77	SER	3.5
1	H	52	TYR	3.4
2	L	125	THR	3.4
1	H	55	ASN	3.2
2	L	122	GLU	3.2
2	L	167	SER	3.2
2	L	159[A]	LEU	3.1
2	L	151	GLY	3.1
2	L	111	ALA	3.1
2	L	110	ALA	3.0
2	L	181	THR	3.0
2	L	152	SER	3.0
2	L	179	THR	3.0
1	H	31[A]	ASP	3.0
2	L	137	ASN	2.9
2	L	126	SER	2.9
1	H	135	THR	2.9
2	L	41	THR	2.9
2	L	143	ILE	2.9
1	H	54	ASN	2.9
2	L	163	THR	2.8
2	L	199	THR	2.8
1	H	53	PRO	2.8
2	L	202	SER	2.8
2	L	150	ASP	2.7
2	L	200	SER	2.7
2	L	165	GLN	2.7
2	L	189	ASN	2.6
1	H	26	ARG	2.6
2	L	107	ARG	2.6
1	H	28	THR	2.6
2	L	135	LEU	2.6
2	L	10	ILE	2.5
2	L	169	ASP	2.5
2	L	164	ASP	2.5
2	L	201	THR	2.5
2	L	168	LYS	2.4
2	L	210	ARG	2.4
1	H	213	THR	2.4
1	H	93	VAL	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	L	178	LEU	2.4
2	L	141	LYS	2.3
1	H	171[A]	VAL	2.3
2	L	109	ASP	2.3
2	L	187	ARG	2.3
2	L	154	ARG	2.3
1	H	136	SER	2.3
2	L	121	SER	2.3
1	H	75	SER	2.2
2	L	166	ASP	2.2
2	L	182	LYS	2.2
2	L	173	SER	2.2
1	H	2	VAL	2.2
2	L	204[A]	ILE	2.1
1	H	187	SER	2.1
2	L	158	VAL	2.1
1	H	86	LEU	2.1
2	L	2	ILE	2.1
2	L	157	GLY	2.1
1	H	12[A]	VAL	2.0
1	H	72	VAL	2.0
2	L	23	CYS	2.0
2	L	184	GLU	2.0
2	L	31[A]	TYR	2.0
1	H	44	THR	2.0
2	L	207	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
2	PCA	L	1	8/9	0.88	0.10	25,27,28,29	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
3	CA	H	251	1/1	0.94	0.17	25,25,25,25	1

6.5 Other polymers [i](#)

There are no such residues in this entry.