



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2026 – 01:23 AM UTC

PDB ID : 3W9G / pdb_00003w9g
Title : Crystal structure of the ankyrin repeat domain of chicken TRPV4
Authors : Itoh, Y.; Hamada-nakahara, S.; Suetsugu, S.
Deposited on : 2013-04-04
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	10052 (2.00-2.00)
Clashscore	190562	11152 (2.00-2.00)
Ramachandran outliers	187476	11031 (2.00-2.00)
Sidechain outliers	187428	11029 (2.00-2.00)
RSRZ outliers	180081	10067 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	260	 4% 88% 10% .
1	B	260	 5% 91% 7% .
1	C	260	 3% 88% 9% .
1	D	260	 25% 80% 17% ..

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 8741 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Vanilloid receptor-related osmotically activated channel protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	256	Total	C	N	O	S	0	6	0
			2065	1307	382	367	9			
1	B	256	Total	C	N	O	S	0	5	0
			2058	1302	379	368	9			
1	C	252	Total	C	N	O	S	0	3	0
			2009	1273	366	361	9			
1	D	252	Total	C	N	O	S	0	0	0
			2001	1267	366	360	8			

There are 40 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	132	MET	-	expression tag	UNP Q9DFS3
A	383	ALA	-	expression tag	UNP Q9DFS3
A	384	ALA	-	expression tag	UNP Q9DFS3
A	385	ALA	-	expression tag	UNP Q9DFS3
A	386	HIS	-	expression tag	UNP Q9DFS3
A	387	HIS	-	expression tag	UNP Q9DFS3
A	388	HIS	-	expression tag	UNP Q9DFS3
A	389	HIS	-	expression tag	UNP Q9DFS3
A	390	HIS	-	expression tag	UNP Q9DFS3
A	391	HIS	-	expression tag	UNP Q9DFS3
B	132	MET	-	expression tag	UNP Q9DFS3
B	383	ALA	-	expression tag	UNP Q9DFS3
B	384	ALA	-	expression tag	UNP Q9DFS3
B	385	ALA	-	expression tag	UNP Q9DFS3
B	386	HIS	-	expression tag	UNP Q9DFS3
B	387	HIS	-	expression tag	UNP Q9DFS3
B	388	HIS	-	expression tag	UNP Q9DFS3
B	389	HIS	-	expression tag	UNP Q9DFS3
B	390	HIS	-	expression tag	UNP Q9DFS3
B	391	HIS	-	expression tag	UNP Q9DFS3
C	132	MET	-	expression tag	UNP Q9DFS3

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	383	ALA	-	expression tag	UNP Q9DFS3
C	384	ALA	-	expression tag	UNP Q9DFS3
C	385	ALA	-	expression tag	UNP Q9DFS3
C	386	HIS	-	expression tag	UNP Q9DFS3
C	387	HIS	-	expression tag	UNP Q9DFS3
C	388	HIS	-	expression tag	UNP Q9DFS3
C	389	HIS	-	expression tag	UNP Q9DFS3
C	390	HIS	-	expression tag	UNP Q9DFS3
C	391	HIS	-	expression tag	UNP Q9DFS3
D	132	MET	-	expression tag	UNP Q9DFS3
D	383	ALA	-	expression tag	UNP Q9DFS3
D	384	ALA	-	expression tag	UNP Q9DFS3
D	385	ALA	-	expression tag	UNP Q9DFS3
D	386	HIS	-	expression tag	UNP Q9DFS3
D	387	HIS	-	expression tag	UNP Q9DFS3
D	388	HIS	-	expression tag	UNP Q9DFS3
D	389	HIS	-	expression tag	UNP Q9DFS3
D	390	HIS	-	expression tag	UNP Q9DFS3
D	391	HIS	-	expression tag	UNP Q9DFS3


- Molecule 2 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	205	Total O 205 205	0	0
2	B	232	Total O 232 232	0	0
2	C	132	Total O 132 132	0	0
2	D	39	Total O 39 39	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Vanilloid receptor-related osmotically activated channel protein




- Molecule 1: Vanilloid receptor-related osmotically activated channel protein




- Molecule 1: Vanilloid receptor-related osmotically activated channel protein



- Molecule 1: Vanilloid receptor-related osmotically activated channel protein





4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	104.95Å 48.27Å 134.03Å 90.00° 101.79° 90.00°	Depositor
Resolution (Å)	37.57 – 2.00 37.57 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (37.57-2.00) 99.7 (37.57-2.00)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.51 (at 2.00Å)	Xtriage
Refinement program	PHENIX	Depositor
R, R_{free}	0.179 , 0.211 (Not available) , 0.207	Depositor DCC
R_{free} test set	1821 reflections (2.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.5	Xtriage
Anisotropy	0.444	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 51.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	8741	wwPDB-VP
Average B, all atoms (Å ²)	48.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 48.88 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 8.0735e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.55	0/2138	0.78	0/2882
1	B	0.56	0/2125	0.80	2/2866 (0.1%)
1	C	0.43	0/2065	0.79	1/2786 (0.0%)
1	D	0.40	0/2041	0.88	5/2754 (0.2%)
All	All	0.49	0/8369	0.81	8/11288 (0.1%)

There are no bond length outliers.

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	260	GLN	CA-C-N	5.59	125.96	119.47
1	B	260	GLN	C-N-CA	5.59	125.96	119.47
1	D	215	SER	CA-C-N	5.51	125.98	120.14
1	D	215	SER	C-N-CA	5.51	125.98	120.14
1	D	165	ARG	N-CA-C	5.16	117.02	109.24
1	C	209	MET	N-CA-C	5.05	116.47	111.07
1	D	148	SER	CA-C-N	5.03	124.63	119.05
1	D	148	SER	C-N-CA	5.03	124.63	119.05

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2065	0	2095	12	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2058	0	2086	10	0
1	C	2009	0	2023	14	0
1	D	2001	0	2016	23	0
2	A	205	0	0	1	0
2	B	232	0	0	2	0
2	C	132	0	0	0	0
2	D	39	0	0	0	0
All	All	8741	0	8220	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:237:LYS:NZ	1:A:241:GLU:OE2	2.26	0.69
1:D:164:LYS:NZ	1:D:170:GLU:OE2	2.25	0.67
1:B:237:LYS:NZ	1:B:241:GLU:OE2	2.30	0.64
1:A:300:LEU:HD11	1:A:338:LYS:HG2	1.81	0.61
1:C:237:LYS:NZ	1:C:241:GLU:OE2	2.34	0.60
1:D:275:LEU:HD21	1:D:300:LEU:HD23	1.84	0.59
1:D:213:ILE:HG23	1:D:243:LEU:HD22	1.86	0.58
1:B:292:GLU:OE1	1:B:338:LYS:NZ	2.36	0.58
1:D:140:LEU:HA	1:D:143:ILE:HD12	1.87	0.57
1:B:300:LEU:HD11	1:B:338:LYS:HG2	1.86	0.56
1:D:136:ASN:HA	1:D:171:PHE:HE1	1.71	0.56
1:C:258:PHE:H	1:C:258:PHE:HD2	1.53	0.55
1:D:165:ARG:NH1	1:D:206:THR:O	2.41	0.54
1:D:165:ARG:NH2	1:D:206:THR:O	2.43	0.52
1:A:292:GLU:OE1	1:A:338:LYS:NZ	2.44	0.51
1:B:152:LEU:HD11	1:B:195:THR:HB	1.93	0.51
1:C:300:LEU:HD11	1:C:338:LYS:HG2	1.92	0.51
1:B:360:LEU:HG	1:B:361:MET:HE2	1.93	0.50
1:D:165:ARG:NH1	1:D:208:ASN:HB2	2.27	0.50
1:A:245:GLU:HG2	1:A:295[A]:HIS:NE2	2.27	0.50
1:D:214:ASN:OD1	1:D:249:ASP:N	2.35	0.49
1:C:144:VAL:HG13	1:C:184:ALA:HB2	1.93	0.49
1:D:293:ASN:OD1	1:D:296:LYS:N	2.27	0.49
1:A:360:LEU:HG	1:A:361:MET:HE2	1.95	0.48
1:D:172:ARG:NH2	1:D:177:GLY:O	2.43	0.47
1:B:262:LYS:NZ	2:B:591:HOH:O	2.47	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:489:HOH:O	1:C:387:HIS:HD2	1.98	0.47
1:C:245:GLU:HG2	1:C:295:HIS:NE2	2.29	0.46
1:D:229:HIS:CE1	1:D:254:ALA:HB2	2.50	0.46
1:B:164:LYS:NZ	2:B:533:HOH:O	2.49	0.46
1:C:234:ARG:HA	1:C:234:ARG:HD3	1.74	0.46
1:C:258:PHE:CD2	1:C:258:PHE:N	2.84	0.46
1:C:364:ALA:HB2	1:C:372:PHE:CE1	2.52	0.45
1:A:145:SER:HB3	1:A:183:LYS:HE3	1.99	0.44
1:D:264:GLU:HG2	1:D:266:GLY:H	1.82	0.44
1:B:181:LEU:HB3	1:B:182:PRO:HD3	2.00	0.43
1:D:173:GLU:HA	1:D:174:PRO:HD3	1.85	0.43
1:D:196:ILE:HB	1:D:197:PRO:HD3	2.01	0.43
1:A:152:LEU:HD11	1:A:195:THR:HB	2.00	0.42
1:C:258:PHE:HD2	1:C:258:PHE:N	2.14	0.42
1:A:204:GLU:HA	1:A:209:MET:HB2	2.01	0.42
1:D:181:LEU:HB3	1:D:182:PRO:HD3	2.00	0.42
1:D:252:ALA:O	1:D:274:PRO:HD3	2.20	0.42
1:A:150:ASP:O	1:A:153:GLU:HG3	2.19	0.42
1:B:378[B]:ARG:NH1	1:B:379:GLU:OE2	2.47	0.42
1:D:137:ARG:HB3	1:D:138:PRO:HD3	2.00	0.42
1:D:134:VAL:HA	1:D:158:PHE:CE1	2.55	0.42
1:D:148:SER:HA	1:D:149:PRO:HD2	1.85	0.42
1:D:219:ASP:O	1:D:257:ARG:HD3	2.20	0.41
1:D:195:THR:O	1:D:199:LEU:HG	2.20	0.41
1:D:193:ASN:O	1:D:196:ILE:HG12	2.20	0.41
1:C:181:LEU:HB3	1:C:182:PRO:HD3	2.02	0.41
1:A:217:PHE:CZ	1:A:230:ILE:HD11	2.56	0.41
1:A:375:ILE:O	1:A:379:GLU:HG3	2.21	0.41
1:A:181:LEU:HB3	1:A:182:PRO:HD3	2.03	0.41
1:C:137:ARG:HB2	1:C:170:GLU:O	2.21	0.41
1:C:272:GLU:OE1	1:C:304:ASP:HB2	2.20	0.40
1:B:296:LYS:HE3	1:B:296:LYS:HB2	1.92	0.40
1:C:193:ASN:O	1:C:196:ILE:HG13	2.22	0.40

There are no symmetry-related clashes.

5.3 Torsion angles

5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	260/260 (100%)	254 (98%)	6 (2%)	0	100	100
1	B	259/260 (100%)	253 (98%)	6 (2%)	0	100	100
1	C	251/260 (96%)	245 (98%)	6 (2%)	0	100	100
1	D	248/260 (95%)	241 (97%)	7 (3%)	0	100	100
All	All	1018/1040 (98%)	993 (98%)	25 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	223/221 (101%)	219 (98%)	4 (2%)	51	58
1	B	222/221 (100%)	218 (98%)	4 (2%)	51	58
1	C	216/221 (98%)	213 (99%)	3 (1%)	59	66
1	D	213/221 (96%)	211 (99%)	2 (1%)	70	78
All	All	874/884 (99%)	861 (98%)	13 (2%)	61	64

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	317	ILE
1	A	327	PHE

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	378[A]	ARG
1	A	378[B]	ARG
1	B	317	ILE
1	B	327	PHE
1	B	378[A]	ARG
1	B	378[B]	ARG
1	C	258	PHE
1	C	317	ILE
1	C	327	PHE
1	D	220	VAL
1	D	327	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	285	HIS
1	B	285	HIS
1	D	347	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no oligosaccharides in this entry.

5.6 Ligand geometry ⓘ

There are no ligands in this entry.

5.7 Other polymers ⓘ

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2		OWAB(Å ²)	Q<0.9
1	A	256/260 (98%)	-0.32	11 (4%)	40 39	15, 27, 68, 105	6 (2%)
1	B	256/260 (98%)	-0.30	12 (4%)	36 35	15, 27, 70, 142	5 (1%)
1	C	252/260 (96%)	0.09	8 (3%)	50 49	29, 43, 65, 119	3 (1%)
1	D	252/260 (96%)	1.43	66 (26%)	1 1	37, 84, 125, 186	0
All	All	1016/1040 (97%)	0.22	97 (9%)	14 12	15, 41, 108, 186	14 (1%)

All (97) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	266	GLY	5.4
1	B	387	HIS	4.9
1	C	258	PHE	4.9
1	D	258	PHE	4.6
1	D	134	VAL	4.5
1	D	265	GLY	4.4
1	D	171	PHE	4.3
1	B	265	GLY	4.3
1	D	389	HIS	4.2
1	A	261	PRO	4.2
1	D	159	LEU	4.2
1	D	207	GLY	3.9
1	D	185	LEU	3.9
1	D	226	THR	3.9
1	C	388	HIS	3.9
1	D	135	PHE	3.8
1	B	260	GLN	3.7
1	D	139	ILE	3.7
1	B	384	ALA	3.5
1	A	387	HIS	3.4
1	D	203	ALA	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	247	GLY	3.2
1	D	266	GLY	3.2
1	D	206	THR	3.2
1	B	261	PRO	3.2
1	D	267	TYR	3.2
1	D	161	THR	3.2
1	C	259	PHE	3.1
1	D	145	SER	3.0
1	D	220	VAL	3.0
1	A	260	GLN	2.9
1	D	210	ARG	2.9
1	D	163	LYS	2.9
1	D	246	LYS	2.9
1	D	160	LEU	2.9
1	A	258	PHE	2.9
1	D	175	SER	2.9
1	D	138	PRO	2.8
1	D	212	PHE	2.8
1	C	267	TYR	2.8
1	D	216	PRO	2.8
1	D	164	LYS	2.8
1	B	258	PHE	2.8
1	C	133	LYS	2.8
1	C	265	GLY	2.8
1	D	174	PRO	2.8
1	D	215	SER	2.7
1	A	266	GLY	2.7
1	A	259	PHE	2.7
1	B	266	GLY	2.7
1	D	166	LEU	2.6
1	A	385	ALA	2.6
1	D	178	LYS	2.6
1	D	300	LEU	2.6
1	D	238	HIS	2.5
1	B	259	PHE	2.5
1	D	183	LYS	2.5
1	D	388	HIS	2.5
1	D	232	ILE	2.5
1	D	237	LYS	2.5
1	D	191	GLY	2.5
1	D	256	GLY	2.5
1	D	213	ILE	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	385	ALA	2.4
1	D	133	LYS	2.4
1	D	155	LEU	2.4
1	D	136	ASN	2.4
1	C	264	GLU	2.4
1	B	386	HIS	2.4
1	D	369	ILE	2.4
1	D	264	GLU	2.3
1	D	348	LEU	2.3
1	A	132	MET	2.3
1	A	262	LYS	2.3
1	D	244	VAL	2.3
1	A	267	TYR	2.3
1	D	230	ILE	2.3
1	B	132	MET	2.3
1	D	292	GLU	2.2
1	D	351	LEU	2.2
1	D	143	ILE	2.1
1	D	198	ILE	2.1
1	D	342	LEU	2.1
1	B	262	LYS	2.1
1	D	285	HIS	2.1
1	D	227	ALA	2.1
1	D	293	ASN	2.1
1	D	236	CYS	2.1
1	D	208	ASN	2.1
1	D	229	HIS	2.1
1	D	162	HIS	2.0
1	D	295	HIS	2.0
1	A	330	LYS	2.0
1	D	153	GLU	2.0
1	D	157	SER	2.0
1	D	248	ALA	2.0
1	D	146	ARG	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.