



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 14, 2026 – 02:23 PM UTC

PDB ID : 4W9H / pdb_00004w9h
Title : pVHL:EloB:EloC in complex with (2S,4R)-1-((S)-2-acetamido-3,3-dimethylbutanoyl)-4-hydroxy-N-(4-(4-methylthiazol-5-yl)benzyl)pyrrolidine-2-carboxamide (ligand 7)
Authors : Gadd, M.S.; Soares, P.; Galdeano, C.; van Molle, I.; Ciulli, A.
Deposited on : 2014-08-27
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity	:	4-5-2 with Phenix2.0
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	2.0
EDS	:	3.0
Buster-report	:	wwPDB partial adaption of 1.1.7 (2018)
Percentile statistics	:	20250101.v01 (using entries in the PDB archive January 1st 2025)
CCP4	:	9.0.010 (Gargrove)
Density-Fitness	:	1.0.12
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.49


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	180053	6658 (2.10-2.10)
Clashscore	190562	7164 (2.10-2.10)
Ramachandran outliers	187476	7099 (2.10-2.10)
Sidechain outliers	187428	7100 (2.10-2.10)
RSRZ outliers	180081	6662 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	104	2% 96% ..
1	D	104	4% 98% .
1	G	104	2% 91% 7% ..
1	J	104	7% 93% 5% ..

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	B	97	
2	E	97	
3	C	162	
3	F	162	
3	I	162	
3	L	162	
4	H	97	
4	K	97	

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 11271 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Transcription elongation factor B polypeptide 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	103	Total	As	C	N	O	S	0	0	0
			809	2	514	136	152	5			
1	D	102	Total	As	C	N	O	S	0	0	0
			794	2	506	132	149	5			
1	G	103	Total	As	C	N	O	S	0	0	0
			816	2	517	136	156	5			
1	J	103	Total	As	C	N	O	S	0	0	0
			812	2	515	136	154	5			

- Molecule 2 is a protein called Transcription elongation factor B polypeptide 1.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
2	B	86	Total	As	C	N	O	S	0	0	0
			685	1	442	109	127	6			
2	E	88	Total	As	C	N	O	S	0	0	0
			692	1	446	111	128	6			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	16	MET	-	initiating methionine	UNP Q15369
E	16	MET	-	initiating methionine	UNP Q15369

- Molecule 3 is a protein called Von Hippel-Lindau disease tumor suppressor.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
3	C	141	Total	As	C	N	O	S	0	1	0
			1134	1	723	203	205	2			
3	F	144	Total	As	C	N	O	S	0	0	0
			1139	1	727	202	207	2			
3	I	143	Total	As	C	N	O	S	0	0	0
			1150	1	733	209	205	2			

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
3	L	143	Total	As	C	N	O	S	0	0	0
			1149	1	733	209	204	2			

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
C	52	GLY	-	expression tag	UNP P40337
C	53	SER	-	expression tag	UNP P40337
F	52	GLY	-	expression tag	UNP P40337
F	53	SER	-	expression tag	UNP P40337
I	52	GLY	-	expression tag	UNP P40337
I	53	SER	-	expression tag	UNP P40337
L	52	GLY	-	expression tag	UNP P40337
L	53	SER	-	expression tag	UNP P40337


- Molecule 4 is a protein called Transcription elongation factor B polypeptide 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
4	H	88	Total	C	N	O	S	0	0	0
			690	446	112	126	6			
4	K	88	Total	C	N	O	S	0	0	0
			694	449	112	126	7			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
H	16	MET	-	initiating methionine	UNP Q15369
K	16	MET	-	initiating methionine	UNP Q15369

- Molecule 5 is N-acetyl-3-methyl-L-valyl-(4R)-4-hydroxy-N-[4-(4-methyl-1,3-thiazol-5-yl)benzyl]-L-prolinamide (CCD ID: 3JF) (formula: C₂₄H₃₂N₄O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
5	C	1	Total	C	N	O	S	0	0
			33	24	4	4	1		
5	F	1	Total	C	N	O	S	0	0
			33	24	4	4	1		
5	I	1	Total	C	N	O	S	0	0
			33	24	4	4	1		
5	L	1	Total	C	N	O	S	0	0
			33	24	4	4	1		

- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	61	Total	O	0	0
			61	61		
6	B	32	Total	O	0	0
			32	32		
6	C	55	Total	O	0	0
			55	55		
6	D	40	Total	O	0	0
			40	40		
6	E	28	Total	O	0	0
			28	28		
6	F	44	Total	O	0	0
			44	44		
6	G	51	Total	O	0	0
			51	51		
6	H	41	Total	O	0	0
			41	41		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	I	60	Total 60	O 60	0	0
6	J	76	Total 76	O 76	0	0
6	K	37	Total 37	O 37	0	0
6	L	50	Total 50	O 50	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Transcription elongation factor B polypeptide 2





- Molecule 1: Transcription elongation factor B polypeptide 2





- Molecule 1: Transcription elongation factor B polypeptide 2





- Molecule 1: Transcription elongation factor B polypeptide 2




- Molecule 2: Transcription elongation factor B polypeptide 1




- Molecule 2: Transcription elongation factor B polypeptide 1




- Molecule 3: Von Hippel-Lindau disease tumor suppressor




- Molecule 3: Von Hippel-Lindau disease tumor suppressor





- Molecule 3: Von Hippel-Lindau disease tumor suppressor





- Molecule 3: Von Hippel-Lindau disease tumor suppressor



- Molecule 4: Transcription elongation factor B polypeptide 1



- Molecule 4: Transcription elongation factor B polypeptide 1



4 Data and refinement statistics

Property	Value	Source
Space group	P 41 2 2	Depositor
Cell constants a, b, c, α , β , γ	92.98Å 92.98Å 364.37Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	93.10 – 2.10 92.98 – 2.10	Depositor EDS
% Data completeness (in resolution range)	99.9 (93.10-2.10) 99.9 (92.98-2.10)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.10	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.13 (at 2.10Å)	Xtrriage
Refinement program	REFMAC 5.8.0073	Depositor
R, R_{free}	0.210 , 0.253 0.217 , 0.259	Depositor DCC
R_{free} test set	4784 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å ²)	33.3	Xtrriage
Anisotropy	0.262	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 41.7	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	11271	wwPDB-VP
Average B, all atoms (Å ²)	41.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 29.79 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.4604e-03. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: CAS, 3JF

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/805	0.81	2/1086 (0.2%)
1	D	0.49	0/789	0.70	0/1063
1	G	0.53	0/812	0.76	0/1095
1	J	0.56	0/808	0.76	0/1090
2	B	0.54	0/689	0.75	0/932
2	E	0.53	0/696	0.77	0/941
3	C	0.53	0/1156	0.73	0/1579
3	F	0.53	0/1159	0.74	0/1587
3	I	0.57	0/1170	0.79	0/1598
3	L	0.57	0/1169	0.82	0/1597
4	H	0.58	0/704	0.82	1/949 (0.1%)
4	K	0.62	0/708	0.83	1/955 (0.1%)
All	All	0.55	0/10665	0.77	4/14472 (0.0%)

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	K	112	CYS	CA-CB-SG	6.66	129.72	114.40
1	A	91	GLU	CA-C-N	-5.66	114.43	120.03
1	A	91	GLU	C-N-CA	-5.66	114.43	120.03
4	H	110	LEU	N-CA-C	5.33	122.14	110.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	809	0	800	1	0
1	D	794	0	779	0	0
1	G	816	0	806	6	0
1	J	812	0	802	4	0
2	B	685	0	672	4	0
2	E	692	0	672	6	0
3	C	1134	0	1114	7	0
3	F	1139	0	1097	9	0
3	I	1150	0	1130	6	0
3	L	1149	0	1127	7	0
4	H	690	0	683	10	0
4	K	694	0	689	6	0
5	C	33	0	32	0	0
5	F	33	0	32	0	0
5	I	33	0	32	0	0
5	L	33	0	32	0	0
6	A	61	0	0	0	0
6	B	32	0	0	2	0
6	C	55	0	0	1	0
6	D	40	0	0	0	0
6	E	28	0	0	1	0
6	F	44	0	0	0	0
6	G	51	0	0	0	0
6	H	41	0	0	1	0
6	I	60	0	0	1	0
6	J	76	0	0	0	0
6	K	37	0	0	0	0
6	L	50	0	0	0	0
All	All	11271	0	10499	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:73:VAL:HG13	2:B:112:CAS:CE2	2.09	0.83
3:C:73:GLN:OE1	3:C:108:ARG:NH1	2.23	0.72
4:H:105:MET:HE2	3:I:180:ILE:HG22	1.76	0.68
3:C:108:ARG:HD2	6:C:451:HOH:O	1.94	0.67
1:G:37:ARG:NH1	1:G:41:GLU:OE1	2.30	0.64
2:E:17:MET:HA	2:E:17:MET:HE2	1.80	0.63
4:H:109:PHE:O	4:H:111:ASP:N	2.32	0.63
4:H:38:THR:HG23	4:H:80:LYS:HD3	1.83	0.60
4:K:45:MET:HE1	4:K:109:PHE:CZ	2.40	0.56
1:G:3:VAL:HG13	1:G:67:ALA:CB	2.34	0.56
2:E:102:GLU:HA	2:E:105:MET:HE3	1.88	0.56
3:L:182:ARG:HA	3:L:185:TYR:CE2	2.41	0.56
1:G:3:VAL:HG13	1:G:67:ALA:HB3	1.89	0.54
2:E:77:PHE:HE1	2:E:112:CAS:CE2	2.20	0.54
3:F:107:ARG:HD3	4:H:109:PHE:CE1	2.43	0.53
3:F:180:ILE:HD12	3:F:184:LEU:HB3	1.90	0.52
3:C:106:GLY:C	4:K:109:PHE:CE1	2.88	0.52
3:L:202:THR:O	3:L:204:GLU:N	2.42	0.52
3:F:73:GLN:H	3:F:141:ASN:HD21	1.58	0.51
2:B:89:GLU:HA	6:B:223:HOH:O	2.10	0.51
3:L:180:ILE:HD12	3:L:184:LEU:HB2	1.92	0.51
3:I:73:GLN:HG2	3:I:108:ARG:NH2	2.26	0.51
1:J:1:MET:HB3	1:J:64:SER:OG	2.11	0.51
4:H:35:HIS:O	4:H:38:THR:HB	2.11	0.50
3:F:70:GLU:CD	3:F:113:ARG:HE	2.19	0.49
1:G:102:VAL:O	3:I:170:VAL:HG23	2.13	0.49
1:J:11:LYS:HG3	1:J:91:GLU:HG3	1.96	0.48
3:F:118:LEU:HD13	3:F:120:ARG:HD3	1.96	0.47
4:K:38:THR:HG23	4:K:80:LYS:HD3	1.96	0.47
1:G:80:ARG:HA	1:G:85:PHE:HA	1.95	0.47
2:E:98:GLU:HB2	6:E:216:HOH:O	2.15	0.47
4:H:101:LEU:HD22	3:I:180:ILE:HG21	1.96	0.47
2:E:86:SER:O	3:F:132:GLN:NE2	2.48	0.46
3:I:182:ARG:CZ	3:I:182:ARG:HB3	2.45	0.46
1:J:52:ASP:OD1	1:J:52:ASP:N	2.49	0.46
4:K:38:THR:HG23	4:K:80:LYS:CD	2.46	0.46
2:B:98:GLU:HB2	6:B:222:HOH:O	2.16	0.45
4:H:87:SER:HA	6:I:454:HOH:O	2.17	0.45
3:L:170:VAL:CG1	3:L:174:ASN:HB2	2.47	0.45
3:F:79:ARG:NE	3:F:150:ASN:OD1	2.50	0.45
3:C:154:PRO:HG2	3:C:156:TYR:CE1	2.52	0.45
3:F:109:ILE:C	3:F:109:ILE:HD12	2.42	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:H:38:THR:CG2	4:H:80:LYS:HD3	2.46	0.44
1:A:80:ARG:HA	1:A:85:PHE:HA	1.99	0.44
1:G:3:VAL:HG12	1:G:5:LEU:HD21	1.99	0.44
2:E:77:PHE:HE1	2:E:112:CAS:AS	2.61	0.44
3:C:191:HIS:O	3:C:192:PRO:C	2.59	0.43
3:F:73:GLN:OE1	3:F:108:ARG:NH2	2.51	0.43
3:L:90:ASN:OD1	3:L:90:ASN:C	2.62	0.42
2:B:73:VAL:CG1	2:B:112:CAS:CE2	2.89	0.42
4:H:63:ARG:NH2	6:H:210:HOH:O	2.54	0.41
4:K:41:THR:CG2	4:K:45:MET:HE3	2.50	0.41
1:J:52:ASP:HB2	1:J:55:LYS:HG2	2.03	0.41
4:K:110:LEU:HB2	4:K:112:CYS:SG	2.61	0.41
3:L:118:LEU:HD13	3:L:120:ARG:HD3	2.04	0.41
4:H:101:LEU:HD22	3:I:180:ILE:CG2	2.52	0.40
3:L:182:ARG:HA	3:L:185:TYR:CD2	2.56	0.40
3:C:112:TYR:HB2	3:C:115:HIS:CE1	2.55	0.40
3:C:181:VAL:O	3:C:182:ARG:C	2.64	0.40

There are no symmetry-related clashes.

5.3 Torsion angles ⓘ

5.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	99/104 (95%)	96 (97%)	3 (3%)	0	100	100
1	D	96/104 (92%)	95 (99%)	1 (1%)	0	100	100
1	G	99/104 (95%)	96 (97%)	3 (3%)	0	100	100
1	J	99/104 (95%)	94 (95%)	5 (5%)	0	100	100
2	B	82/97 (84%)	80 (98%)	2 (2%)	0	100	100
2	E	84/97 (87%)	82 (98%)	2 (2%)	0	100	100
3	C	139/162 (86%)	134 (96%)	5 (4%)	0	100	100

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
3	F	141/162 (87%)	138 (98%)	3 (2%)	0	100	100
3	I	140/162 (86%)	138 (99%)	2 (1%)	0	100	100
3	L	140/162 (86%)	135 (96%)	4 (3%)	1 (1%)	18	15
4	H	84/97 (87%)	78 (93%)	5 (6%)	1 (1%)	10	7
4	K	84/97 (87%)	82 (98%)	1 (1%)	1 (1%)	10	7
All	All	1287/1452 (89%)	1248 (97%)	36 (3%)	3 (0%)	43	44

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
4	H	110	LEU
4	K	85	ASN
3	L	203	GLN

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	86/90 (96%)	86 (100%)	0	100	100
1	D	83/90 (92%)	83 (100%)	0	100	100
1	G	88/90 (98%)	87 (99%)	1 (1%)	65	74
1	J	87/90 (97%)	86 (99%)	1 (1%)	65	74
2	B	75/85 (88%)	74 (99%)	1 (1%)	61	69
2	E	74/85 (87%)	73 (99%)	1 (1%)	59	67
3	C	126/147 (86%)	124 (98%)	2 (2%)	55	64
3	F	124/147 (84%)	124 (100%)	0	100	100
3	I	127/147 (86%)	125 (98%)	2 (2%)	55	64
3	L	126/147 (86%)	124 (98%)	2 (2%)	55	64
4	H	75/86 (87%)	72 (96%)	3 (4%)	28	29
4	K	76/86 (88%)	72 (95%)	4 (5%)	20	19

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	1147/1290 (89%)	1130 (98%)	17 (2%)	57 65

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	101	LEU
3	C	150	ASN
3	C	196	LYS
2	E	58	ASN
1	G	37	ARG
4	H	17	MET
4	H	38	THR
4	H	112	CYS
3	I	73	GLN
3	I	148	PHE
1	J	52	ASP
4	K	39	SER
4	K	61	ASN
4	K	85	ASN
4	K	112	CYS
3	L	148	PHE
3	L	195	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	61	ASN
3	C	174	ASN
2	E	35	HIS
2	E	61	ASN
3	F	96	GLN
3	F	141	ASN
3	F	174	ASN
3	F	191	HIS
3	I	96	GLN
3	I	174	ASN
3	L	174	ASN

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

14 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
1	CAS	J	89	1	5,8,9	1.25	0	1,9,11	1.66	0
3	CAS	L	77	3	5,8,9	0.62	0	1,9,11	0.22	0
3	CAS	C	77	3	5,8,9	0.55	0	1,9,11	0.46	0
1	CAS	G	60	1	5,8,9	0.83	0	1,9,11	0.56	0
1	CAS	D	60	1	5,8,9	0.64	0	1,9,11	0.61	0
1	CAS	J	60	1	5,8,9	0.66	0	1,9,11	0.45	0
2	CAS	E	112	2	6,9,9	1.48	1 (16%)	2,11,11	1.63	0
1	CAS	G	89	1	5,8,9	1.15	0	1,9,11	1.97	0
1	CAS	A	89	1	5,8,9	1.20	0	1,9,11	0.16	0
2	CAS	B	112	2	6,9,9	1.24	1 (16%)	2,11,11	2.05	1 (50%)
3	CAS	I	77	3	5,8,9	0.67	0	1,9,11	0.44	0
3	CAS	F	77	3	5,8,9	0.62	0	1,9,11	0.30	0
1	CAS	D	89	1	5,8,9	1.22	0	1,9,11	0.78	0
1	CAS	A	60	1	5,8,9	0.85	0	1,9,11	0.32	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	J	89	1	-	0/0/7/9	-
3	CAS	L	77	3	-	0/0/7/9	-
3	CAS	C	77	3	-	0/0/7/9	-
1	CAS	G	60	1	-	0/0/7/9	-
1	CAS	D	60	1	-	0/0/7/9	-

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CAS	J	60	1	-	0/0/7/9	-
2	CAS	E	112	2	-	0/6/9/9	-
1	CAS	G	89	1	-	0/0/7/9	-
1	CAS	A	89	1	-	0/0/7/9	-
2	CAS	B	112	2	-	0/6/9/9	-
3	CAS	I	77	3	-	0/0/7/9	-
3	CAS	F	77	3	-	0/0/7/9	-
1	CAS	D	89	1	-	0/0/7/9	-
1	CAS	A	60	1	-	0/0/7/9	-

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	112	CAS	OXT-C	-2.40	1.23	1.30
2	E	112	CAS	OXT-C	-2.39	1.23	1.30

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	112	CAS	OXT-C-O	-2.21	119.06	124.08

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	112	CAS	2	0
2	B	112	CAS	2	0

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	3JF	C	301	-	34,35,35	1.93	6 (17%)	49,51,51	2.75	9 (18%)
5	3JF	L	301	-	34,35,35	1.86	5 (14%)	49,51,51	3.17	10 (20%)
5	3JF	I	301	-	34,35,35	1.89	5 (14%)	49,51,51	2.98	8 (16%)
5	3JF	F	301	-	34,35,35	1.86	7 (20%)	49,51,51	2.48	7 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	3JF	C	301	-	-	0/31/43/43	0/3/3/3
5	3JF	L	301	-	-	0/31/43/43	0/3/3/3
5	3JF	I	301	-	-	0/31/43/43	0/3/3/3
5	3JF	F	301	-	-	0/31/43/43	0/3/3/3

All (23) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	3JF	CBA-CBB	-6.00	1.35	1.48
5	F	301	3JF	CBA-CBB	-5.79	1.36	1.48
5	L	301	3JF	CBA-CBB	-5.47	1.36	1.48
5	I	301	3JF	CA-C	-5.42	1.40	1.52
5	I	301	3JF	CBA-CBB	-5.41	1.37	1.48
5	L	301	3JF	CA-C	-5.19	1.40	1.52
5	F	301	3JF	CA-C	-4.87	1.41	1.52
5	C	301	3JF	CA-C	-4.78	1.41	1.52
5	I	301	3JF	CBB-SAU	-4.21	1.62	1.73
5	L	301	3JF	CA-N	3.67	1.54	1.47
5	C	301	3JF	CAO-CAY	-3.62	1.43	1.51
5	I	301	3JF	CAO-CAY	-3.51	1.43	1.51
5	L	301	3JF	CAO-CAY	-3.50	1.43	1.51
5	I	301	3JF	CA-N	3.32	1.53	1.47
5	F	301	3JF	CA-N	3.31	1.53	1.47
5	F	301	3JF	CAO-CAY	-3.24	1.44	1.51
5	C	301	3JF	CA-N	3.12	1.53	1.47

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	C	301	3JF	CAN-SAU	-3.07	1.66	1.72
5	L	301	3JF	CAN-SAU	-2.89	1.67	1.72
5	F	301	3JF	CD2-N	2.33	1.50	1.47
5	C	301	3JF	CAZ-NAR	-2.29	1.34	1.38
5	F	301	3JF	CAB-CAZ	2.18	1.54	1.49
5	F	301	3JF	CAZ-NAR	-2.09	1.35	1.38

All (34) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	I	301	3JF	CAN-SAU-CBB	15.02	96.82	88.96
5	L	301	3JF	CAN-SAU-CBB	14.61	96.61	88.96
5	C	301	3JF	CAN-SAU-CBB	12.48	95.49	88.96
5	I	301	3JF	SAU-CAN-NAR	-10.77	106.83	116.08
5	F	301	3JF	CAN-SAU-CBB	10.51	94.46	88.96
5	L	301	3JF	SAU-CAN-NAR	-10.08	107.42	116.08
5	C	301	3JF	SAU-CAN-NAR	-9.41	107.99	116.08
5	F	301	3JF	SAU-CAN-NAR	-8.93	108.40	116.08
5	L	301	3JF	CAZ-CBB-SAU	-8.60	104.37	109.64
5	C	301	3JF	CAZ-CBB-SAU	-5.53	106.25	109.64
5	F	301	3JF	CAZ-CBB-SAU	-5.45	106.30	109.64
5	I	301	3JF	CAZ-CBB-SAU	-4.98	106.59	109.64
5	C	301	3JF	CBA-CBB-SAU	4.56	127.25	118.77
5	F	301	3JF	CAN-NAR-CAZ	4.39	115.65	110.71
5	L	301	3JF	CBA-CBB-SAU	4.34	126.84	118.77
5	L	301	3JF	CAN-NAR-CAZ	3.90	115.10	110.71
5	C	301	3JF	CAN-NAR-CAZ	3.73	114.91	110.71
5	F	301	3JF	CBA-CBB-SAU	3.37	125.03	118.77
5	C	301	3JF	CBA-CBB-CAZ	-3.18	126.50	130.81
5	I	301	3JF	CG-CD2-N	2.91	105.80	103.00
5	I	301	3JF	CBA-CBB-SAU	2.87	124.11	118.77
5	C	301	3JF	CD2-N-CA	-2.68	108.28	111.83
5	F	301	3JF	CG-CB-CA	2.67	106.87	103.75
5	L	301	3JF	CAJ-CAL-CBA	2.57	123.54	120.80
5	L	301	3JF	CAX-CBE-NAT	-2.35	104.54	107.40
5	L	301	3JF	CAL-CBA-CAM	-2.35	115.58	118.57
5	C	301	3JF	CG-CB-CA	2.23	106.35	103.75
5	C	301	3JF	CAX-CBE-NAT	-2.21	104.72	107.40
5	I	301	3JF	CAY-CAO-NAS	-2.16	108.53	113.07
5	L	301	3JF	CG-CB-CA	2.12	106.22	103.75
5	I	301	3JF	CAN-NAR-CAZ	2.12	113.09	110.71
5	I	301	3JF	CG-CB-CA	2.09	106.19	103.75

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	301	3JF	CAX-CBE-NAT	-2.04	104.93	107.40
5	L	301	3JF	CAY-CAO-NAS	-2.03	108.80	113.07


There are no chirality outliers.



There are no torsion outliers.


There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	101/104 (97%)	-0.03	2 (1%) 65 67	26, 35, 59, 85	0
1	D	100/104 (96%)	0.24	4 (4%) 42 44	27, 45, 69, 84	0
1	G	101/104 (97%)	0.16	2 (1%) 65 67	27, 41, 61, 71	0
1	J	101/104 (97%)	-0.08	7 (6%) 23 24	22, 31, 56, 86	0
2	B	85/97 (87%)	0.30	5 (5%) 28 30	26, 37, 62, 107	0
2	E	87/97 (89%)	0.32	5 (5%) 29 31	27, 40, 70, 84	0
3	C	140/162 (86%)	0.32	6 (4%) 40 41	25, 37, 67, 87	1 (0%)
3	F	143/162 (88%)	0.20	6 (4%) 40 42	23, 37, 69, 99	0
3	I	142/162 (87%)	0.26	4 (2%) 55 57	24, 37, 58, 69	0
3	L	142/162 (87%)	0.47	20 (14%) 6 6	23, 36, 77, 88	0
4	H	88/97 (90%)	0.54	8 (9%) 15 15	27, 39, 73, 84	0
4	K	88/97 (90%)	0.38	11 (12%) 8 8	23, 34, 65, 79	0
All	All	1318/1452 (90%)	0.26	80 (6%) 27 29	22, 38, 69, 107	1 (0%)

All (80) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
4	H	110	LEU	7.7
4	H	109	PHE	6.8
4	H	111	ASP	6.7
3	C	62	VAL	5.0
1	D	81	ALA	4.8
4	K	57	THR	4.8
1	J	102	VAL	4.6
3	F	62	VAL	4.6
4	K	110	LEU	4.5
3	L	170	VAL	4.5
3	L	142	VAL	4.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	83	ASP	4.4
3	L	178	LEU	4.2
4	K	109	PHE	4.2
3	F	185	TYR	4.1
3	L	177	ARG	4.0
3	L	62	VAL	3.9
2	E	57	THR	3.9
3	L	181	VAL	3.8
3	L	175	TYR	3.8
3	I	170	VAL	3.7
2	E	88	THR	3.6
2	E	48	GLY	3.5
3	I	62	VAL	3.5
3	L	184	LEU	3.3
2	B	87	SER	3.3
1	J	103	MET	3.2
3	L	179	ASP	3.1
4	H	44	ALA	3.1
3	C	202	THR	3.1
1	G	83	ASP	3.1
2	B	88	THR	3.0
1	J	101	ASP	3.0
3	C	191	HIS	3.0
1	A	83	ASP	2.9
3	L	176	ARG	2.9
3	L	172	PRO	2.8
3	F	201	LEU	2.8
4	K	45	MET	2.8
2	B	47	SER	2.8
3	L	173	GLU	2.8
2	B	17	MET	2.7
3	F	191	HIS	2.7
4	H	17	MET	2.7
1	J	99	LEU	2.7
4	K	46	LEU	2.7
4	H	112	CYS	2.7
4	H	57	THR	2.6
3	I	181	VAL	2.6
3	L	171	LYS	2.6
3	L	185	TYR	2.6
4	K	85	ASN	2.6
1	J	84	THR	2.6

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
3	L	180	ILE	2.5
3	L	144	GLY	2.5
4	K	88	THR	2.5
3	L	204	GLU	2.4
3	C	201	LEU	2.3
2	E	47	SER	2.3
2	E	87	SER	2.3
1	G	81	ALA	2.3
1	D	84	THR	2.3
4	K	44	ALA	2.3
3	F	205	ARG	2.3
3	L	195	GLN	2.3
3	C	144	GLY	2.3
4	K	87	SER	2.2
1	J	100	PRO	2.2
4	H	42	ILE	2.2
3	C	176	ARG	2.2
1	J	83	ASP	2.2
1	A	101	ASP	2.2
3	F	202	THR	2.2
1	D	102	VAL	2.1
3	I	185	TYR	2.1
4	K	107	ALA	2.1
2	B	85	ASN	2.1
4	K	47	SER	2.1
3	L	191	HIS	2.1
3	L	174	ASN	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
1	CAS	D	89	9/10	0.88	0.16	51,58,92,94	0
2	CAS	E	112	10/10	0.89	0.19	35,42,90,92	0
1	CAS	G	89	9/10	0.89	0.17	32,37,74,78	0
2	CAS	B	112	10/10	0.91	0.20	47,54,88,96	0
1	CAS	A	89	9/10	0.93	0.14	40,46,75,75	0
1	CAS	A	60	9/10	0.95	0.12	29,32,53,53	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
1	CAS	D	60	9/10	0.96	0.10	31,33,55,58	0
1	CAS	J	60	9/10	0.96	0.11	28,32,52,57	0
1	CAS	G	60	9/10	0.98	0.09	37,39,50,52	0
3	CAS	C	77	9/10	0.98	0.08	31,34,54,56	0
3	CAS	I	77	9/10	0.98	0.09	26,29,43,45	0
3	CAS	F	77	9/10	0.98	0.07	27,29,49,50	0
1	CAS	J	89	9/10	0.98	0.08	28,31,52,57	0
3	CAS	L	77	9/10	0.98	0.09	26,28,44,48	0

6.3 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors($\text{\AA}^2$)	Q<0.9
5	3JF	C	301	33/33	0.95	0.07	24,29,40,40	0
5	3JF	I	301	33/33	0.95	0.08	24,30,36,38	0
5	3JF	L	301	33/33	0.95	0.08	23,29,37,39	0
5	3JF	F	301	33/33	0.96	0.07	24,31,36,37	0




The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.

Electron density around 3JF C 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)




**Electron density around 3JF I 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)






Electron density around 3JF L 301:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)

**Electron density around 3JF F 301:**

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.